View Article Online / Journal Homepage / Table of Contents for this issue

## Stereospecific Cyclopropane Ring Hydrogenolysis

By M. N. AKHTAR and W. R. JACKSON\*

(Department of Chemistry, David Keir Building, The Queen's University of Belfast, Belfast BT9 5AG)

Summary Hydrogenolysis of the cyclopropane ring in 3-acetoxynortricyclene has been shown to be stereo-specifically cis.

THE ring opening of cyclopropanes by hydrogen over a metal catalyst is of wide application<sup>1</sup> and of particular interest to us since it is a possible step in the rearrangement of cycloalkanes over metal surfaces.<sup>2</sup> The one example in which the stereochemistry of hydrogenolysis of a cyclo-

propane ring has been established showed it to be non-stereoselective<sup>3</sup> over a platinum catalyst in acetic acid solution.

We have investigated the hydrogenolysis of 3-acetoxynortricyclene (1) over a platinum catalyst in acetic acid solution containing a small amount of perchloric acid at  $80^{\circ}$ . A mixture (32:68) of the 2-exo-(2) and 7-acetoxy-(3) norbornanes was obtained; no 2-endo-acetoxynorbornane was detected (by g.l.c. or n.m.r.). A sample of the endo-acetate was shown to be isomerised only slowly to the exo-isomer (ca. 5%) under the reaction conditions, indicating that the isomer ratio is almost exclusively kinetically controlled. The experiment was repeated using deuterium and the mixture of deuteriated (2) and (3) so obtained was separated by preparative g.l.c.

At 220 MHz the n.m.r. spectrum of (3) showed the following absorptions:  $\tau$  5.12 (s, 7-H), 7.86 (s, 1- and 4-H), 8.19 (d, syn-exo-H), 8.30 (s, COMe), 8.62 (d, anti-exo-H), 8.85 (d, syn-endo-H), and 9.06 (d, anti-endo-H) with  $^{2}J_{exo-endo}$  7.5 Hz. Assignments were made by examination of the n.m.r. spectra of specifically labelled deuterio-7acetoxynorbornanes. syn-exo-Dideuterio-(3) was prepared by reduction  $(LiAlD_4^5)$  of syn-norbornen-7-ol<sup>4</sup> and subsequent acetylation. The absorption at  $\tau$  8.19 had disappeared and was assigned to the syn-exo-protons while the doublet at  $\tau$  8.85 which had collapsed to a singlet was assigned to the syn-endo-protons. anti-exo-Dideuterio-(3) was prepared by addition of deuterium over a palladium catalyst to 7-anti-acetoxynorbornene<sup>5</sup> and its spectrum showed no absorption at  $\tau$  8.62 and a singlet at  $\tau$  9.06. Further, addition of deuterium to 7-acetoxynorbornadiene over a platinum catalyst gave a compound which showed singlets at  $\tau$  8.19 and 9.06 with smaller singlets at  $\tau$  8.62 and 8.85 suggesting that exo-addition to the anti-olefinic bond and endo-addition to the syn-olefinic bond were predominant, in agreement with previous suggestions.<sup>6</sup> Integration of the spectrum of deuteriated (3) obtained by deuteriolysis of (1) now showed only single protons for the syn- and antiendo-absorptions.† Thus hydrogenolysis of the cyclopropane ring in 7-acetoxynortricyclene involves predominant (>90%) cis-hydrogen addition from the outer edge of the cyclopropane ring. This result thus parallels the stereochemistry of hydrogenation of many olefins1 and provides an interesting contrast with the previous, nonspecific, reaction.<sup>3</sup>



It was not possible to elucidate fully the stereochemistry of deuterium incorporation into the 2-exo-acetate (2) because of the complexity of its n.m.r. spectrum. However, the absorption due to the 2-endo-proton in the deuteriated compound was modified in a manner consistent with loss of the large  ${}^{3}J_{endo-endo}$  coupling.<sup>7</sup> This observation is in agreement with the formation of 3,5-di-endo-deuterio-2acetoxynorbornane, as would be predicted for a stereospecific deuterium incorporation similar to that observed in (3).

## (Received, 5th May 1972; Com. 766.)

† The evidence does not exclude the formation of 2,5-di-endo-deuterio-7-acetoxynorbornane. However, formation of this isomer must involve a highly improbable stereospecific rearrangement and incorporation of deuterium is almost certainly into the 2,6-endo positions.

<sup>1</sup> M. Freifelder, 'Practical Catalytic Hydrogenation', Wiley-Interscience, New York, 1971, p. 525; P. N. Rylander, 'Catalytic Hydrogenation Over Platinum Metals', Academic Press, New York, 1967, p. 468.
<sup>2</sup> H. A. Quinn, M. A. McKervey, W. R. Jackson, and J. J. Rooney, J. Amer. Chem. Soc., 1970, 92, 2922.
<sup>3</sup> Z. Majerski and P. von R. Schleyer, Tetrahedron Letters, 1968, 6195.
<sup>4</sup> W. C. Baird, J. Org. Chem., 1966, 31, 2411.
<sup>5</sup> B. Franzus and E. S. Snyder, J. Amer. Chem. Soc., 1965, 87, 3423.
<sup>6</sup> H. C. Brown and J. H. Kawakami, J. Amer. Chem. Soc., 1970, 92, 201.
<sup>7</sup> J. I. Musher, Mol. Phys., 1962, 6, 93.