Contents lists available at SciVerse ScienceDirect



Inorganic Chemistry Communications



journal homepage: www.elsevier.com/locate/inoche

Synthesis and characterization of new soluble polyfluorinated seco-porphyrazines

Selma Kayaköy, Ergün Gonca*

Department of Chemistry, Fatih University, TR34500 B.Cekmece, Istanbul, Turkey

ARTICLE INFO

Article history: Received 15 February 2012 Accepted 9 April 2012 Available online 19 April 2012

Keywords: Seco-porphyrazines Fluorocarbons Pentafluorophenyl Zinc Copper

ABSTRACT

The synthesis of new polyfluorinated magnesium porphyrazine bearing pentafluorophenyl moieties was achieved by cyclotetramerization of 3,4-bis(pentafluorophenyl)pyrroline-2,5-diimine in the presence of Mg(BuO)₂. Acidmediated demetallation of the magnesium porphyrazine resulted in peripheral oxidation of one pyrrole ring to reveal the *seco*-porphyrazine, [2,3,7,8,12,13,17,18-octakis(pentafluorophenyl)-2-*seco*-2,3-dioxoporphyrazine]. Further reaction of that product with copper (II) acetate, zinc (II) acetate and cobalt (II) acetate has led to the metallated *seco*-derivatives, [2,3,7,8,12,13,17,18-octakis(pentafluorophenyl)-2-*seco*-2,3-dioxoporphyrazinato] M(II) [M = Cu(II), Zn(II), Co(II)]. The spectroscopic and aggregation behavior of all the target porphyrazines were investigated in different solvents and at different concentrations in chloroform. The compounds were characterized by using many spectroscopic techniques such as FT-IR, ¹H NMR, ¹³C NMR, ¹⁹F NMR, UV-vis, mass and elemental analysis. The porphyrazines were soluble in various organic solvents such as chloroform, dichloromethane, pyridine, and THF.

© 2012 Elsevier B.V. All rights reserved.

Porphyrins (P) and their structural derivatives –phthalocyanines (Pc) and porphyrazines (Pz)– constitute a distinct class of macrocyclic tetrapyrrole systems with unique physical and chemical properties. Phthalocyanines have received a great deal of interest owing to their extensive use in a number of applications involving chemical sensors, electrochromic devices, batteries, semiconductors, catalysts, and liquid crystals [1,2]. Metallated phthalocyanines (MPcs) are also currently undergoing clinical trials for use in the photodynamic therapy (PDT) of cancer [3–8]. Because of their chemical diversity, these compounds, and in particular porphyrins and phthalocyanines, have been at the focus of multidisciplinary interest for many years. However, although the molecular structures of all three types of macrocycles are very similar, benzo- and especially aza-substitution have a strong impact on the overall chemical behavior of the corresponding tetrapyrrole ligands [9–11].

The porphyrazines have specific optical properties because they have symmetrical rich 18π -electron aromatic macrocycle, which can play as a host to different metal ions in its central cavity. Therefore, they have been extensively used in optical recording media, photosensors, optical filters and copy preventing inks. A *seco*-porphyrazine is defined as a porphyrazine in which one of the pyrrole rings has been effectively cut and replaced by two acyclic substituents attached to the macrocyclic core.

Fluorinated metallated tetrapyrrole compounds currently receive a great deal of attention because of their interesting electron transfer, photosensitizing properties, along with magnetic and thermal characteristics [12–17]. The presence of pentafluorophenyl groups on the macrocycle ring can increase the catalytic activity and stability [18]. Fluoro-substituted tetrapyrrole compounds are known for their high solubility, even in polar, aprotic solvents. The increased solubility may be due to fluorine, which has the highest electronegativity of all elements [19].

Our group has been heavily interested in the preparation of new soluble porphyrazine derivatives. Among these, Pzs with peripheral functional groups such as 3,5-bis(trifluoromethyl)benzylthio [20] and 3-thiopropylpentafluorobenzoate [21] can be cited. Recently, we have synthesized novel *seco*-porphyrazines substituted with 1-naphthyl [22], *o*-tolyl and *p*-tolyl [23], 4-tert-butylphenyl [24], 4-biphenyl [25], and 3,5-bis(trifluoromethyl)phenyl [26] groups on the peripheral positions as encountered by Barrett, Hoffman and coworkers, with peripheral amino derivatives [27–30].

In the present paper, octasubstituted symmetric magnesium porphyrazine (**3**), unsymmetrical metal-free *seco*-porphyrazine (**4**), and metallated *seco*-porphyrazines (**5–7**) containing polyfluorophenyl groups on the periphery were synthesized. Magnesium porphyrazine was prepared by a direct cyclotetramerization of 3,4-bis(pentafluorophenyl)pyrroline-2,5-diimine (**2**) in the presence of Mg(BuO)₂. Even though the first step of this process to obtain **3** has been actualized in the requested direction, demetallation to metal-free compound leads to an oxidized **4** as encountered by Nie et al. [31] with peripheral amino derivatives. Together with further metallated products (**5–7**), these new soluble complexes were characterized by elemental analysis, together with FT-IR, ¹H NMR, ¹³C NMR, ¹⁹F NMR, UV–vis, and mass spectral data.

3, **4** and **5**–**7** with eight pentafluorophenyl groups bound to the periphery are 2,3-bis(pentafluorophenyl)dinitrile derivative (1) [32] which was obtained by oxidative coupling of pentafluorophenylacetonitrile in the presence of I_2 . The solid product

^{*} Corresponding author. Tel.: + 90 212 866 33 00; fax: + 90 212 866 34 02. *E-mail address:* egonca@fatih.edu.tr (E. Gonca).

^{1387-7003/\$ –} see front matter 0 2012 Elsevier B.V. All rights reserved. doi:10.1016/j.inoche.2012.04.002

(1) was obtained in a yield of 64% (Scheme 1). It was a mixture of maleo and fumaronitriles and the latter was present in higher ratio as expected from the presence of pentafluorophenyl substituents. The product was brown colored and was very soluble in chloroform, dichloromethane and acetone, but insoluble in *n*-hexane. Its reluctance to turn into porphyrazine under the usual conditions (i.e. $Mg(OR)_2$ in refluxing *n*-propanol or *n*-butanol) points out the lower ratio of the maleonitrile component. This problem was solved by activation of the fumaronitrile derivative to the corresponding diiminopyrroline derivative (2) [33] which cyclizes quite easily to porphyrazine at reflux temperature of *n*-propanol or *n*-butanol [34]. In order to convert the diiminopyrroline derivative (2) into porphyrazine, we have made use of its template reaction in the presence of magnesium butanolate, which is the typical method generally applied in cyclotetramerization of these tetrapyrrole derivatives [35-37]. The dark blue octakis(pentafluorophenyl)porphyrazinatomagnesium (3) [38] was soluble in chloroform, dichloromethane, acetone, THF and pyridine (Fig. 1). The conversion of **3** to metal-free *seco*-porphyrazine derivative (**4**) [39] was achieved by treatment with trifluoroacetic acid followed by neutralization with ammonia and aqueous precipitation. The purity of the product could be raised by dissolving the crude substance in minimum amount of chloroform and then reprecipitating it by dropwise addition of the solution to *n*-hexane. Further reaction of **4** with copper (II) acetate, zinc (II) acetate and cobalt (II) acetate has led to the metallated secoderivatives, [2,3,7,8,12,13,17,18-octakis(pentafluorophenyl)-2-seco-2,3dioxoporphyrazinato] M(II) [M = Cu(II), Zn(II), Co(II)] (5-7) [40-42] (Scheme 1) (Fig. 2).

The newly synthesized compounds were characterized by using standard spectral methods (FT-IR, UV–vis, mass, ¹H, ¹³C and ¹⁹F NMR) in addition to elemental analysis. Spectral investigations for all new products were consistent with the assigned structures.

Elemental analysis results correspond closely with the calculated values for (1–7) (Table 1).

In the FT-IR spectrum of 2,3-bis(pentafluorophenyl)dinitrile derivative (1), the characteristic stretching vibration of C=N is observed at 2218 cm⁻¹, the aromatic C=C peak is at 1610 cm⁻¹ and the stretching vibrations of C – F are at 1346, 1166 and 1125 cm⁻¹. In the FT-IR spectrum of 3,4-bis(pentafluorophenyl)pyrroline-2,5-diimine (2), stretching vibration of N–H is observed at 3330 cm⁻¹, the aromatic C=C peak is at 1610 cm⁻¹ and the characteristic C – F peaks are at 1344, 1184, 1123 cm⁻¹ [20,21,26,43,44]. In the FT-IR spectrum of **3** the N–H groups in the inner core show absorption at 3335 cm⁻¹.



Fig. 1. [2,3,7,8,12,13,17,18-octakis(pentafluorophenyl)porphyrazinato]Mg(II) (3).

¹H NMR investigations of **2** and **4** provide the characteristic chemical shifts for the structures as expected. The N–H protons of **2** were also identified in the ¹H NMR spectrum with a broad singlet peak at δ = 4.80 ppm, presenting the diiminopyrroline derivative. ¹H NMR spectrum of **4** showed chemical shift belonging to metal-free porphyrazine ring protons (broad singlet) at – 1.35 ppm [45,46].

In the ¹³C NMR spectra of diamagnetic porphyrazines **3**, **4** and **6**, six different single chemical shifts for carbon atoms were clearly seen.

¹⁹F NMR spectroscopy has been a very useful technique for investigating the fluorinated compound. ¹⁹F NMR spectrum of **6** showed three different peaks at -142.8 ppm, -152.3 ppm and -161.9 ppm, respectively, relative to the fluorine atoms in the *ortho*, *para* and *meta* positions of the phenyl substituents and the spectrum showed the expected signals of the five fluorine atoms attached to the aromatic ring [47,48]. Integration of the peaks gave a 2:1:2 ratio as expected.



Scheme 1. (i) NaOMe, MeOH, I₂; (ii) NH₃, cat Na, Ethylene glycol; (iii) Mg turnings, I₂, n-BuOH; (iv) CF₃CO₂H; (v) EtOH and Cu(OAc)₂, Zn(OAc)₂, or Co(OAc)₂.



Fig. 2. [2,3,7,8,12,13,17,18-octakis(pentafluorophenyl)-2-*seco*-2,3-dioxoporphyrazine] and metal derivatives [M = 2H, Cu(II), Zn(II) or Co(II)] **(4–7**).

In addition to these supportive results for the structures, mass spectrometry results of pentafluorophenyl substituted porphyrazines (**3–7**) gave the characteristic molecular ion peaks, at m/z: 1664.1 [M]⁺, 1674.2 [M]⁺, 1736.8 [M]⁺, 1738.7 [M]⁺ and 1731.1 [M]⁺, respectively, confirming the proposed structures.

The most revealing data for tetrapyrrole systems are obtained by their electronic spectra in solution. In the electronic absorption spectra of the complexes (**3**, **5**–**7**), Q bands were observed at 668, 676, 680, and 672 nm, respectively in chloroform. A second intense and broad π – π * transition in the range 356–376 nm which is called Soret or B band is also a characteristic band as expected for tetrapyrroles (Table 2). As a consequence of the change in the symmetry of porphyrazine core (**3**) from D_{4h} to C_{2v}, the electronic absorption spectra of **4** display a split in the Q band at 620 and 696 nm [49]. UV–vis spectra of **5–7** in chloroform are shown in Fig. 3.

The aggregation behavior of porphyrazines in solution, which can be followed effectively by absorption studies, is a good indication of the interactions between the aromatic macrocycles of the porphyrazines. The aggregation behavior of **5** was investigated at different concentrations in chloroform (Fig. 4). In chloroform, as the concentration was increased, the intensity of the Q-band absorption increased in parallel, and there were no new bands because of the aggregated ones [50]. It is seen that the Beer–Lambert law was obeyed for compound **5** for concentrations ranging from 3×10^{-5} to 2.5×10^{-6} mol dm⁻³ (Fig. 4). And also the aggregation treatments of **5** were worked in dichloromethane at different concentrations ranging from 3×10^{-5} to 2.5×10^{-6} mol dm⁻³. There were only slight differences in B and Q bands as nm and because of the aggregated ones there were no new bands. UV–vis spectra of

Table 1				
Elemental	analysis	results	of	1–7. ^a

Compound	С	Н	Ν
1 2 3	46.76 (46.85) 44.90 (44.98) 46.25 (46.17) 45.78 (45.90)	0.83 (0.71)	6.90 (6.83) 9.77 (9.84) 6.80 (6.73) 6.81 (6.69)
4 5 6 7	43.78 (43.90) 44.18 (44.27) 44.11 (44.23) 44.28 (44.39)	0.10 (0.12)	6.57 (6.45) 6.33 (6.45) 6.59 (6.47)

^a Required values are given in parentheses.

Table 2

UV-vis data for the porphyrazines (3-7) in chloroform.

Compound		λ/nm (log ϵ/dm^3 m	$\lambda/nm~(log~\epsilon/dm^3~mol^{-1}~cm^{-1})$	
3	356 (4.91)	668 (4.88)	696 (4.61)	
4	348 (4.93)	620 (4.58)		
5	360 (4.98)	676 (4.79)		
6	376 (4.98)	680 (4.91)		
7	368 (4.95)	672 (4.89)		

7 $(1 \times 10^{-5} \text{ M})$ in solvents of different polarities (chloroform, dichloromethane, pyridine, and THF) are given in Fig. 5. There is almost no difference with respect to the changes in the nature of the solvent.

The substitution of polyfluorinated groups into the peripheral position of symmetrical porphyrazine and unsymmetrical *seco*-porphyrazines informs high solubility in organic solvents such as chloroform, dichloromethane, pyridine, and THF. The solubility of **3** in chloroform, DCM, THF, and acetone was determined as 44, 54, 60, and 63 mg/mL, respectively. But, it was not soluble in methanol. The increased solubility might be because of the higher solubility of pentafluorophenyl substituents in organic polar solvents. When the polarities of the solvents are increased, the solubility of complexes also increases as expected [19–21,26,46–48,50].

The spectral properties of the novel magnesium, metal-free secoporphyrazine and metallated *seco*-porphyrazines (M = Cu, Zn, Co) with eight pentafluorophenyl substituents at the periphery have been presented in this work. The 40 peripheral fluorine atoms in the structure increase the solubility of the complexes in halogenated solvents, acetone and THF. Fluorinated metallated porphyrazines have currently been receiving a great deal of attention due to their interesting electron transport characteristics [12,51]. While unsubstituted tetrapyrrole compounds exhibit p-type behavior due to the doping with electron accepting molecules, thin films of some metal hexadecafluoro-tetrapyrrole compounds exhibit n-type behavior. These properties resulted in a number of studies aiming at different applications like photovoltaic cells, rectifying junction and gas sensors. Also, fluorocarbons display enhanced hydrophobicity, chemical resistance and thermal stability in comparison to their hydrocarbon counterparts [43,52]. Consequently, porphyrazines reported here can be proposed as active compounds for catalytic and photosensor applications in homogeneous media [17,18,53].

Acknowledgment

This work is supported by the Scientific Research Fund of Fatih University under the project number P50021102_B.



Fig. 3. UV-vis spectra of 5-7 in chloroform.



Fig. 4. UV-vis spectra of 5 in chloroform at various concentrations.



Fig. 5. UV-vis spectra of 7 in various solvents.

References

- [1] R.P. Linstead, J. Chem. Soc. (1934) 1016-1017.
- [2] M.J. Stillman, T. Nyokong, in: C.C. Leznoff, A.B.P. Lever (Eds.), Phthalocyanines: Properties and Applications, vol. 1, VCH, New York, 1989, pp. 139–257.
- [3] J. Griffiths, J. Schofield, M. Wainwright, S.B. Brown, Dye. Pigment. 33 (1997) 65–78.
- [4] T.J. Dougherty, Photochem. Photobiol. 58 (1993) 895–900.
- [5] R.A. Hsi, D.I. Rosenthal, E. Glatstein, Drugs 57 (1999) 725-734.
- [6] I. Rosenthal, Photochem. Photobiol. 53 (1991) 859-870.
- [7] Q. Peng, J. Moan, Br. J. Cancer 72 (1995) 565-574.
- [8] R.W. Boyle, J. Rousseau, S.V. Kudrevich, M. Obochi, J.E. van Lier, Br. J. Cancer 73 (1996) 49-53.
- [9] D. Dolphin, The porphyrins, vols. 1-7, Academic Press, New York, 1978-1979.
- [10] C.C. Leznoff, A.B.P. Lever, Phthalocyanines: properties and applications, vols. 1–4, VCH, Weinheim, 1989–1996.
- [11] S.L.J. Michel, S. Baum, A.G.M. Barrett, B.M. Hoffman, in: K.D. Karlin (Ed.), Progress in Inorganic Chemistry, vol. 50, J. Wiley & Sons, New York, 2001, pp. 473–591.
- [12] Z.A. Bao, A.J. Lovinger, J. Brown, J. Am. Chem. Soc. 120 (1998) 207-208.
- [13] S. Hiller, D. Schlettwein, N.R. Armstrong, D. Wöhrle, J. Mater. Chem. 8 (1998) 945-954.
- [14] J.R.L. Smith, G. Reginato, Org. Biomol. Chem. 1 (2003) 2543–2549.
- [15] K.R. Venugopala Reddy, M.N.K. Harish, Fassiulla, M.H. Moinuddin Khan, J. Keshavayya, J. Fluorine Chem. 128 (2007) 1019–1025.
- [16] J. Piskorz, P. Skupin, S. Lijewski, M. Korpusinski, M. Sciepura, K. Konopka, S. Sobiak, T. Goslinski, J. Mielcarek, J. Fluorine Chem. 135 (2012) 265–271.

- [17] T. Goslinski, J. Piskorz, J. Photochem. Photobiol. C: Photochem. Rev. 12 (2011) 304–321.
- [18] J.R. Smith, G. Reginato, Org. Biomol. Chem. 1 (2003) 2543-2549.
- [19] T. Sugimori, S. Horike, M. Handa, K. Kasuga, Inorg. Chim. Acta 278 (1998) 253–255.
- [20] D. Koçak, E. Gonca, J. Fluorine Chem. 131 (2010) 1322-1326.
- [21] H. Kunt, E. Gonca, Polyhedron (2012), http://dx.doi.org/10.1016/j.poly.2012.03.015.
- [22] A. Nazlı, E. Gonca, A. Gül, J. Porphyrins Phthalocyanines 10 (2006) 996–1002.
 [23] E. Gonca, Ü.G. Baklacı, H.A. Dinçer, J. Porphyrins Phthalocyanines 12 (2008)
- 116–122.
- [24] E. Gonca, B. Keskin, J. Coord. Chem. 62 (2009) 2875–2882. [25] F. Gonca, Ü.G. Baklaci, H.A. Dincer, Polyhedron 27 (2008) 243:
- [25] E. Gonca, Ü.G. Baklacı, H.A. Dinçer, Polyhedron 27 (2008) 2431–2435.
 [26] E. Gonca, M. Altunkaya, Polyhedron 30 (2011) 1035–1039.
- [27] A.G. Montalban, S.J. Lange, L.S. Beall, N.S. Mani, D.J. Williams, A.J.P. White, A.G.M. Barrett, B.M. Hoffman, J. Org. Chem. 62 (1997) 9284–9289.
- [28] A.G. Montalban, S.M. Baum, A.G.M. Barrett, B.M. Hoffman, Dalton Trans. (2003) 2093–2102
- [29] D.M. Eichhorn, S. Yang, W. Jarrell, T.F. Baumann, L.S. Beall, A.J.P. White, D.J. Williams, A.G.M. Barrett, B.M. Hoffman, J. Chem. Soc. Chem. Commun. (1995) 1703–1704.
- [30] A.G. Montalban, H.G. Meunier, R.B. Ostler, A.G.M. Barrett, B.M. Hoffman, G. Rumbles, J. Phys. Chem. A 103 (1999) 4352–4358.
- [31] H. Nie, C.L. Stern, A.G.M. Barrett, B.M. Hoffman, Chem. Commun. (1999) 703-704.
- [32] Experimental data for compound 1: Yield: 1.65 g (64%). FT-IR, ν_{max}/(cm⁻¹): 2218 (C=N), 1662, 1614 (C=C, aromatic), 1575, 1512, 1465, 1380, 1340, 1275, 1161, 1119, 1055, 936, 902, 844, 706, 675, 588. ¹³C NMR (δ, ppm): 107.5, 118.5, 121.6, 137.4, 142.7, 143.3. ¹⁹F NMR (δ, ppm): -144.6 (*o*-fluoro), -154.4 (*p*-fluoro), -163.6 (*m*-fluoro). MS (ESI) *m/z*: 410.9 [M]⁺.
- [33] Experimental data for compound **2**: Yield: 0.31 g (48%). FT-IR, $\nu_{max}/(cm^{-1})$: 3330 (N–H), 1664, 1610 (C=C, aromatic), 1570, 1515, 1467, 1382, 1346, 1272, 1166, 1125, 1063, 942, 908, 842, 708, 671, 586. ¹H NMR (δ , ppm): 4.80 (br s, 3H, NH). ¹³C NMR (δ , ppm): 107.5, 125.3, 137.4, 142.2, 143.3, 164.3. ¹⁹F NMR (δ , ppm): – 144.8 (*o*-fluoro), – 154.5 (*p*-fluoro), – 163.8 (*m*-fluoro). MS (ESI) *m*/z: 427.8 [M]⁺.
- [34] T.F. Baumann, A.G.M. Barrett, B.M. Hoffman, Inorg. Chem. 36 (1997) 5661-5665.
- [35] W. Wolf, E. Degener, S. Petersen, Angew. Chem. 72 (1960) 963-966.
- [36] V.N. Kopranenkov, L.S. Goncharova, E.A. Luk'yanets, J. Org. Chem. 15 (1979) 1076–1082.
- [37] L.E.S. Schramm, B.M. Hoffman, Inorg. Chem. 19 (1980) 383-385.
- [38] Experimental data for compound 3: Yield: 117 mg (56%). FT-IR, v_{max}/(cm⁻¹): 1660, 1610 (C=C, aromatic), 1572, 1518, 1461, 1386, 1341, 1278, 1161, 1118, 1052, 936, 906, 843, 708, 671, 586. ¹³C NMR (ô, ppm): 107.2, 118.8, 121.1, 137.8, 142.3, 143.7. ¹⁹F NMR (ô, ppm): -142.7 (o-fluoro), -152.5 (p-fluoro), -161.6 (m-fluoro). MS (ESI) m/z: 1664.1 [M]⁺.
- [39] Experimental data for compound 4: Yield: 114 mg (68%). FT-IR, ν_{max}/(cm⁻¹): 3335 (N-H), 1722 (C=O), 1664, 1614 (C=C, aromatic), 1585, 1509, 1414, 1344, 1302, 1271, 1184, 1123, 1066, 906, 842, 705, 686, 582. ¹H NMR (δ, ppm): -1.35 (br s, 2H, NH). ¹³C NMR (δ, ppm): 107.4, 118.7, 121.3, 137.6, 142.1, 143.5. ¹⁹F NMR (δ, ppm): -141.5 (o-fluoro), -152.3 (p-fluoro), -161.8 (m-fluoro). MS (ESI) m/z: 1674.2 [M]⁺.
- [40] Experimental data for compound 5: Yield: 94 mg (54%). FT-IR, ν_{max}/(cm⁻¹): 1720 (C=O), 1666, 1614 (C=C, aromatic), 1580, 1512, 1415, 1348, 1305, 1274, 1188, 1126, 1075, 904, 840, 705, 685, 582. MS (ESI) m/z: 1736.8 [M]⁺.
- [41] Experimental data for compound 6: Yield: 129 mg (74%). FT-IR, v_{max}/(cm⁻¹): 1718 (C=O), 1662, 1608 (C=C, aromatic), 1585, 1502, 1408, 1344, 1309, 1272, 1184, 1126, 1066, 903, 847, 703, 686, 582. ¹³C NMR (δ, ppm): 107.4, 118.6, 121.4, 137.6, 142.5, 143.8. ¹⁹F NMR (δ, ppm): -142.8 (o-fluoro), -152.3 (pfluoro), -161.9 (m-fluoro). MS (ESI) m/z: 1738.7 [M]⁺.
- [42] Experimental data for compound 7: Yield: 100 mg (58%). FT-IR, ν_{max}/(cm⁻¹):
 1723 (C=O), 1660, 1613 (C=C, aromatic), 1581, 1511, 1416, 1349, 1301, 1274, 1188, 1123, 1071, 905, 842, 706, 682, 580. MS (ESI) m/z: 1731.1 [M]⁺.
- [43] A. Mueller, T. Kowalewski, K.L. Wooley, Macromolecules 31 (1998) 776-786.
- [44] L. Gao, X. Qian, J. Fluorine Chem. 113 (2002) 161-165.
- [45] H. Eichhorn, M. Rutloh, D. Wöhrle, J. Stumpe, J. Chem. Soc. Perkin Trans. 2 (1996) 1801–1810.
- [46] İ. Özçeşmeci, O. Güney, A.İ. Okur, A. Gül, J. Porphyrins Phthalocyanines 13 (2009) 753–759.
- [47] M. Selçukoğlu, E. Hamuryudan, Dye. Pigment. 74 (2007) 17-20.
- [48] M. Selçukoğlu, İ. Özçeşmeci, E. Hamuryudan, Polyhedron 29 (2010) 2710-2715.
- [49] Y. Rio, M.S. Rodriguez-Morgade, T. Torres, Org. Biomol. Chem. 6 (2008) 1877-1894.
- [50] H. Enkelkamp, R.J.M. Nolte, J. Porphyrins Phthalocyanines 4 (2000) 454–459.
- [51] E. Kol'tsov, T. Basova, P. Semyannikov, I. Iqumenov, Mater. Chem. Phys. 86 (2004) 222-227.
- [52] R.D. Chambers, Fluorine in organic chemistry, in: G.A. Olah (Ed.), Interscience Monographs on Organic Chemistry, John Wiley & Sons, New York, 1973.
- [53] S.P. Foxon, S.J.R. Lindsay, P.O. Brien, G. Reginato, J. Chem. Soc. Perkin Trans. 2 (2001) 1145–1153.