

The authors thank the National Science Council of the Republic of China for financial support (Grant No. NSC 85-2113-M-003-005, NSC 85-2113-M-003-002).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: DE1050). Services for accessing these data are described at the back of the journal.

References

- Johnson, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee, USA.
- Larson, A. C., Lee, F. L., Le Page, Y., Webster, M., Charland, J.-P. & Gabe, E. J. (1990). *NRCVAX. Crystal Structure System*. Chemistry Division, NRC, Ottawa, Canada.
- Latif, N., Girgis, N. S. & Michael, F. (1970). *Tetrahedron*, **26**, 5765–5772.
- Latif, N., Mishriky, N. & Mohsen, K. A. (1974). *J. Chem. Soc. Perkin Trans. 1*, pp. 875–879.
- Latif, N., Zeid, I. F. & Assad, F. (1970). *Chem. Ind. (London)*, 1539.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Riche, C. (1989). *NONIUS. Program for Data Reduction for Nonius CAD-4 diffractometer*. Institut de Chimie des Substances Naturelles du CNRS, Gif sur Yvette, France.

Acta Cryst. (1997). **C53**, 959–961

(*tert*-Butoxycarbonylamino)acetonitrile†

GEETA SOOD, CARL H. SCHWALBE AND WILLIAM FRASER

Pharmaceutical Sciences Institute, Aston University, Aston Triangle, Birmingham B4 7ET, England. E-mail: c.h.schwalbe@aston.ac.uk

(Received 21 October 1996; accepted 12 February 1997)


Abstract

The title compound, $C_7H_{12}N_2O_2$, is a useful intermediate for the synthesis of peptidic nucleic acids containing a backbone composed of repeating (2-aminoethyl)glycine units. Conjugation within the urethane system is facilitated by near planarity about both C—O and C—N bonds. There is a synperiplanar relationship between the O11 atom of the carbonyl group and the tertiary C7 atom, along with an antiperiplanar relationship between the C5 atom and the C10 methyl group.

Comment

Peptidic nucleic acids (PNAs) are useful nucleic acid mimetics which contain aminoethylglycine units in place of the conventional DNA sugar–phosphate backbone (Hyrup & Nielsen, 1996). PNAs containing the

usual DNA bases (Egholm *et al.*, 1993) along with pseudoisocytosine (Egholm *et al.*, 1995) possess anti-sense/antigene properties (Noble *et al.*, 1995; Knudsen & Nielsen, 1996). Our interest in modulating the hydrogen-bonding interactions between PNA and DNA oligomers prompted an investigation of PNAs containing other purine bases such as xanthine and diamino-purine. The title compound, (1), is a key intermediate for the synthesis of the PNA backbone structure (Meltzer, Liang & Matsudaira, 1995; Ravikumar, 1994) which is available as an oil through acylation of aminoacetonitrile hydrochloride using di-Boc anhydride or on dehydration of 3-(*tert*-Boc-amino)ethanamide (Houssin, Bernier & Heinichart, 1988) (where Boc is *butoxycarbonyl*). We have successfully prepared (1) in crystalline form for the first time and here we compare its crystal structure with that of other Boc-protected analogues.



In common with other urethanes, the O6—C5—N4 bond angle is approximately 10° lower than the trigonal value. The *trans* orientation of the O11 atom compared to the H atom attached to N4 is a general feature among Boc-protected amino acids. Like Boc-protected glycine (Semertzidis *et al.*, 1989) and alanine (Benedetti *et al.*, 1981) structures, (1) has a synperiplanar relationship about the C5—O6 bond between the carbonyl O11 atom and the tertiary C7 atom, as well as an antiperiplanar disposition of one methyl group (C10) to the carbonyl C5 atom. However, the locations of the multiply bonded C atom, C2, in (1) and the carboxyl C atom in the amino acids, show more variability: C2—C3—N4—C5 is (–)-synclinal in (1) but ranges from 62.9 to 110.7° in the latter. N4, the only proton donor group, donates a fairly weak intermolecular hydrogen bond to a nearby N1 atom related by $(2 - x, -0.5 + y, 0.5 - z)$ with $H4 \cdots N1$ 2.37 (2) Å and $N4 \cdots N1$ 3.132 (2) Å.




Fig. 1. ORTEP view (Johnson, 1976) of the molecule. Displacement ellipsoids are drawn at the 50% probability level. H atoms are drawn as small spheres of arbitrary radius.

† Alternative name: *tert*-butyl 3-cyano-2-azapropanoate.

Experimental

To a mixture of aminoacetonitrile hydrochloride (30.0 g, 0.325 mol) and di-Boc anhydride (70.9 g, 0.325 mol) in CH₂Cl₂ (1 l), Et₃N (125 ml, 0.9 mol) was added dropwise and the mixture stirred overnight at room temperature. After rotary evaporation of the product mixture, the residue was dissolved in ether, washed with water followed by brine and then dried (MgSO₄). Evaporation of the solvent gave the title compound (43.3 g, 85%). Recrystallization from EtOAc gave the analytical sample (m.p. 328–331 K). Thin-layer chromatography (EtOAc) *R_f* 0.58. IR (KBr disc): $\nu_{\max}$ 3390, 3010, 2995, 2252, 1720, 1534, 935, 839, 783 cm⁻¹ (all strong bands). ¹H NMR [250.1 MHz; (CDCl₃)]: δ 1.46 [s, 9 H, (CH₃)₃], 4.05 (d, 2 H, *J* = 6.0 Hz, CH₂), 5.30 p.p.m. (*m*, 1 H, NH, D₂O exchangeable). ¹³C NMR [250.1 MHz; (CDCl₃)]: δ 28.1 (CH₃)₃, 29.0 (CH₂), 81.1 (C), 115.6 (CN), 155.0 p.p.m. (CO). MS (CI⁺): *m/z* (I_r) 157 (*M*⁺H, 93%), 118 (100%), 74 (26%), 72 (6%), 56 (8%); Analysis calculated for C₇H₁₂N₂O₂: C 53.8, H 7.7, N 17.9%. Found: C 53.8, H 7.6, N 17.8%.

Crystal data

C ₇ H ₁₂ N ₂ O ₂	Cu K α radiation
<i>M_r</i> = 156.19	λ = 1.54178 Å
Orthorhombic	Cell parameters from 25 reflections
<i>P</i> 2 ₁ 2 ₁ 2 ₁	θ = 22.7–29.1°
<i>a</i> = 8.5261 (7) Å	μ = 0.729 mm ⁻¹
<i>b</i> = 10.0120 (7) Å	<i>T</i> = 293 (2) K
<i>c</i> = 10.2216 (14) Å	Tabular
<i>V</i> = 872.5 (2) Å ³	0.6 × 0.6 × 0.2 mm
<i>Z</i> = 4	Colourless
<i>D_x</i> = 1.189 Mg m ⁻³	
<i>D_m</i> not measured	

Data collection

Enraf–Nonius CAD-4 diffractometer	1624 reflections with <i>I</i> > 2σ(<i>I</i>)
ω/2θ scans	<i>R</i> _{int} = 0.0255
Absorption correction: ψ scans (North, Phillips & Mathews, 1968)	$\theta_{\max}$ = 69.91°
<i>T</i> _{min} = 0.788, <i>T</i> _{max} = 0.864	<i>h</i> = 0 → 10
1862 measured reflections	<i>k</i> = 0 → 12
1658 independent reflections	<i>l</i> = -12 → 12
	3 standard reflections frequency: 120 min intensity decay: 14%

Refinement

Refinement on <i>F</i> ²	$\Delta\rho_{\max}$ = 0.10 e Å ⁻³
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)] = 0.0286	$\Delta\rho_{\min}$ = -0.09 e Å ⁻³
<i>wR</i> (<i>F</i> ²) = 0.0843	Extinction correction: <i>SHELXL93</i> (Sheldrick, 1993)
<i>S</i> = 1.084	Extinction coefficient: 0.095 (5)
1658 reflections	Scattering factors from <i>International Tables for Crystallography</i> (Vol. C)
149 parameters	Absolute configuration: Flack (1983)
All H-atom parameters refined	Flack parameter = 0.5 (2)
$w = 1/[\sigma^2(F_o^2) + (0.0586P)^2 + 0.0396P]$	
where $P = (F_o^2 + 2F_c^2)/3$	
(Δ/σ) _{max} = -0.001	

Table 1. Selected geometric parameters (Å, °)

N1—C2	1.134 (2)	C5—O6	1.3400 (14)
C2—C3	1.471 (2)	O6—C7	1.4774 (13)
C3—N4	1.436 (2)	C7—C8	1.514 (2)
N4—C5	1.354 (2)	C7—C10	1.515 (2)
C5—O11	1.2090 (15)	C7—C9	1.511 (2)
N1—C2—C3	177.7 (2)	O6—C7—C8	110.13 (11)
N4—C3—C2	112.50 (11)	O6—C7—C10	102.34 (10)
C5—N4—C3	120.07 (11)	C8—C7—C10	112.11 (14)
O11—C5—O6	126.77 (11)	O6—C7—C9	109.64 (11)
O11—C5—N4	123.50 (11)	C8—C7—C9	112.93 (14)
O6—C5—N4	109.73 (9)	C10—C7—C9	109.16 (12)
C5—O6—C7	120.76 (9)		
C2—C3—N4—C5	-84.25 (15)	C5—O6—C7—C10	-178.53 (11)
O11—C5—O6—C7	-2.5 (2)		

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989). Cell refinement: *CAD-4 Software*. Data reduction: *CADABS* (Gould & Smith, 1996). Program(s) used to solve structure: *MULTAN11/84* (Main, Germain & Woolfson, 1984). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993). Molecular graphics: *ORTEPII* (Johnson, 1976). Software used to prepare material for publication: *SHELXL93*.

We thank the EPSRC for a Total Technology Studentship (GS) and the EPSRC Mass Spectrometry Service (Swansea). We also thank Celltech Therapeutics (Slough) for continuous support of the DNA molecular recognition project and Dr M. A. W. Eaton and Dr J. Turner for helpful discussions.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BM1121). Services for accessing these data are described at the back of the journal.

References

- Benedetti, E., di Blasio, B., Pavone, V., Pedone, C., Toniolo, C. & Bonora, G. M. (1981). *Biopolymers*, **20**, 1635–1649.
- Egholm, M., Behrens, G., Christensen, L., Berg, R. H., Nielsen, P. E. & Buchardt, O. (1993). *J. Chem. Soc. Chem Commun.* pp. 800–801.
- Egholm, M., Christensen, L., Dueholm, K. L., Buchardt, O., Coull, J. & Nielsen, P. E. (1995). *Nucleic Acids Res.* **23**, 217–222.
- Enraf–Nonius (1989). *CAD-4 Software*. Version 5. Enraf–Nonius, Delft, The Netherlands.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Gould, R. O. & Smith, D. E. (1986). *CADABS. Program for CAD-4 Data Reduction*. University of Edinburgh, Scotland.
- Houssin, R., Bernier, J.-L. & Heinichart, J.-P. (1988). *Synthesis*, pp. 259–261.
- Hyrup, B. & Nielsen, P. E. (1996). *Bioorg. Med. Chem. Lett.* **4**, 5–23.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Knudsen, H. & Nielsen, P. E. (1996). *Nucleic Acids Res.* **24**, 494–500.
- Main, P., Germain, G. & Woolfson, M. M. (1984). *MULTAN11/84. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Universities of York, England, and Louvain, Belgium.
- Meltzer, P. C., Liang, A. Y. & Matsudaira, P. (1995). *J. Org. Chem.* **60**, 4305–4308.
- Noble, S. A., Bonham, M. A., Bisi, J. E., Bruckenstein, D. A., Brown, P. H., Brown, S. C. & Cadilla, R. G. (1995). *Drug Dev. Res.* **34**, 184–195.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Ravikumar, V. T. (1994). *Synth. Commun.* **24**, 1767–1772.

Semertzidis, M., Matsoukas, J., Nastopoulos, V., Hondrelis, J., Voliotis, S. & Leban, I. (1989). *Acta Cryst.* C45, 1474–1475.
 Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.

Acta Cryst. (1997). C53, 961–963

Isotopically Substituted Bis(ethylenedithio)tetrathiafulvalene at 15 K by Time-of-Flight Neutron Diffraction

DIANNA M. YOUNG,^a ARTHUR J. SCHULTZ,^a ARAVINDA M. KINI^b AND JACK M. WILLIAMS^b

^aIntense Pulsed Neutron Source, Argonne National Laboratory, Argonne, IL 60439, USA, and ^bChemistry and Materials Science Divisions, Argonne National Laboratory, Argonne, IL 60439, USA. E-mail: ajschultz@anl.gov

(Received 19 November 1996; accepted 21 February 1997)


Abstract

The title compound, C₁₀D₈S₈, (ET), has been synthesized whereby all eight S atoms, the four ethylene C atoms and all eight H atoms have been isotopically substituted with ³⁴S, ¹³C and ²H, respectively. Analysis of the neutron scattering lengths derived from single-crystal time-of-flight neutron diffraction data obtained at *T* = 15 K confirms the isotopic distribution. The title phase crystallizes in the space group *P*₂₁/*n* with bond lengths and angles consistent with neutral ET.

Comment

The title compound, (I), (ET), participates as the electron-donor molecule in a large number of organic superconducting salts (Williams *et al.*, 1991). In order to investigate the mechanism for the electron pairing in this class of superconductors, a variety of isotopically substituted ET molecules with different distributions of isotopes throughout the molecule have been synthesized (Kini *et al.*, 1996). Salts of these organic donor molecules have been studied to determine the isotopic effect on *T*_c in the superconducting crystals. In the crystal used in this neutron diffraction experiment, the isotopic substitution of the ET molecule involved the replacement of the four C atoms in the terminal 1,2-ethanedithyl groups with ¹³C, all eight S atoms with ³⁴S and all eight H atoms with deuterium, ²H. The substitution increases the mass of the ET molecule, hereafter denoted by ²H₈¹³C₄³⁴S₈–ET, by 28 a.m.u., which is about 7% of the original mass. From the study by Kini *et al.* (1996), it was shown that the C and S substitution has a normal BCS-like mass (*M*) effect (*T*_c proportional to *M*^{−α}),

whereas deuterium substitution produces an inverse isotopic effect (*T*_c increases). We present here the neutron structure of ²H₈¹³C₄³⁴S₈–ET at 15 K in order to confirm the isotopic distribution and to provide high-precision low-temperature structural parameters.



The only previous structure of neutral ET is the room-temperature single-crystal X-ray structure in space group *P*₂₁/*c* by Kobayashi, Kobayashi, Yukiyoishi, Saito & Inokuchi (1986). We had transformed the atomic coordinates from *P*₂₁/*c* to *P*₂₁/*n* in our previous comparison of the ET structure with the isomorphous Au(DDDT)₂ (DDDT = 5,6-dihydro-1,4-dithiin-2,3-dithiolate) crystal structure (Schultz *et al.*, 1987). The transformation of the atomic coordinates from the *P*₂₁/*c* unit cell in Kobayashi *et al.* (1986) to the *P*₂₁/*n* unit cell in this paper is (*x* − *z*, −*y*, −*z*) + (1/2, 1/2, 0).

The *P*₂₁/*n* room-temperature unit cell of neutral ET reported by Kobayashi *et al.* (1986) is *a* = 6.614 (11), *b* = 13.985 (2), *c* = 15.721 (3) Å, β = 93.81 (2)° and *V* = 1451 (5) Å³. The low-temperature data of isotopically substituted ET gave a slightly smaller volume, as expected, of 1416 (1) Å³ and *a* = 6.661 (1), *b* = 13.652 (2), *c* = 15.613 (2) Å, β = 94.281 (2)°. The bond lengths of the low-temperature isotopically substituted ET appear to be slightly longer than those obtained from room-temperature data. Shorter bond distances of the room-temperature neutral ET are probably attributable to anisotropic thermal motion. Isotopically substituted ET bond distances at 15 K are C1—C2 = 1.359 (2), C5—C6 = 1.360 (2), S2—C4 = 1.771 (3) and S4—C6 = 1.769 (3) Å. Kobayashi *et al.* (1986) reported 1.319, 1.328, 1.757 and 1.751 ± (0.007–0.012) Å, respectively.

The ET molecules pack as face-to-face dimeric pairs. Many of the S—S contacts between different molecules have distances close to the sum of the van der Waals radii (3.70 Å). The S··S contacts are listed in Table 2.

The crystal for this data collection was obtained from a batch of crystals which underwent mass spectrometry to verify isotopic substitution. In order to confirm the degree of isotopic substitution of this crystal, scattering lengths for each of the atoms were refined. Scattering lengths in units of 10^{−12} cm from Sears (1986) are: *b*(¹²C) = 0.66511 (16), *b*(¹³C) = 0.619 (9), *b*(³²S) = 0.2804 (2), *b*(³⁴S) = 0.348 (3), *b*(H) = −0.37406 (11), *b*(D) = 0.6671 (4). The average refined scattering length in this structure of the ethanedithyl ¹³C atoms is 0.656 (3) × 10^{−12} cm, which appears to be closer in magnitude to ¹²C but is also within three standard deviations of the ¹³C scattering length. Thus, the refined ethanedithyl