Gold(I)-Catalyzed Rearrangement of *N*-Aryl 2-Alkynylazetidines to Pyrrolo[1,2-*a*]indoles

LETTERS 2013 Vol. 15, No. 4 836–839

ORGANIC

Nicolas Kern, Marie Hoffmann, Aurélien Blanc,* Jean-Marc Weibel, and Patrick Pale*

Laboratoire de Synthèse, Réactivité Organiques & Catalyse, associé au CNRS, Institut de Chimie, Université de Strasbourg, 4 rue Blaise Pascal, 67070 Strasbourg, France

ablanc@unistra.fr; ppale@unistra.fr

Received December 22, 2012



Various *N*-aryl-2-alkynylazetidines were very efficiently converted to pyrrolo[1,2-*a*]indoles with gold catalysts, especially the 2-biphenyl-dicyclohexylphosphino-gold(I) hexafluoroantimonate, in dichloromethane at room temperature. Additionally, two formal syntheses of bioactive nonnatural compounds, i.e. 7-methoxymitosene and an 5-HT_{2C} receptor agonist, have been achieved.

The now well-known activation of an alkene or alkyne by π -coordination with gold salts or complexes¹ and our involvement in gold multifaceted catalysis,² i.e. cascade reactions implying both π and σ Lewis acidities of gold,³ led us to imagine a rapid access to pyrrolo[1,2-*a*]indoles **2**, based on the opening of *N*-aryl 2-alkynylazetidines **1** (Scheme 1). The process is based on π and σ Lewis acidities of gold, which could favor alkynylazetidine opening and allene formation (Scheme 1) (**A** and/or **A'** upon π or σ Au activation). The benzazocine intermediate **B** would then be cyclized upon another gold activation, leading after protodeauration of **C** to the pyrrolo[1,2-*a*]indole.

The pyrrolo[1,2-*a*]indole tricyclic structure is a common motif found in a number of naturally occurring products,

among which mitomycins and its derivatives,⁴ known as mitosanes and mitosenes,⁵ are the most well-known due to their antitumor activities (Figure 1).⁶ Yuremamine, isolated from the stem bark of *Mimosa hostilis*, is another recent interesting example, due to its traditional use promoting hallucinogenic and psychoactive effects.⁷ Another psychoactive alkaloid harmalidine and its derivatives have been isolated from the seeds of *Pegalum harmala*.⁸ The pyrrolo[1,2-*a*]indole motif can also be found in some nonnatural compounds, exhibiting useful biological activities (Figure 1).^{9,10} Due to their interesting properties, these compounds have given rise to various synthetic approaches.^{11,12} Most of them relied on multistep sequences,

For selected recent reviews on gold chemistry, see: (a) Lu, B.-L.;
 Dai, L.; Shi, M. Chem. Soc. Rev. 2012, 41, 3318. (b) Corma, A.; Leyva-Pérez, A.; Sabater, M. J. Chem. Rev. 2011, 1657. (c) Krause, N.; Winter, C. Chem. Rev. 2011, 1994. (d) Gagosz, F. Actual. Chim. 2010, 347, 12. (e) Belmont, P.; Parker, E. Eur. J. Org. Chem. 2009, 6075. (f) Hashmi, A. S. K.; Hutchings, G. J. Angew. Chem., Int. Ed. 2006, 45, 7896.

^{(2) (}a) Kern, N.; Blanc, A.; Miaskiewicz, S.; Robinette, M.; Weibel, J.-M.; Pale, P. J. Org. Chem. **2012**, 77, 4323. (b) Kern, N.; Blanc, A.; Weibel, J.-M.; Pale, P. Chem. Commun. **2011**, 47, 6665. (c) Britton, J.; Camp, J. E. Chemistry Today **2012**, 30 (3, Suppl.), 6.

⁽³⁾ Yamamoto, Y. J. Org. Chem. 2007, 72, 7817.

^{(4) (}a) Kudo, S.; Marumo, T.; Tomioka, T.; Kato, H.; Fujimoto, Y. Antibiot. Chemother. **1958**, *8*, 228. (b) Lefemine, D. V.; Dann, M.; Barbatschi, F.; Hausmann, W. K.; Zbinovsky, V.; Monnikendam, P.; Adam, J.; Bohonos, N. J. Am. Chem. Soc. **1962**, *84*, 3184. (c) Tulinsky, A. J. Am. Chem. Soc. **1962**, *84*, 3188.

^{(5) (}a) Webb, S.; Cosulich, D. B.; Mowat, J. H.; Patrick, J. B.; Broschard, R. W.; Meyer, W. E.; Williams, R. P.; Wolf, C. F.; Fulmor, W.; Pidackas, C.; Lancaster, J. E. *J. Am. Chem. Soc.* **1962**, *84*, 3185. (b) Webb, S.; Cosulich, D. B.; Mowat, J. H.; Patrick, J. B.; Lancaster, J. E.; Broschard, R. W.; Meyer, W. E.; Williams, R. P.; Wolf, C. F.; Fulmor, W.; Pidackas, C. *J. Am. Chem. Soc.* **1962**, *84*, 3187.

^{(6) (}a) Bradner, W. T. *Cancer Treat. Rev.* **2001**, *27*, 35. (b) For the initially proposed but now proven mode of action, see: Iyer, V. N.; Szybalski, W. Science **1964**, *145*, 55.

⁽⁷⁾ Vepsalainen, J. J.; Auriola, S.; Tukiainen, M.; Ropponen, N.; Callaway, J. *Planta Med.* **2005**, *71*, 1053.

⁽⁸⁾ Siddiqui, S.; Khan, O. Y.; Siddiqui, B. S.; Faizi, S. Phytochem. 1987, 26, 1548.

⁽⁹⁾ Peters, R.; Waldmeier, P.; Joncour, A. Org. Process Res. Dev. 2005, 9, 508.

⁽¹⁰⁾ Allen, G. R.; Poletto, J. F.; Weiss, M. J. J. Am. Chem. Soc. 1964, 86, 3877.

especially in the mitomycin area,13 and any additional convergent and rapid route would be beneficial to this area.

Scheme 1. Hypothesis for Au-Catalyzed Conversion of N-Aryl Alkynylazetidines to Pyrrolo[1,2-a]indoles





Figure 1. Natural products and some bioactive non-natural compounds exhibiting the pyrrolo[1,2-*a*]indole motif.

In the search for optimal conditions, the simple 2-(hept-1-ynyl)-2-methyl-1-phenylazetidine 1a was readily prepared in three simple steps with a good overall yield (see Supporting Information) and submitted to various Au catalysts (Table 1).¹⁴ Simple gold(I) or gold(III) chlorides gave the expected compound 2a in dichloromethane in a clean reaction but with low conversions despite long reaction times (Table 1, entries 1-2). The more soluble and more electrophilic cationic phosphinogold(I) complexes showed a higher reactivity but a high dependence on solvent nature (entries 3-8).

Table 1. Screening of Reaction Conditions for the Au-Catalyzed Transformation of N-Phenyl 2-Alkynylazetidine 1a into 2a



1

2

3

4	"	$(CH_2CI)_2$, rt	0.5	78
5	//	THF, rt	0.33	86
6	//	toluene, rt	18	37^b
7	//	acetone, rt	18	65^b
8	"	CH ₃ CN, rt	18	55^b
9	$L^{1}AuNTf_{2}(4)$	CH_2Cl_2 , rt	0.5	92
10	$L^{1}AuSbF_{6}.MeCN(4)$	CH_2Cl_2 , rt	0.33	98
11	$L^{2}AuSbF_{6}.MeCN(4)$	CH_2Cl_2 , rt	2	98
12	$L^{1}AuSbF_{6}.MeCN(2.5)$	CH_2Cl_2 , rt	0.5	98
13	$AgSbF_{6}(4)$	CH ₂ Cl ₂ , rt	18	$< 10^{\circ}$

 ${}^{a}C = 0.2 \text{ mol/L}$. ^bConversion, the starting material accounting for the mass balance. ^c Degradation occurred leading to unidentified byproducts.

The Gagosz catalyst¹⁵ rapidly gave 2a in high yield in chlorinated solvents (Table 1, entries 3-4) or in THF (entry 5), but slow reactions and low conversions were observed in less polar solvents (entry 6) as well as in more polar and coordinating solvents (entries 7-8). Switching to the bulky and stabilizing (Cy_2) JohnPhos ligand L¹ induced a fast and very efficient reaction (entry 9). Combining this ligand or the bulkiest JohnPhos ligand L^2 and a more electrophilic counterion, such as SbF_6^- , gave almost quan*titatively* the pyrrolo[1,2-*a*]indole **2a** (entries 10–11) and allowed the catalytic charge to be decreased to only 2.5% (entry 12).¹⁶ Control experiments showed that the silver salt used to prepare the cationic gold species was not responsible for the rearrangement, but led to degradation (entry 13).

With these results in hand, the scope of this gold(I)catalyzed rearrangement of N-aryl 2-alkynylazetidines to pyrrolo[1,2-a]indoles was then examined. Various N-aryl 2-alkynylazetidines, 1a-j, were thus prepared and submitted to the optimal conditions described above (Table 2).

Steric and electronic effects at the acetylene moiety on one hand and electronic effects at the N-aryl group on the other hand should influence the first opening-cyclization step and would thus shed some light on the actual mechanism (see Scheme 1). Their role was thus examined with substrates **1a**-**c** and **1d**-**g**. A branched alkyl chain next to the acetylene moiety did not significantly change the

⁽¹¹⁾ For selected references on the synthesis of pyrrolo[1,2-a]indoles, see: (a) Chen, K.; Zhang, Z.; Shi, M. Chem. Commun. 2012, 48, 7696. (b) Zeldin, R. M.; Toste, F. D. Chem. Sci. 2011, 2, 1706. (c) Patil, D. V.; Cavitt, M. A.; France, S. Org. Lett. 2011, 13, 5820.

⁽¹²⁾ For a leading reference on vuremamine synthetic studies, see: Johansen, M. B.; Kerr, M. A. Org. Lett. 2008, 10, 3497.

⁽¹³⁾ For a review on mitomycin synthesis, see: (a) Andrez, J.-C. Beilst. J. Org. Chem. 2009, 5, n°33.

⁽¹⁴⁾ For gold-catalyzed formation of pyrrolo[1,2-a]indoles, see: (a) Kusama, H.; Miyashita, Y.; Takaya, J.; Iwasawa, N. Org. Lett. **2006**, 8, 289. (b) Yan, Z.-Y.; Xiao, Y.; Zhang, L. Angew. Chem., Int. Ed. **2012**, 51, 8624.

⁽¹⁵⁾ Mezailles, N.; Ricard, L.; Gagosz, F. Org. Lett. 2005, 7, 4133.

⁽¹⁶⁾ A further decrease in catalyst amount led to slower reactions (e.g., with 1%, the reaction required almost 10 days to give a 64% conversion, but still in a spot-to-spot transformation).

	$R^1 \longrightarrow R^3$ $R^1 \longrightarrow R^2$	Biph(C	y ₂)PAuSbF ₆ .CH ₃ CN (2.5 mol %) CH ₂ Cl ₂ , rt	R^3		2
entry	alkynylazetidin	ies, 1	pyrrolo[1,2-a]-ind	oles, 2	time (h)	yield (%)
1	C ₅ H ₁₁	1a	C ₅ H ₁₁	2a	0.5	98
2	Cy- <u></u> →N	1b	Cy N	2b	0.5	91
3	Ph-=-	1c	Ph N	2c	2	80^a
4		1d	MeO	2d	0.5	99
5	C ₆ H ₁₃	1e	CI	2e	1	82
6		1f		2f 1:1 2f'	0.5	83
7	CI C ₅ H ₁₁	1g		2g 1:4 2g'	1	92
8	C ₅ H ₁₁	1h	C ₅ H ₁₁	2h	0.75	91
9	ОМе С ₅ H ₁₁ -=	1i	MeO	2i	3	91 ^{<i>b</i>}
10		1j	OTBS N OBn	2j	3	82 ^{<i>b</i>}

 Table 2. Scope of the Gold(I)-Catalyzed Formation of

 Pyrrolo[1,2-a]indoles from N-Aryl 2-Alkynylazetidines

 a Reaction run at 40 °C to complete the conversion. b Reaction run at 0 °C to avoid degradation.

reaction rate compared to the simple linear alkyl group (Table 2, entry 2 vs 1). With a phenyl group conjugated with the acetylene moiety, the reaction proceeded as well, although the isolated yield was lower due to product instability (entry 3 vs 1). With an electron-donating group at the *para* position relative to the azetidine nitrogen, the rearrangement proceeded as rapidly but very efficiently (entry 4) while an electron-withdrawing group slightly slowed the reaction leading to a lower isolated yield (entry 5). With an electron-donating or -withdrawing

group at the *meta* position relative to the azetidine nitrogen, mixtures of rearranged products were obtained (entries 6-7). The methoxylated derivative reacted more rapidly than its chlorinated analog, but a better yield was obtained with the latter, suggesting contradictory electronic influences. However, regioisomeric ratios were different. from 1:1 for a methoxy group to 1:4 for the chloro substituent (entry 6 vs 7). In the latter, the major product was the 3-chloro-9-pentylpyrrolo[1,2-a]indole 2g', the less hindered isomer, suggesting that steric interactions influenced the reaction. To conclude on these aspects, the meta-isopropyl derivative 1h was rearranged under standard conditions (entry 8). A single regioisomer was observed and isolated in high yield, characterized as the 3-isopropyl-9pentylpyrrolo[1,2-a]indole 2h, again the less hindered isomer. These results thus confirmed the major role of steric interactions in this rearrangement, probably during the first arylation step (Scheme 1) in which the meta-aryl substituent would experience 1,3-allylic strain with the acetylenic substituent. We were able to ensure that β -substitution of the azetidine did not affect the reaction. As expected, the rearrangement of *cis*-1i-j very efficiently afforded the corresponding tricyclic indoles in high yields, although this reaction has to be performed at 0 °C to avoid OBn elimination (entries 9-10).

To go further, and with total synthesis perspectives.¹⁷ we applied this Au-catalyzed rearrangement toward a few pyrrolo[1,2-a]indoles, either known intermediates in the synthesis of bioactive compounds or models of natural products (Table 3). We selected 7-ethoxy-2,3-dihyropyrrolo-[1,2-*a*]indole,⁹ an intermediate en route to an agonist of the 5-HT_{2c} receptor (Figure 1), linked to the regulation of food intake and anxiety.¹⁸ This compound could easily be obtained through the Au-catalyzed rearrangement of the simple N-(4-ethoxyphenyl)-2-ethynylazetidine 1k in 85% yield (Table 3, entry 1). Yet, the psychoactive alkaloid harmalidine¹⁹ seemed to have never been synthesized (Figure 1). The present Au-catalyzed rearrangement would offer the *first* approach to such a compound and analogs, starting from either N-(aryl) 2-(4-hydroxybut-1-ynyl)-3,3dimethylazetidine 11 or 1m. The hindered tert-butyldimethylsilyl-protected derivative 11 gave 21 and 21' in good yield but as a 1:1 mixture (entry 2). To obtain the required regioselectivity toward the harmalidine core, we introduced a bromo substituent (1m), blocking one ortho position. The latter indeed furnished the expected regioisomer in good yield (entry 3). Looking for the mitomycins skeleton formation, we then engaged the free and TIPS-protected propargyl alcohols **1n** and **1o** in the gold-catalyzed rearrangement. We rapidly observed the formation of the same compound in both reactions (entries 4-5), which was identified as the 3,3'-methylbispyrrolo[1,2-a]indole 3 coming from the condensation of the desired pyrrolo[1,2-a]indole motif on itself followed by a loss of formaldehyde.²⁰

⁽¹⁷⁾ Rudolph, M.; Hashmi, A. S. K. *Chem. Soc. Rev.* 2012, *41*, 2448.
(18) Tecot, L. H.; Sun, L. M.; Akana, S. F.; Strack, A. M.; Lowenstein, D. H.; Dallman, M. F.; Julluis, D. *Nature* 1995, *374*, 542.

Lowenstein, D. H.; Dallman, M. F.; Julluis, D. *Nature* **1995**, *374*, 542. (19) Siddiqui, S.; Khan, O. Y.; Siddiqui, B. S.; Faizi, S. *Phytochem.* **1987**, *26*, 1548.

Rewardingly, the benzyl protected analog 1p gave the expected carbon core 2p of the mitomycin family, stable enough to be isolated in good yield (entry 6). Finally, the simple mitosene derivative 2q was efficiently produced from 2-ethynyl-1-(2,4,5-trimethoxy-3-methylphenyl)-azetidine 1q, although forcing conditions had to be used (entry 7). This result achieved the formal synthesis of the bioactive 7-methoxymitosene²¹ (Figure 1).

Table 3. Gold(I)-Catalyzed Formation of Pyrrolo[1,2-a]indoles of Biological Interest^a



^{*a*} Performed with (Cy₂)JohnPhosAuSbF₆·MeCN (4 mol %) at rt in CH₂Cl₂. ^{*b*} Performed with JohnPhosAuSbF₆·MeCN (10 mol %) at 80 °C. ^{*c*} R = H. ^{*d*} R = TIPS (triisopropylsily). ^{*e*} R = Bn (benzyl). ^{*f*} Isolated yields after 72 h of stirring using 10 mol % of catalyst added in four portions.

To gain insight into the proposed mechanism,²² we prepared the perdeuterated compound 1r and monitored the deuterium labeling into the rearranged product 2r (Scheme 2). In the latter, a single deuterium label was found at the allylic position within the pyrrolidine ring, at

- (21) Liang, L.; Wang, Y.; Zhang, L. Org. Lett. 2012, 14, 3736.
- (22) Hashmi, A. S. K. Angew. Chem., Int. Ed. 2010, 49, 5232.





the expected position of the gold atom after rearrangement (**C**, Scheme 1). This result corroborates the second part of the proposed mechanism, although the allyl gold part in **C** seems to not be protodemetalated through an SE' mechanism as recently reported.²³

However, the first step is still unclear. On one hand, cyclizations of propargyl anilines initiated by π -coordination have been reported.²⁴ Such a pathway would lead to \mathbf{A}' in our case. On the other hand, the alkynyl azetidine opening can also be viewed as a gold-catalyzed amino-Claisen rearrangement, according to a few precedents in the literature.²⁵ The strain release of the four-membered ring would thus be the driving force of this process, and azaphilic gold interaction would initiate such a pathway leading to A, in analogy to very recent calculations performed for gold(I)-catalyzed Claisen rearrangement.²⁶ The fact that β -lactams, precursors of the azetidine derivatives (see Supporting Information), failed to give pyrrolo[1,2-a]indolones supports the azaphilic activation pathway (A in Scheme 1), as in this case the carbophilic mechanism should have been favored by the weakly donating nitrogen.

In conclusion, we have developed a novel and efficient Au-catalyzed rearrangement of *N*-aryl-2-alkynylazetidines to pyrrolo[1,2-*a*]indoles. Further investigations of this rearrangement and applications in alkaloid synthesis are ongoing in our laboratory. In the latter aspect, two formal syntheses (5-HT_{2C} receptor agonist and 7-methoxymitosene) have been achieved.

Acknowledgment. We gratefully acknowledge the CNRS and the Agence Nationale de la Recherche (Grant ANR-11-JS07-001-01). N.K. and M.H. thank the French Ministry of Research for a PhD fellowship.

Supporting Information Available. Complete experimental procedures, characterization data, and spectral data. This material is available free of charge via the Internet at http://pubs.acs.org.

⁽²⁰⁾ Pawlak, J. M.; Khau, V. V.; Hutchison, D. R.; Martinelli, M. J. J. Org. Chem. 1996, 61, 9055.

⁽²³⁾ Hasmi, A. S. K.; Schuster, A. M.; Litters, S.; Rominger, F.;
Pernpointner, M. *Chem.—Eur. J.* 2011, *17*, 5661.
(24) (a) Menon, R. S.; Findlay, A. D.; Bissember, A. C.; Banwell,

 ^{(24) (}a) Menon, R. S.; Findlay, A. D.; Bissember, A. C.; Banwell,
 M. G. J. Org. Chem. 2009, 74, 8901. (b) Nevado, C.; Echavarren, A.
 Chem.—Eur. J. 2005, 11, 3155.

^{(25) (}a) Sherry, B. D.; Toste, F. D. J. Am. Chem. Soc. **2004**, *126*, 15978. (b) Tejedor, D.; Mendez-Abt, G.; Cotos, L.; Garcia-Tellado, F. Chem. Soc. Rev. **2013**, *42*, 458. For a gold-catalyzed tandem Claisencyclization reaction, see: (c) Sherry, B. D.; Maus, L.; Lafortza, B. N.; Toste, F. D. J. Am. Chem. Soc. **2006**, *128*, 8132. (d) Saito, A.; Konishi, T.; Hanzawa, Y. Org. Lett. **2010**, *12*, 372.

⁽²⁶⁾ Vidhani, D. V.; Cran, J. W.; Krafft, M. E.; Manoharan, M.; Alabugin, I. V. J. Org. Chem. **2013**dx.doi.org/10.1021/jo302152j.

The authors declare no competing financial interest.