## Preparation of [(mes)Os(CNR)Cl<sub>2</sub>] and Reactions of the Methyl **Isocyanide Complex with Organolithium and Grignard Reagents.** X-ray Crystal Structure of $[(mes)Os(=C(NHCH_3)C_8H_5)(C_8H_5)_2]$

Helmut Werner,\* Ulrich Wecker, Michael Schulz, and Stefan Stahl

Institut für Anorganische Chemie der Universität Würzburg, Am Hubland, D-8700 Würzburg, Germany

Received March 4. 1991

The isocyanide complexes  $[(mes)Os(CNR)Cl_2]$  (2-4) are prepared almost quantitatively from  $[(mes)OsCl_2]_n$ (1) and CNR in dichloromethane. Reactions of 2 (R = CH<sub>3</sub>) with CH<sub>3</sub>Li and C<sub>6</sub>H<sub>5</sub>Li in hydrocarbon-ether give the dimethyl and diphenyl derivates  $[(mes)OsR_2(CNCH_3)]$  (5, 6) in excellent yields. Treatment of 2 with CH<sub>3</sub>MgI leads to the formation of a mixture of  $[(mes)OsCH_3(CNCH_3)I]$  (7) and  $[(mes)Os(CNCH_3)I_2]$ (9), which can be separated by column chromatography. Analogously, from 2 and  $C_6H_5MgI$ , both [(mes)OsC<sub>6</sub>H<sub>5</sub>(CNCH<sub>3</sub>)I] (8) and 9 have been obtained. In contrast, reaction of 2 with C<sub>6</sub>H<sub>5</sub>MgBr gives, besides [(mes)OsC<sub>6</sub>H<sub>5</sub>(CNCH<sub>3</sub>)Br] (11), the carbene complex [(mes)Os(=C(NHCH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>)(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>] (10) as the main product. The X-ray crystal structure of 10 has been determined (monoclinic space group P2<sub>1</sub>/n) with a = 12.402 (3) Å, b = 14.237 (2) Å, c = 13.622 (2) Å,  $\beta = 91.54$  (1)°). The molecule possesses a piano-stool configuration with an Os-C(carbene) distance of 1.992 (5) Å.

#### Introduction

As an extension of earlier work from our laboratory on (arene)osmium(II) and osmium(0) complexes,<sup>1,2</sup> we have recently described a series of related compounds in which, besides the six-membered ring, one or two CO but no phosphane or phosphite ligands are coordinated to the metal.<sup>3-5</sup> We have found that in contrast to the halfsandwiches [(arene)Os(CO)(PR<sub>3</sub>)] and [(arene)OsH<sub>2</sub>(PR<sub>3</sub>)], the corresponding dicarbonyl and carbonyl dihydrido metal derivatives,  $[(arene)Os(CO)_2]$  and  $[(arene)OsH_2-$ (CO)], are photochemically more labile and thus useful starting materials for C-H activation studies both in matrix and in solution.<sup>4,6</sup> A surprising result was that the carbonyl hydrido methyl complex [(mes)OsH(CH<sub>3</sub>)(CO)], which is the photoproduct from both  $[(mes)Os(CO)_2]$  and  $[(mes)OsH_2(CO)]$  in CH<sub>4</sub> matrices<sup>4</sup> and which can also be prepared from  $[(mes)OsCl_2]_n$  or  $[(mes)Os(CO)Cl_2]$  on more conventional routes,<sup>5,7</sup> does not react with ligands such as CO or PMe<sub>3</sub> to give an acetyl(hydrido) metal derivative.

We therefore turned our attention to isocyanide osmium complexes in the hope that they offer a better chance for functionalization of the C-bound ligand. Previously, we

Scheme I





 $= C_{e}H_{e}$ 



have already reported the preparation of benzene osmium compounds  $[(C_6H_6)Os(CNR)(PMe_3)]$  which behave as metal bases and react with NH<sub>4</sub>PF<sub>6</sub>, CH<sub>3</sub>I and ClAuPPh<sub>3</sub> to form salts of the corresponding cations  $[(C_6H_6)OsX-(CNR)(PMe_3)]^+$  (X = H, CH<sub>3</sub>, AuPPh<sub>3</sub>).<sup>8</sup> The reactions of the PF<sub>6</sub> salts of these cations with excess PMe<sub>3</sub> lead to displacement of the benzene ring and formation of the octahedral complexes trans- $[OsX(CNR)(PMe_3)_4]PF_6.^8$ 

In the present paper we describe the synthesis of mesitylene(isocyanide)dichloroosmium(II) compounds and discuss the reactivity of one representative (with CNCH<sub>3</sub>) as ligand) toward organolithium and Grignard reagents. The result of the X-ray structural analysis of the carbene complex  $[(mes)Os(=C(NHCH_3)C_6H_5)(C_6H_5)_2]$ , which derives from [(mes)Os(CNCH<sub>3</sub>)Cl<sub>2</sub>] upon treatment with  $C_{6}H_{5}MgBr$ , is also reported.

#### Results

Preparation of the Dichloro Complexes [(mes)Os- $(CNR)Cl_2$ ]. The bridging halide complex  $[(mes)OsCl_2]_n$ 

0276-7333/91/2310-3278\$02.50/0 © 1991 American Chemical Society

<sup>(1) (</sup>a) Werner, H.; Werner, R. Angew. Chem. 1978, 90, 721; Angew Chem., Int. Ed. Engl. 1978, 17, 683. (b) Werner, H.; Werner, R. J. Organomet. Chem. 1980, 194, C7. (c) Werner, R.; Werner, H. Angew. Organomet. Chem. 1980, 194, C7. (c) Werner, R.; Werner, H. Angew.
Chem. 1981, 93, 826; Angew. Chem., Int. Ed. Engl. 1981, 20, 793. (d)
Werner, H.; Werner, R. Chem. Ber. 1982, 115, 3766. (e) Werner, R.;
Werner, H. Chem. Ber. 1982, 115, 3781. (f) Werner, R.; Werner, H. Chem.
Ber. 1983, 116, 2074. (g) Schubert, U.; Werner, R.; Werner, L.; Werner,
H. J. Organomet. Chem. 1983, 253, 363. (h) Werner, R.; Werner, R.;
Burschka, C. Chem. Ber. 1984, 117, 152. (i) Werner, R.; Werner, H.
Chem. Ber. 1984, 117, 161. (j) Werner, H.; Roder, K. J. Organomet.
Chem. 1985, 281, C38. (k) Werner, H.; Werner, R. Chem. Ber. 1985, 138, 4543. (l) Werner, H.; Zheng. Soc. Chem. Commun. 1985. Chem. 1985, 281, C38. (k) Werner, H.; Werner, R. Chem. Ber. 1985, 118, 4543. (l) Werner, H.; Zenkert, K. J. Chem. Soc., Chem. Commun. 1985, 1607. (m) Werner, H.; Kletzin, H.; Höhn, A.; Paul, W.; Knaup, W.; Ziegler, M. L.; Serhadli, O. J. Organomet. Chem. 1986, 306 227. (n) Werner, H.; Weinand, R.; Otto, H. J. Organomet. Chem. 1986, 307, 49. (o) Weinand, R.; Werner, H. Chem. Ber. 1986, 119, 2055. (p) Werner, H.; Roder, K. J. Organomet. Chem. 1986, 310, C51. (q) Werner, H.; Zenkert, K. J. Organomet. Chem. 1988, 345, 151. (r) Zenkert, K.; Werner, H. Chem. Ber. 1988, 121, 811. (s) Werner, H.; Kletzin, H.; Roder, K. J. Organomet. Chem. 1988, 355, 401. (t) Boder, K. Werner, H. J. Organ. Organomet. Chem. 1988, 355, 401. (t) Roder, K.; Werner, H. J. Organomet. Chem. 1989, 362, 321.

<sup>(2)</sup> Werner, H. Angew. Chem. 1983, 95, 932; Angew. Chem., Int. Ed.

 <sup>(2)</sup> werner, H. Angew. Chem. 1983, 95, 952; Angew. Chem., Int. Ea.
 Engl. 1983, 22, 927.
 (3) (a) Roder, K.; Werner, H. Angew. Chem. 1987, 99, 719; Angew.
 Chem., Int. Ed. Engl. 1987, 26, 686. (b) Roder, K.; Werner, H. Chem. Ber.
 1989, 122, 833.

<sup>(4)</sup> McCamley, A.; Perutz, R. N.; Stahl, S.; Werner, H. Angew. Chem.
(989, 101, 1721; Angew. Chem., Int. Ed. Engl. 1989, 28, 1690.
(5) Stahl, S.; Werner, H. Organometallics 1990, 9, 1876.
(6) Stahl, S. Dissertation, Universität Würzburg, 1990.
(7) Stahl, S.; Werner, H. J. Am. Chem. Soc. 1991, 113, 2944.

<sup>(8)</sup> Werner, H.; Weinand, R. Z. Naturforsch. 1983, 38b, 1518.



(1), which is either a polymer<sup>5</sup> or a dimer,<sup>9</sup> reacts with isocyanides CNR (R = CH<sub>3</sub>, t-C<sub>4</sub>H<sub>9</sub>, C<sub>6</sub>H<sub>5</sub>) under similar conditions as with CO, CH<sub>3</sub>CN, DMSO, and phosphines<sup>5,10</sup> to give the mononuclear compounds [(mes)Os(CNR)Cl<sub>2</sub>] (2–4) (Scheme I). They are orange-yellow solids that are moderately air-stable and easily soluble in polar solvents such as CH<sub>2</sub>Cl<sub>2</sub> or CHCl<sub>3</sub>. The IR spectra of 2–4 show a strong CN stretching frequency at 2140–2180 cm<sup>-1</sup>, which appears at about the same wavenumbers as that of the benzene osmium derivatives [(C<sub>6</sub>H<sub>6</sub>)Os(CNR)I<sub>2</sub>] and [(C<sub>6</sub>H<sub>6</sub>)Os(CNR)(PMe<sub>3</sub>)I]PF<sub>6</sub>.<sup>8</sup>

Reactions of Compound 2 with Organolithium and Grignard Reagents. In two preceding papers on the chemistry of mesitylene osmium carbonyl and olefin complexes we have already described that the reactions of [(mes)Os(CO)Cl<sub>2</sub>] with organolithium and Grignard reagents proceed differently.<sup>5,11</sup> This observation has also been made for 2 (which was chosen as a representative of the dichloro isocyanide compounds) as starting material. Treatment of a suspension of 2 with an ether solution of  $CH_3Li$  or with  $C_6H_5Li$  in cyclohexane/ether give the dimethyl- and diphenylosmium complexes 5 and 6 (Scheme II) in 78% and 89% yield. Both compounds are yellow air-sensitive solids, which have been characterized by elemental analysis and mass spectra. The thermal stability decreases in the order 2 > 6 > 5, which is similar to that found for the carbonyl derivatives  $[(mes)Os(CO)X_2]$ . In the <sup>1</sup>H NMR spectrum of 5, the signal of the osmiumbound  $CH_3$  protons is found upfield ( $\delta$  0.88), indicative of the expected high electron density at the metal center.

The course of the reaction of 2 with RMgI (R = CH<sub>3</sub>, C<sub>6</sub>H<sub>5</sub>) and C<sub>6</sub>H<sub>5</sub>MgBr depends on the type of Grignard reagent used. If a suspension of 2 in benzene is treated with an ether solution of CH<sub>3</sub>MgI in the molar ratio of 1:1.5, two products, [(mes)OsCH<sub>3</sub>(CNCH<sub>3</sub>)I] (7) and [(mes)Os(CNCH<sub>3</sub>)I<sub>2</sub>] (9) (Scheme III), are obtained, which can be separated by chromatographic techniques. In contrast to the behavior of [(mes)Os(CO)Cl<sub>2</sub>] toward CH<sub>3</sub>MgI, even with an excess of the Grignard reagent the formation of the dimethylosmium complex 5 cannot be observed. Compound 9 has also been prepared by metathesis from 2 and KI in almost quantitative yield.

Most interestingly, the reactions of 2 with phenylmagnesium halides give different mixture of products, depending on the type of Grignard reagent used. Whereas with  $C_6H_5MgI$  (molar ratio 1:1), in analogy to the reaction of 2 with  $CH_3MgI$ , a mixture of  $[(mes)OsC_6H_5(CNCH_3)I]$ (8) and 9 is formed, the major component upon treatment



Figure 1. SCHAKAL<sup>19</sup> drawing of complex 10.

Table I. Selected Bond Distances and Angles with Esd's of

			10					
Bond Distances, Å								
Os-C1	2.300 (5)	Os-C5	2.280 (5)	Os-C24	2.092 (5)			
Os-C2	2.321(5)	Os-C6	2.247 (5)	N-C10	1.326 (6)			
Os-C3	2.239 (5)	Os-C10	1.992 (5)	N-C17	1.473 (6)			
Os-C4	2.271 (5)	Os-C18	2.100 (4)	C10-C11	1.503 (7)			
Bond Angles, deg								
C10-Os-C18		87.0 (2)	C11-C	C11-C10-Os 1				
C10-Os-C24		87.3 (2)	C11-C	C11-C10-N 1				
C18-Os-C24		90.1 (2)	Os-C10	Os-C10-N 12				
C10-N-C17		128.0 (4)						
Torsional Angles, <sup>a</sup> deg								
CENT-Os-C10-N		-164.3 (5	5) C18-O	C18-Os-C10-C11				
CENT-Os-C10-C11		16.4 (7	7) C24-0	C24-Os-C10-N				
C18-Os-C10-N		-30.1 (7	) C24-0	s-C10-C1	1 -119.2 (6)			

<sup>a</sup>CENT = centroid of the mesitylene ring.

of 2 with  $C_6H_5MgBr$  followed by chromatography on  $Al_2O_3$ is the carbene osmium complex [(mes)Os(=C-(NHCH\_3)C\_6H\_5)(C\_6H\_5)\_2] (10). In addition, also the bromo phenyl compound [(mes)OsC\_6H\_5(CNCH\_3)Br] (11) (Scheme III) has been isolated. Although a 3.4-fold excess of the Grignard reagent has been used, the formation of the diphenyl metal derivative 6 cannot be detected.

The carbene complex 10 is an orange air-sensitive solid that is quite volatile and decomposes at ca. 100 °C. The most characteristic feature in the <sup>1</sup>H NMR spectrum is the doublet at  $\delta$  1.66 for the NHCH<sub>3</sub> protons, the splitting originating from coupling with the NH nucleus. In the <sup>13</sup>C NMR, a signal at 221.9 appears, which is assigned to the carbene carbon atom. We note that for the complex [Os(=CHCH<sub>3</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>], which has been recently described by Roper et al.,<sup>12</sup> the CHCH<sub>3</sub> signal is found at  $\delta$  227, whereas in the cationic carbene osmium derivatives

 $[(C_6H_6)Os(=CCH_2CH_2CHR'O)(PR_3)I]^+$ , which are somewhat related in structure to 10, the carbone carbon resonance appears at  $\delta$  260–285.<sup>13</sup>

Molecular Structure of 10. A single-crystal X-ray diffraction investigation of compound 10 confirms the structural proposal shown in Scheme III. The SCHAKAL drawing (Figure 1) reveals that the osmium is coordinated by the mesitylene ring, the two phenyl groups, and the aminocarbene ligand, the phenyl substituent of which is directed away from the metal-bound  $C_6H_5$  units. The bond angles C10–Os–C18, C10–Os–C24, and C18–Os–C24 (see Table I) are near to 90°, which is in agreement with the

<sup>(9)</sup> Bennett, M. A.; Weerasuria, A. M. A. J. Organomet. Chem. 1990, 394, 481.

<sup>(10)</sup> Werner, H.; Stahl, S.; Kohlmann, W. J. Organomet. Chem. 1991, 409, 285.

<sup>(11)</sup> Werner, H.; Stahl, S.; Schulz, M. Chem. Ber. 1991, 124, 707.

 <sup>(12)</sup> Gallop, M. A.; Roper, W. R. Adv. Organomet. Chem. 1986, 25, 121.
 (13) Werner, H.; Knaup, W.; Schulz, M. Chem. Ber. 1991, 124, 1121.

piano-stool arrangement of the molecule.

The Os=C bond length [1.992 (5) Å] is similar to that in  $[Os(=CHPh)(CO)(PPh_3)_2Cl_2]$  [1.94 (1) Å],<sup>12</sup> [Os(= $CF_2$  (NO) (PPh<sub>3</sub>)<sub>2</sub>Cl] [1.967 (4) Å],<sup>12</sup> and [(C<sub>6</sub>H<sub>6</sub>) Os- $(=CCH_2CH_2CH_2O)(PMe-t-Bu_2)I]^+$  [1.953 (2) Å]<sup>13</sup> and almost identical with that in  $[Os_3(CO)_9(=C(OCH_3) CH_3(\mu-H)(\mu-COCH_3)$  [1.99 (2) Å].<sup>14</sup> The C10-N distance [1.326 (6) Å] is significantly shorter than the distance C17-N [1.473 (6) Å], indicating a substantial double-bond character of the carbon-nitrogen bond. In agreement with the proposed sp<sup>2</sup> hybridization of the carbon earbon atom, the groups attached to C10 are exactly coplanar. The dihedral angle between the best planes through the atoms Os, C10, N, C11 and the mesitylene ring is 100.4 (2)°. Other torsional angles are given in Table I.

Although the mesitylene ring is exactly planar, the Os-C1 to Os-C6 distances differ by ca. 0.08 Å. The greater bond lengths (Os-C1 and Os-C2) are found opposite to C10, an observation reflecting the strong trans influence of the carbene ligand.

#### **Concluding Remarks**

The work presented in this paper has shown that interaction of the new isocyanide complexes [(mes)Os-(CNR)Cl<sub>2</sub>] with methylating and phenylating reagents can lead to the formation of either  $OsR_2$  or Os(R)X species. Whereas from 2 (R = CH<sub>3</sub>) and LiR (R = CH<sub>3</sub>, C<sub>6</sub>H<sub>5</sub>) the dimethyl- and diphenylosmium compounds 5 and 6 are obtained, treatment of the same starting material with  $CH_3MgI$  or  $C_8H_5MgX$  (X = Br, I) gives in each case at least two products. Besides the iodo methyl and phenyl and the bromo phenyl osmium derivatives (7, 8, 11), from 2 and  $C_6H_5MgBr$  a carbone complex 10 is formed, which probably originates from nucleophilic attack of the Grignard reagent at the isocyanide carbon atom followed by protonation on the nitrogen. As far as Fischer-type carbene complexes are concerned, both Darensbourg<sup>15</sup> and Casey<sup>16</sup> have previously observed that not only organolithium but also Grignard reagents can be used to transform a CO into an oxycarbene ligand. We are not aware that an analogous generation of an aminocarbene ligand from a coordinated isocyanide and RMgX has been reported.

The possibility that the solvent benzene takes part in the formation of the  $C(NHCH_3)C_6H_5$  ligand can be excluded. If the reaction of 2 with  $C_6H_5MgBr$  is performed in C<sub>6</sub>D<sub>6</sub>-ether, no deuterium is incorporated into the product. If on the other hand the aluminum oxide used for the chromatographic workup of the reaction mixture is deactivated with  $D_2O_1$ , instead of 10 the monodeuterated complex  $[(\text{mes})Os(=C(\text{NDCH}_3)C_6H_5)(C_6H_5)_2]$  (10-d<sub>1</sub>) has been isolated. The IR spectrum of  $10-d_1$  shows a characteristic ND stretching frequency at 2485 cm<sup>-1</sup>, compared with the NH absorption of 10 at 3345 cm<sup>-1</sup>.

We finally note that two carbene rhodium compounds of composition  $[(C_5Me_5)Rh(=C(NHR)R')X_2]$  (12a, X = Cl; 12b, X = Br), which are related in structure to 10, have recently been described in the literature.<sup>17,18</sup> Following their work on intermolecular oxidative addition of C-H

(16) Casey, C. P.; Brunsvold, W. R. Inorg. Chem. 1977, 16, 391.
(17) Jones, W. D.; Feher, F. J. Organometallics 1983, 2, 686.
(18) Jones, W. D.; Duttweiler, R. P.; Feher, F. J.; Hessell, E. T. New J. Chem. 1989, 13, 725.
(20) Authors F. Valler, Visite Visite Provided to Universitation of the temperature of temperatu

bonds to metal centers, Jones and co-workers prepared the (isocyanide)p-tolylrhodium derivatives  $[(C_5Me_5)Rh(p C_6H_4CH_3$ (CNR)X] (R = CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>; X = Cl, Br), which after addition of NH4X react to give the dichloro and dibromo carbene rhodium complexes 12a,b in 90% yield. The aminocarbene ligand is thought to be formed by insertion of the isocyanide into the rhodium-tolyl bond and subsequent protonation of the iminoacyl nitrogen atom with  $NH_4^+$ . A similar phenyl migration has not been observed with 6, 8, or 11 in the presence of nucleophiles, from which we conclude that the generation of the carbene ligand in 10 occurs by external attack of the Grignard reagent at the coordinated isocyanide. We are currently exploring the possibility of preparing other arene carbene osmium complexes from various isocyanide metal precursors and also of using the recently described carbonylosmium compounds [(arene)OsR<sub>2</sub>(CO)]<sup>11</sup> as starting materials.

## **Experimental Section**

All reactions were carried out under an atmosphere of argon by Schlenk tube techniques. The starting material [(mes)OsCl<sub>2</sub>]<sub>n</sub> (1) was prepared as described recently.<sup>5</sup> NMR spectra were recorded on a JEOL FX 90 Q and a Bruker AC 200 instruments, IR spectra on a Perkin-Elmer 1420 infrared spectrometer, and mass spectra on a Varian MAT CH 7 instrument.

Preparation of Complexes [(mes)Os(CNR)Cl<sub>2</sub>] (2-4). A suspension of 1 (900 mg, 1.18 mmol for n = 2) in 40 mL of dichloromethane was treated with the isocyanide ( $R = CH_3$ , 259 mg, 6.30 mmol;  $R = t - C_4 H_9$ , 480 mg, 5.75 mmol;  $R = C_6 H_5$ , 605 mg, 5.87 mmol) and stirred for 3 h at 40 °C ( $R = CH_3$ ) or for 2.5 h (R = t-C<sub>4</sub>H<sub>9</sub> and C<sub>6</sub>H<sub>5</sub>) at room temperature. The solution was filtered and the residue repeatedly washed with CH<sub>2</sub>Cl<sub>2</sub> (ca. 30 mL). The filtrate was concentrated to ca. 5-10 mL in vacuo, and ether (ca. 20 mL) was added. An orange-yellow precipitate was obtained, which was filtered off, washed with ether and hexane, and dried in vacuo, yielding 800 mg (80%) of 2, 1.02 g (93%) of 3, and 1.11 g (97%) of 4; mp of 3 166 °C dec and of 4 214 °C dec.

2: Anal. Calcd for C<sub>11</sub>H<sub>15</sub>Cl<sub>2</sub>NOs: C, 31.28; H, 3.54; N, 3.32. Found: C, 31.18; H, 3.66; N, 3.45 IR (KBr)  $\nu$ (CN) 2187 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 90 MHz) δ 5.26 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 3.78 (s, CNCH<sub>3</sub>), 2.31  $(s, C_6H_3Me_3).$ 

3: Anal. Calcd for C<sub>14</sub>H<sub>21</sub>Cl<sub>2</sub>NOs: C, 36.20; H, 4.56; N, 3.02. Found: C, 36.46; H, 4.61; N, 3.01. MS (70 eV), m/z 465 (2, M<sup>+</sup>), 382 (15,  $M^+$  – CNC<sub>4</sub>H<sub>9</sub>); IR (KBr)  $\nu$ (CN) 2162 cm<sup>-1</sup>; <sup>1</sup>H NMR  $(CDCl_3, 90 \text{ MHz}) \delta 5.23 \text{ (s, } C_6H_3Me_3), 2.30 \text{ (s, } C_6H_3Me_3), 1.54 \text{ (s,}$  $t-C_4H_9$ ).

4: Anal. Calcd for C<sub>16</sub>H<sub>17</sub>Cl<sub>2</sub>NOs: C, 39.67; H, 3.54; N, 2.89. Found: C, 39.88; H, 3.83; N, 3.01. MS (70 eV), m/z 485 (0.3, M<sup>+</sup>), 120 (34, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub><sup>+</sup>); IR (KBr)  $\nu$ (CN) 2142 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 90 MHz)  $\delta$  7.39 (m, C<sub>6</sub>H<sub>5</sub>), 5.34 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 2.39 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>).

Preparation of [(mes)Os(CH<sub>3</sub>)<sub>2</sub>(CNCH<sub>3</sub>)] (5). A suspension of 2 (200 mg, 0.47 mmol) in 10 mL of benzene was treated at 5 °C with 0.75 mL of a 1.25 M solution of CH<sub>3</sub>Li in ether (0.94 mmol). After stirring for 2.5 h, a small amount of  $Al_2O_3$  (ca. 100 mg) was added and the solvent was removed in vacuo. The residue was dissolved in 3 mL of benzene and the solution chromatographed on  $Al_2O_3$  (basic, activity grade III). With hexane a yellow fraction was eluted from which after removing the solvent a yellow microcrystalline solid was isolated, yield 139 mg (78%), mp 76 °C dec. Anal. Calcd for C<sub>13</sub>H<sub>21</sub>NOs: C, 40.92; H, 5.55; N, 3.67. Found: C, 41.02; H, 5.46; N, 3.73. MS (70 eV), m/z 383 (14, M<sup>+</sup>), 368 (13, M<sup>+</sup> – CH<sub>3</sub>), 312 (1, (mes)Os<sup>+</sup>); IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu$ (CN) 2102 cm<sup>-1</sup>; <sup>1</sup>H NMR ( $C_{6}D_{6}$ , 90 MHz)  $\delta$  4.38 (s,  $C_{6}H_{3}Me_{3}$ ), 2.68 (s, CNCH<sub>3</sub>), 1.96 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 0.88 (s, OsCH<sub>3</sub>).

**Preparation of**  $[(mes)Os(C_6H_5)_2(CNCH_3)]$  (6). A suspension of 2 (100 mg, 0.24 mmol) in 5 mL of benzene was treated at 5 °C with 0.24 mL of a 2.0 M solution of  $C_6H_5Li$  in cyclohexane/ether. After stirring for 30 min, the reaction mixture was worked up as described for 5. With hexane an orange fraction was eluted, which contained a small amount of 10 (9 mg, 6%). With dichloromethane, a yellow fraction was eluted from which after removing the solvent a yellow microcrystalline solid was isolated, yield 108 mg (89%); mp 129 °C dec. Anal. Calcd for C<sub>23</sub>H<sub>25</sub>NOs: C, 54.63;

<sup>(14)</sup> Jensen, C. M.; Lynch, T. J.; Knobler, C. B.; Kaesz, H. D. J. Am. Chem. Soc. 1982, 104, 4679.

<sup>(15)</sup> Darensbourg, D. J.; Darensbourg, M. Y. Inorg. Chim. Acta 1971, 5,247

<sup>(19)</sup> Author: E. Keller, Kristallographisches Institut der Universität Freiburg, W-7800 Freiburg, FRG.

H, 4.98; N, 2.77. Found: C, 54.49; H, 4.93; N, 2.67. MS (70 eV), m/z 507 (36, M<sup>+</sup>), 466 (2, M<sup>+</sup> - CNCH<sub>3</sub>), 430 (2, M<sup>+</sup> - C<sub>6</sub>H<sub>5</sub>), 389 (3, (mes)OsC<sub>6</sub>H<sub>5</sub><sup>+</sup>), 353 (100, M<sup>+</sup> - 2C<sub>6</sub>H<sub>5</sub>); IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu$ (CN) 2123 cm<sup>-1</sup>; <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 90 MHz)  $\delta$  7.87 (m, 4 H, C<sub>6</sub>H<sub>5</sub>), 7.19 (m, 6 H, C<sub>6</sub>H<sub>5</sub>), 4.41 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 2.47 (s, CNCH<sub>3</sub>), 1.65 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>).

**Preparation of [(mes)OsCH<sub>3</sub>(CNCH<sub>3</sub>)I] (7) and [(mes)-Os(CNCH<sub>3</sub>)I<sub>2</sub>] (9).** A suspension of 2 (150 mg, 0.36 mmol) in 5 mL of benzene was treated at 5 °C with 2.2 mL of a 0.25 M solution of CH<sub>3</sub>MgI in ether (0.55 mmol). After warming to room temperature, the reaction mixture was stirred for 15 min, and then a small amount of Al<sub>2</sub>O<sub>3</sub> (ca. 100 mg) was added (to destroy the excess of CH<sub>3</sub>MgI). The solvent was removed in vacuo and after the residue was dissolved in 3 mL of C<sub>6</sub>H<sub>6</sub>/CH<sub>2</sub>Cl<sub>2</sub> (1:1), the solution was chromatographed on Al<sub>2</sub>O<sub>3</sub> (basic, activity grade III). With benzene, an orange fraction and, subsequently, with dichloromethane a red fraction were eluted. Both fractions were brought to dryness in vacuo to give an orange-yellow microcrystalline solid of 7 (yield 41 mg, 23%) and a red microcrystalline solid of 9 (yield 163 mg, 75%), mp of 7 148 °C dec.

7: Anal. Calcd for  $C_{12}H_{18}INO_{5}$ : C, 29.21; H, 3.68; N, 2.84. Found: C, 29.30; H, 3.61; N, 2.63. MS (70 eV), m/z 495 (7, M<sup>+</sup>), 480 (13, M<sup>+</sup> – CH<sub>3</sub>), 439 (15, (mes)OsI<sup>+</sup>); IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu$ (CN) 2140 cm<sup>-1</sup>; <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 90 MHz)  $\delta$  4.42 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 2.59 (s, CNCH<sub>3</sub>), 2.00 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 1.73 (s, OsCH<sub>3</sub>).

9: Anal. Calcd for  $C_{11}H_{15}I_2NOs$ : C, 21.83; H, 2.50; N, 2.31. Found: C, 22.43; H, 2.72; N, 2.48. IR (KBr)  $\nu$ (CN) 2173 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 90 MHz)  $\delta$  5.36 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 3.92 (s, CNCH<sub>3</sub>), 2.56 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>).

A second procedure for the preparation of 9 is as follows: A solution of 2 (100 mg, 0.24 mmol) in 15 mL of acetone was treated with an excess of KI (ca. 300 mg). After it was stirred for 3 h at 60 °C, the solution was cooled and then the solvent was removed in vacuo. The residue was extracted twice with 6 mL of  $CH_2Cl_2$  each, and the combined extracts were concentrated to ca. 3 mL. After addition of hexane, a dark red precipitate was obtained, which was filtered off, repeatedly washed with hexane, and dried in vacuo, yield 129 mg (90%).

Preparation of [(mes)OsC<sub>6</sub>H<sub>5</sub>(CNCH<sub>3</sub>)I] (8). A suspension of 2 (105 mg, 0.25 mmol) in 10 mL of benzene was treated at 5 °C with an equimolar amount of C<sub>6</sub>H<sub>5</sub>MgI in ether. After warming to room temperature and stirring for 30 min, some Al<sub>2</sub>O<sub>3</sub> was added. The solvent was removed in vacuo, the residue was dissolved in 2 mL of benzene, and the solution was chromatographed on  $Al_2O_3$  (basic, activity grade III). With hexane, an orange fraction was eluted, which contained a small quantity (13 mg, 9%) of 10. With  $C_6H_6/CH_2Cl_2$  (5:1) a yellow fraction was eluted from which after removing the solvent a yellow microcrystalline solid 8 (41 mg, 30%) was obtained. Finally, a red fraction was eluted from which 9 (88 mg, 58%) was isolated; mp of 8 176 °C dec. Anal. Calcd for C<sub>28</sub>H<sub>31</sub>INOs: C, 36.76; H, 3.63; N, 2.52. Found: C, 36.76; H, 3.68; N, 2.72. MS (70 eV), m/z 557 (17, M<sup>+</sup>), 516 (0.3, M<sup>+</sup> - $CNCH_3$ , 480 (5, M<sup>+</sup> - C<sub>6</sub>H<sub>5</sub>), 439 (9, (mes)OsI<sup>+</sup>), 430 (20, M<sup>+</sup> -I), 389 (14, (mes)OsC<sub>6</sub>H<sub>5</sub><sup>+</sup>), 353 (1, (mes)OsCNCH<sub>3</sub><sup>+</sup>); IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu$ (CN) 2160 cm<sup>-1</sup>; <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 90 MHz)  $\delta$  7.91 (m, 2 H, C<sub>6</sub>H<sub>5</sub>), 7.14 (m, 3 H,  $C_{6}H_{5}$ ), 4.42 (s,  $C_{6}H_{3}Me_{3}$ ), 2.45 (s,  $CNCH_{3}$ ), 1.82 (s,  $C_6H_3Me_3$ ).

**Preparation of [(mes)Os(=C(NHCH<sub>3</sub>)C<sub>6</sub>H<sub>5</sub>)(C<sub>6</sub>H<sub>6</sub>)<sub>2</sub>] (10) and [(mes)OsC<sub>6</sub>H<sub>6</sub>(CNCH<sub>3</sub>)Br] (11). A suspension of 2 (105 mg, 0.25 mmol) in 10 mL of benzene was treated with 3.6 mL of a 0.25 M solution of C<sub>6</sub>H<sub>6</sub>MgBr in ether (0.9 mmol). After stirring for 30 min, a small amount of Al<sub>2</sub>O<sub>3</sub> (ca. 100 mg) was added and the solvent was removed in vacuo. The residue was dissolved in 2 mL of benzene and the solution was chromatographed on Al<sub>2</sub>O<sub>3</sub> (basic, activity grade III). With benzene/hexane (1:1) an orange fraction was eluted, which was brought to dryness in vacuo. Recrystallization of the residue from hexane gave orange crystals of 10, yield 68 mg (71%), mp 99 °C dec. With dichloromethane, a second orange fraction was eluted from which after removing the solvent an orange microcrystalline powder of 11 was obtained, yield 29 mg (23%), mp 188 °C dec.** 

10: Anal. Calcd for  $C_{29}H_{31}NOs$ : C, 59.67; H, 5.35; N, 2.40. Found: C, 59.62; H, 5.55; N, 2.46. MS (70 eV), m/z 585 (3, M<sup>+</sup>), 507 (18, M<sup>+</sup> - C<sub>6</sub>H<sub>6</sub>), 353 (11, (mes)Os(CNCH<sub>3</sub>)<sup>+</sup>); IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu$ (NH) 3345 cm<sup>-1</sup>; <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 200 MHz)  $\delta$  7.82–6.84 (m, C<sub>6</sub>H<sub>5</sub>), 4.47 (s, C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 1.66 (d, J(HH) = 5.3 Hz, NCH<sub>3</sub>), 1.38 (s,

Table II. Crystallographic Data for 10

formula	C <sub>29</sub> H <sub>31</sub> NOs
formula wt	583.77
cryst size, mm	$0.15 \times 0.4 \times 0.5$
cryst syst	monoclinic
space group	$P2_1/n$
cell dimens determn	23 reflecns, $12^{\circ} < \theta < 15^{\circ}$
a, Å	12.402 (3)
b, Å	14.237 (2)
c, Å	13.622 (2)
$\beta$ , deg	91.54 (1)
V, Å <sup>3</sup>	2404.3
Z	4
$d_{\rm calcd}, {\rm g \ cm^{-3}}$	1.61
diffractometer	Enraf-Nonius CAD 4
radiatn (graphite monochromated)	Mo Kα (0.70930Å)
temp, °C	$20 \pm 1$
$\mu,  {\rm cm}^{-1}$	53.2
scan method	$\omega/2\theta$
$2\theta$ (max), deg	46
total no. of rfins scanned	3682
no. of unique rflns	3320
no, of obsd rflns $(F_0 > 3\sigma(F_0))$	2618
no, of params refined	280
R	0.028
R <sub>w</sub>	0.031
rfln/param ratio	9.35
residual electron density, e Å <sup>-3</sup>	+0.67/-0.42

 Table III. Positional Parameters and Their Estimated

 Standard Deviations<sup>a</sup>

atom	x	У	z	β, Ų
Os	0.23643 (2)	0.19085 (2)	0.00061 (2)	3.110 (5)
Ν	0.2175 (5)	0.2397 (5)	0.2084(5)	4.2 (2)
C1	0.2561(7)	0.1859 (7)	-0.1667 (6)	4.9 (2)
C2	0.1666(7)	0.1322 (6)	-0.1466 (6)	4.4 (2)
C3	0.1766 (7)	0.0604 (6)	-0.0744 (6)	4.4 (2)
C4	0.2796 (7)	0.0390 (6)	-0.0305 (6)	4.4 (2)
C5	0.3685(7)	0.0944 (6)	-0.0547 (6)	4.4 (2)
C6	0.3576 (7)	0.1712 (7)	-0.1176 (6)	4.5 (2)
C7	0.4532 (7)	0.2304 (7)	-0.1402 (7)	5.9 (2)
C8	0.0569 (8)	0.1487 (8)	-0.1967 (8)	6.9 (3)
C9	0.2905 (8)	-0.0466 (6)	0.0350 (6)	5.3 (2)
C10	0.2684 (6)	0.1864 (6)	0.1445 (5)	3.6 (2)
C11	0.3525 (6)	0.1238 (6)	0.1915 (5)	3.6 (2)
C12	0.4605 (6)	0.1311 (6)	0.1693 (6)	4.2 (2)
C13	0.5359 (7)	0.0714 (7)	0.2137 (7)	5.6 (2)
C14	0.5025 (8)	0.0057 (8)	0.2800 (7)	6.4 (2)
C15	0.3958 (9)	-0.0028 (7)	0.3020 (7)	6.1 (3)
C16	0.3206 (7)	0.0557 (6)	0.2585 (6)	4.7 (2)
C17	0.2455 (7)	0.2553 (7)	0.3128 (6)	5.6 (2)
C18	0.0779 (5)	0.2275 (6)	0.0357 (6)	3.4 (2)
C19	0.0231 (6)	0.3104 (7)	0.0027 (6)	4.3 (2)
C20	-0.0835 (6)	0.3247 (7)	0.0220 (6)	5.3 (2)
C21	-0.1416 (6)	0.2640 (7)	0.0758 (6)	5.0 (2)
C22	-0.0923 (6)	0.1842 (7)	0.1107 (6)	4.9 (2)
C23	0.0155 (6)	0.1671 (6)	0.0919 (6)	4.4 (2)
C24	0.2811 (6)	0.3324 (5)	0.0054 (6)	3.8 (2)
C25	0.2512 (7)	0.3939 (6)	-0.0745 (7)	4.8 (2)
C26	0.2861(7)	0.4868 (7)	-0.0784 (8)	5.8 (2)
C27	0.3531 (7)	0.5241 (7)	-0.0073 (8)	6.1 (3)
C28	0.3840 (7)	0.4674 (7)	0.0718 (8)	5.7 (2)
C29	0.3494 (6)	0.3754 (7)	0.0775 (7)	4.9 (2)

<sup>a</sup> Anisotropically refined atoms are given in the form of the isotropic equivalent displacement parameter defined as  $B_{eq} = \frac{4}{3^{e}} [a^2B(1,1) + b^2B(2,2) + c^2B(3,3) + ab(\cos \gamma)B(1,2) + ac(\cos \beta)B(1,3) + bc(\cos \alpha)B(2,3)].$ 

 $\rm C_6H_3Me_3),$  signal of the NH proton not observed;  $^{13}\rm C$  NMR (C<sub>6</sub>D<sub>6</sub>, 50.3 MHz)  $\delta$  221.9 (s, Os=C), 155.0, 152.7, 143.9, 129.0, 127.4, 127.2, 126.8, 123.3, 121.2 (all s, C<sub>6</sub>H<sub>5</sub>), 99.9 (s, CMe of C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 89.0 (s, CH of C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>), 36.3 (s, NCH<sub>3</sub>), 17.6 (s, CCH<sub>3</sub> of C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>).

11: Anal. Calcd for  $C_{17}H_{20}BrNOs$ : C, 40.16; H, 3.96; N, 2.75. Found: C, 40.53; H, 4.27; N, 2.87. MS (70 eV), m/z 509 (0.3, M<sup>+</sup>), 120 (33,  $C_{6}H_{3}Me_{3}^{+}$ ); IR ( $CH_{2}Cl_{2}$ )  $\nu$ (CN) 2167 cm<sup>-1</sup>; <sup>1</sup>H NMR ( $C_{6}D_{6}$ , 90 MHz)  $\delta$  7.86 (m, 2 H of  $C_{6}H_{5}$ ), 7.18 (m, 3 H of  $C_{6}H_{5}$ ), 4.47 (s,  $C_{6}H_{3}Me_{3}$ ), 2.50 (s, CNCH<sub>3</sub>), 1.88 (s,  $C_{6}H_{3}Me_{3}$ ). X-ray Structural Analysis of 10. Single crystals were grown from hexane at -15 °C. Crystal data collection parameters are summarized in Table II. Intensity data were corrected for Lorentz and polarization effects. An empirical absorption correction was applied; the minimal transmission was 81%. The structure was solved by direct methods (SHELXS-36). Atomic coordinates (see Table III) and anisotropic thermal parameters of the non-hydrogen atoms were refined by full-matrix least squares. The positions of all hydrogen atoms were calculated according to ideal geometry and were refined by the riding method. For other details see Table II.

Acknowledgment. We thank the Deutsche Forschungsgemeinschaft (SFB 347) and the Fonds der Chemischen Industrie for financial support, the Fonds in particular for a Doktorandenstipendium for U. Wecker and M. Schulz. We also gratefully acknowledge support by Dr. G. Lange and F. Dadrich (mass spectra), Dr. C. Burschka (X-ray analysis), T. Daniel and D. Schneider (NMR spectra), U. Neumann, R. Schedl, and C. P. Kneis (elemental analyses), and DEGUSSA AG (chemicals).

Supplementary Material Available: Drawing of complex 10 showing the atom numbering scheme and tables of bond distances, bond angles, least-squares planes, positional parameters, and general displacement parameter expressions (11 pages); tables of observed and calculated structure factors (14 pages). Ordering information is given on any current masthead page.

# Cluster Chemistry. 69.<sup>1</sup> Unusual Transformation of Triarylphosphine Ligands on a Pentanuclear Ruthenium Cluster. X-ray Structures of Two Phases of $Ru_5(\mu_4$ -PPh)( $\mu_3$ -PhC<sub>2</sub>Ph)( $\mu$ -PPh<sub>2</sub>)<sub>2</sub>(CO)<sub>10</sub>

Michael I. Bruce\* and Michael J. Liddell

Jordan Laboratorles, Department of Physical and Inorganic Chemistry, University of Adelaide, Adelaide, South Australia 5001, Australia

Brian W. Skelton and Allan H. White

Department of Physical and Inorganic Chemistry, University of Western Australia, Nedlands, Western Australia 6009, Australia

Received March 18, 1991

The reaction between  $\operatorname{Ru}_5(\mu_5-C_2\operatorname{PPh}_2)(\mu-\operatorname{PPh}_2)(\operatorname{CO})_{13}$  and  $\operatorname{PPh}_3$  gave a monosubstituted complex, which in refluxing  $\operatorname{CH}_2\operatorname{Cl}_2$  was transformed into  $\operatorname{Ru}_5(\mu_4-\operatorname{PPh})(\mu_3-\operatorname{PhC}_2\operatorname{Ph})(\mu-\operatorname{PPh}_2)_2(\operatorname{CO})_{10}$  (4, Ar = Ph). This complex contains a distorted-octahedral PRu<sub>5</sub> cluster with the PPh<sub>2</sub> groups bridging basal Ru-Ru edges, together with a  $\operatorname{C}_2\operatorname{Ph}_2$  ligand spanning an Ru<sub>3</sub> face. A similar reaction with P(tol)<sub>3</sub> (tol =  $\operatorname{C}_6\operatorname{H}_4\operatorname{Me}_p$ ) gave  $\operatorname{Ru}_5(\mu_4-\operatorname{PPh})(\mu_3-\operatorname{PhC}_2\operatorname{Ar})(\mu-\operatorname{PPh}_2)(\mu-\operatorname{PAr}_2)(\operatorname{CO})_{10}$  (4, Ar = tol). The net reaction occurring on the Ru<sub>5</sub> cluster is between PAr<sub>3</sub> (Ar = Ph, tol) and  $\operatorname{C}_2\operatorname{PPh}_2$  to form PPh, PAr<sub>2</sub>, and PhC<sub>2</sub>Ar ligands. The molecular structure of 4 (Ar = Ph) was determined from X-ray studies on solvent-free and solvated modifications. Ru<sub>5</sub>( $\mu_4$ -PPh)( $\mu_3$ -PhC<sub>2</sub>Ph)( $\mu$ -PPh<sub>2</sub>)\_2(\operatorname{CO})\_{10}: triclinic, space group  $\operatorname{PI}_1$ , a = 17.282 (4) Å, b = 13.558 (2) Å, c = 13.498 (4) Å,  $\alpha = 61.94$  (1)°,  $\beta = 81.06$  (2)°,  $\gamma = 85.03$  (2)°, Z = 2, R = 0.037, R' = 0.036 for 6084 observed reflections. Ru<sub>5</sub>( $\mu_4$ -PPh)( $\mu_3$ -PhC<sub>2</sub>Ph)( $\mu$ -PPh<sub>2</sub>)\_2(\operatorname{CO})\_{10}·CH<sub>2</sub>Cl<sub>2</sub>: triclinic, space group PI, a = 21.399 (10) Å, b = 12.658 (15) Å, c = 12.447 (8) Å,  $\alpha = 59.00$  (7)°,  $\beta = 79.20$  (4)°,  $\gamma = 74.22$  (8)°, Z = 2, R = 0.067, R' = 0.085 for 5442 observed reflections.

### Introduction

We have recently shown that the preferred sites of CO substitution by tertiary phosphines or phosphites in  $\operatorname{Ru}_5(\mu_5\text{-}C_2\operatorname{PPh}_2)(\mu\operatorname{-PPh}_2)(\operatorname{CO})_{13}$  (1; Scheme I) are at Ru(3) (thermally induced reactions) or at Ru(4) (in Me<sub>3</sub>NO-facilitated reactions) to give isomers of  $\operatorname{Ru}_5(\mu_5\text{-}C_2\operatorname{PPh}_2)(\mu\operatorname{-PPh}_2)(\operatorname{CO})_{12}(\operatorname{PR}_3)$  (2a, 2c).<sup>1</sup> In studies with P(OEt)<sub>3</sub>, no subsequent modification of the group 15 ligand occurred; the product obtained with PMe<sub>2</sub>Ph was noted as being unstable, but the decomposition products were not characterised. We have now found that facile rearrangement reactions of triarylphosphines occur with 1; these are described below.

## **Results and Discussion**

Two methods of substituting CO ligands in 1 by  $PPh_3$ were examined. The first method, using  $Me_3NO$  to remove

Scheme I. Reactions of  $Ru_5(\mu_8-C_2PPh_2)(\mu-PPh_2)(CO)_{13}$  (1) with P(OEt)<sub>3</sub>



a CO ligand, permitted the isolation of dark brown crystalline  $\operatorname{Ru}_5(\mu_5-C_2PPh_2)(\mu-PPh_2)(CO)_{12}(PPh_3)$  (3, Ar = Ph). In the FAB mass spectrum, a molecular ion was found at

<sup>(1)</sup> Part 68: Bruce, M. I.; Liddell, M. J.; Nicholson, B. K.; Skelton, B. W.; Tiekink, E. R. T.; White, A. H. J. Organomet. Chem., in press.