COMMUNICATION

www.rsc.org/obc

B

Tandem RCM-Pauson–Khand reaction for access to tricycles in one step †

Marta Rosillo, Luis Casarrubios, ‡ Gema Domínguez and Javier Pérez-Castells* Departamento de Química, Facultad de CC. Experimentales y de la Salud,

Universidad San Pablo-CEU, Boadilla del Monte, 28668-Madrid, Spain. *E-mail: jpercas@ceu.es*

Received 28th February 2003, Accepted 26th March 2003 First published as an Advance Article on the web 4th April 2003

Starting from conveniently designed dienvnes complexed to cobalt, a tandem RCM-intramolecular Pauson-Khand reaction yields tricyclic compounds. The methodology allows the synthesis of 6,5,5 and 7,5,5 systems.

An important strategy for the synthesis of natural products consists of applying transition metal catalyzed or mediated reactions to simple starting materials to obtain polycyclic scaffolds that can be further functionalized. Two powerful reactions that help in this aim are the RCM¹ and the Pauson-Khand² reaction. By means of the RCM reaction cycles going from 5 to 8 members, macrocycles are readily obtained. These cycloalkenes can be part of Pauson-Khand reactions. If an alkynyl moiety is suitably situated the RCM process followed by the intramolecular Pauson-Khand reaction would give a tricyclic structure, bearing a cyclopentenone. One problem would be the competition between RCM and enyne metathesis, but this could be avoided if the triple bond is complexed to cobalt prior to the RCM process. Green³ and Young⁴ have recently shown the feasibility of carrying out an RCM in structures containing cobalt hexacarbonyl-alkyne complexes.

This communication shows our first results in the tandem RCM-intramolecular Pauson-Khand reaction on acyclic dienes bearing a complexed triple bond. Cobalthexacarbonyl acts first as a protecting group and second as the mediator for the PKR. The methodology allows, the formation of tricyclic 6,5,5 and 7,5,5 structures including, in some examples, oxygen or nitrogen (Scheme 1). Related tricyclic structures via Pauson-Khand reactions for the total synthesis of natural products such as dendrobine,⁵ asteriscanolide⁶ and epoxydictymene⁷ have been obtained by other groups.



Our first aim was the synthesis of substrates containing a propargyl ether moiety, which normally gives good results in Pauson-Khand reactions. Thus, starting from 5-hexen-1-ol and from ε-caprolactone we carried out the synthesis of compounds 4 and 5, respectively, following the reactions depicted in Scheme 2. Thus, compound 4 was obtained in 54% overall yield after three steps, whereas compound 5 required seven steps and was obtained in 38% overall yield.

‡ Present address: Centro de investigación Lilly S. A. Avda. de la industria 30. 28108 Alcobendas, Madrid, Spain



Scheme 2

Compound 4 was submitted to reaction with 5% Grubbs' catalyst (Cy₃P)₂Cl₂Ru=CHPh, to explore the competition among the different possible metathesis processes. We obtained a complex crude mixture which could not be separated and that contained at least three metathesis products. Thus, we complexed compound 4 with cobalt and submitted the cobalthexacarbonyl-alkyne complex 6 to the same conditions. The reaction did not take place unless the cobalt complex was previously purified by means of a silica column chromatography. In that case a clean conversion to 7 was observed, obtaining this product in 90% yield after purification (Scheme 3).



The next step was to effect the Pauson-Khand reaction on compound 7. We obtained the desired compound 8⁸ as only one diastereomer in 45% yield using our reported procedure.9 On

10.1039/b302277d

ö

[†] Electronic supplementary information (ESI) available: spectral data and ¹H-NMR and ¹³C-NMR spectra for compounds 8, 10, 11, 14a and 14b. See http://www.rsc.org/suppdata/ob/b3/b302277c

the other hand, we carried out the reaction of complex 6 with Grubbs' catalyst and when TLC showed the disappearance of starting material we added the promoters for the Pauson-Khand reaction. We compared three reaction conditions: molecular sieves and trimethylamine N-oxide (TMANO) as copromoters in toluene at rt (method A),9 TMANO or NMO in DCM at rt (methods B and C), and Sugihara's procedure using cyclohexylamine in refluxing DCM (method D).¹⁰ We obtained the same diastereomer of 8 in 47% 55%, 81% and 15% yield, respectively. Compound 5 was reacted with $Co_2(CO)_8$ and the pure resulting complex 9 (79%) was treated under conditions A, B and C. In this case the RCM was not complete after 18 hours. NMR analysis of an aliquot of the reaction showed a 75% conversion. Addition of more ruthenium catalyst did not improve the conversion.¹¹ Thus we proceeded with the Pauson-Khand reaction and we obtained a 55% (method A) and 70% (methods B,C) yield of 10 as a mixture of diastereomers that could not be separated, along with 20% 19% and 11% of compound 11, which was obtained as a single diastereomer, respectively.¹² The formation of a seven membered ring does not allow complete diastereoselection in the Pauson-Khand process (Scheme 4).



Scheme 4

The next step consisted of the synthesis of a nitrogen containing substrate. Thus, from 6-heptenal we carried out the synthesis of propargylamine 12 by reaction of the corresponding propargyl imine with a vinyl Grignard. Compound 12, obtained in moderate yield, was protected as the Boc derivative and complexed with cobalt to give complex 13. The tandem reaction of this substrate using methods A, B and C formed the desired tricyclic product 14 as a (3:2), (4:3) and (2:1) mixture of diastereomers, respectively,¹³ which were separated and assigned with NOE experiments (Scheme 5).¹⁴ The major isomer was in this case the *cis, trans* compound 14a. Again, the formation of the seven membered ring has led to a mixture of diastereomers.

In summary a new access to tricyclic structures by means of a RCM-Pauson–Khand process is described. The methodology is



currently being extended to new precursors of systems with different ring sizes.

Acknowledgements

Grants from the DGES (MCYT-Spain, PB98-0053) and the Universidad San Pablo-CEU (2/01) have supported this work. M. R. C. acknowledges the DGES for predoctoral fellowship.

Notes and references

- Recent reviews on RCM reactions: (a) T. M. Trnka and R. H. Grubbs, Acc. Chem. Res., 2001, 34, 18–29; (b) A. Fürstner, Angew. Chem., Int. Ed., 2000, 39, 3012–3043; (c) M. E. Maier, Angew. Chem., Int. Ed., 2000, 39, 2073–2077; (d) S. K. Armstrong, J. Chem. Soc., Perkin Trans. 1, 1998, 371–388; (e) R. H. Grubbs and S. Chang, Tetrahedron, 1998, 54, 4413–4450.
- Recent reviews on the Pauson–Khand reaction: (a) T. Sugihara, M. Yamaguchi and M. Nishizawa, *Chem. Eur. J.*, 2001, 7, 1589– 1595; (b) K. M. Brummond and J. L. Kent, *Tetrahedron*, 2000, 56, 3263–3282; (c) K. K. Chung, *Coord. Chem. Rev.*, 1999, 188, 297–341.
 J. R. Green, *Synlett*, 2001, 353–356.
- 4 J. A. Burlison, J. M. Gray and D. G. J. Young, *Tetrahedron Lett.*, 2001, 42, 5363–5365.
- 5 J. Cassayre and S. Z. Zard, J. Am. Chem. Soc., 1999, 121, 6072-6073.
- 6 M. E. Krafft, Y. Y. Cheung and K. A. Abboud, J. Org. Chem., 2001, 66, 7443–7448.
- 7 T. F. Jamison, S. Shambayati, W. E. Crowe and S. L. Schreiber, J. Am. Chem. Soc., 1997, 119, 4353–4363.
- 8 This product was described previously: S. Shambayati, W. E. Crowe and S. L. Schreiber, *Tetrahedron Lett.*, 1990, **31**, 5289–5292.
- 9 L. Pérez-Serrano, L. Casarrubios, G. Domínguez and J. Pérez-Castells, *Org. Lett.*, 1999, 1, 1187–1188.
- 10 T. Sugihara, M. Yamada, H. Ban, M. Yamaguchi and C. Kaneko, *Angew. Chem., Int. Ed. Engl.*, 1997, 36, 2801–2802.
- 11 We also tried the reaction with a second generation catalyst $(Cy_3P)(NHC)Cl_2Ru=CHPh:$ M. Scholl, S. Ting, W. C. Lee and R. H. Grubbs, *Org. Lett.*, 1999, 1, 953–956 This complex did not give any conversion after 20 hours even when heating the reaction at 40 °C. Starting material was recovered.
- 12 Compound 11 arises from the Pauson–Khand reaction of 9.
- 13 Calculated by integration of the signals corresponding to the olefinic proton in the ¹H-NMR spectrum of the crude mixture.
- 14 Compound 14a: no NOE increments in H_{8a} are observed when irradiating H_{8b} or H_{4a} . Compound 14b: when irradiating H_{8b} a NOE increment of 8% was observed in the signal of H_{8a} and a 12% increment was observed in the signal of H_{4a} . The assignment of the signals was done with a COSY experiment.

Published on 04 April 2003. Downloaded by University of California - Santa Cruz on 30/10/2014 23:55:16.