Reaction Mechanisms

the selectivity and reactivity of Cu^I enolate complexes

containing or lacking phen ligands in reactions with different iodoarenes, reactions with substrates that probe for aryl

radical intermediates, and DFT calculations all indicate that

the active species is a C-bound Cu^I enolate complex ligated by

a single phen unit, and that this intermediate reacts with the

iodoarene to generate a Cu^{III} intermediate without the intermediacy of aryl radicals. The relative reactivity of the different enolates correlates with the energy differences between the observed species and the reactive C-bound enolates, along with the barriers to oxidative addition of ArI

The syntheses of phen-ligated Cu^{I} complexes containing the enolates from β -dicarbonyl compounds and cyanoacetate,

as well as the less stabilized enolates from α phenylesters, is

outlined in Scheme 1. Treatment of copper tert-butyloxide

with 2 equivalents of phen in benzene, with subsequent

addition of 2,4-pentanedione, ethyl acetoacetate, diethyl

malonate, dimethyl malonate, ethyl cyanoacetate, ethyl phenylacetate, and methyl phenylacetate formed the Cu^I

to form aryl-Cu^{III} enolate intermediates.

complexes **1–7** in 59–82% yield.^[8]

RT, 5 min

Copper(I) Enolate Complexes in α-Arylation Reactions: Synthesis, Reactivity, and Mechanism**

Zheng Huang and John F. Hartwig*

The α -arylation of enolates has become a widely used method to prepare α -aryl carboxylic acid and ketone derivatives that comprise important classes of biologically active compounds. Palladium-catalyzed α -arylation of carbonyl compounds has been developed over the past decade, but the coppermediated α -arylation of activated methylene compounds has been known for nearly a century [Eq. (1)].^[1-2] Recently,

$$X + R^{1} R^{2} \xrightarrow{Cu(I), \text{ ligand}}_{\text{base}} R^{1} R^{2} = CO_{2}R, \quad (1)$$

significant improvements in the scope and reaction conditions of these copper-catalyzed Hurtley-type reactions have been achieved using a combination of copper complexes with ancillary ligands, such as 2-phenylphenol,^[3] chelating Schiff bases,^[4] L-prolines,^[5] and 2-picolinic acid.^[6]

Despite the long history of Hurtley couplings, the mechanism of this process has been little explored. Setsune et al. proposed that the active copper species in the coupling of aryl halides with diethyl malonate is a Cu¹ enolate generated in situ and that the Cu¹ species reacts with the aryl halide by electron transfer.^[7] Buchwald et al. found that CuI and 2-phenylphenol catalyzed the arylation of diethyl malonate, but this system did not catalyze the arylation of the cyclic isopropylidene malonate and 1,3-cyclopentanedione. Accordingly, they proposed that a bidentate coordination mode through two O atoms was required for the arylation reactions to occur.^[3] Thus, mechanistic proposals have been made, but no explicit structural information on the copper enolate complexes that could be intermediates in the Hurtley couplings has been reported.

We report the synthesis of a series of Cu^{I} enolate complexes supported by 1,10-phenanthroline (phen). Their structures consist of an unusual combination of one cationic Cu^{I} center ligated by two phen ligands and one free, anionic enolate unit. Kinetic data on the stoichiometric reactions of the Cu^{I} enolate complexes with iodobenzene, comparisons of

[*] Dr. Z. Huang, Prof. Dr. J. F. Hartwig ^[+]					
Department of Chemistry					
University of Illinois Urbons Champaign					

University of Illinois Urbana-Champaign, Urbana, IL 61801 (USA) [⁺] Current address: University of California, Department of Chemistry 718 Latimer Hall, Berkeley, CA 94720 (USA) E-mail: jhartwig@berkeley.edu

- [**] We thank the NIH (GM-58108) for support of this work.
- Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201106719.

e arylation of the clopentanedione. 2 equivate coordination for the arylation n on the copper es in the Hurtley 1 74% 2 equiv

CuO/Bu



Scheme 1. Synthesis of Cu¹ enolate complexes 1-7.

All complexes were characterized by IR and NMR spectroscopy and elemental analysis, and complexes 1, 4, 5, and 6 were characterized by single-crystal X-ray diffraction (Figure 1).^[9] The ¹H NMR spectra and microanalytical data for each complex showed that the complexes contain a 2:1:1 ratio of phen/Cu/enolate. The solid-state structures of 1, 4, and 6 contain a cationic Cu¹ center ligated by two phen ligands and a free, unligated anionic enolate unit.^[10] The sp²

[N₄Cu]



Figure 1. ORTEP diagrams of 1 (top left, C50–C51 = 1.400(3) Å, C51– C52 = 1.392(6) Å, \pm C50-C51-C52 = 128.3(5)°), 4 (top right, C27– C26 = 1.400(3) Å, C27–C28 = 1.398(3) Å, \pm C26-C27-C28 = 124.71(17)°), 5 (bottom left, C1–N1 = 1.151(4) Å, C1–C2 = 1.396(5) Å, C2– C3 = 1.399(5) Å, \pm C1-C2-C3 = 119.9(4)°), and 6, (bottom right, C28– C29 = 1.432(6) Å, C27–C28 = 1.391(5) Å, \pm C27-C28-C29 = 126.3(4)°). Thermal ellipsoids shown at 50% probability.

carbanions in the enolates (C51 in 1, C27 in 4, C28 in 6) are nearly trigonal planar, and, consistent with a conjugated enolate, the C–C bond distances are close to those in an aromatic ring.

In contrast to the enolates in complexes 1, 4, and 6, the cyanoester anion in 5 is bound to copper through the N atom of the cyano group (Figure 1), and this binding creates an unusual five-coordinate, trigonal bipyramidal, formally 20 valence electron Cu^{I} site. The Cu–N bonds in the equatorial plane (1.947(3) to 2.071(3) Å) are considerably shorter than those in the axial positions (2.368(3) and 2.411 Å).

The conductivity of solutions containing the complexes was used to assess the relationship between the solid-state and solution structures. The conductivities of 1.0 mM solutions of complexes **1–7** in DMSO solvent ranged from 18.8 to $24.5 \ \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ versus $23.5 \ \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ for [NBu₄]-[BPh₄] and $0.3 \ \Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ for ferrocene. Thus, complexes **1–7** are all predominantly salts in DMSO, including complex **5**, which adopts the neutral, five-coordinate structure in the solid state.

To assess the potential of the enolate complexes 1–7 to be intermediates in the α -arylation reactions, we investigated their reactions with iodobenzene (PhI). These results are summarized in Table 1. The malonate, cyanoester, and phenylacetate complexes 3–7 reacted with PhI at 25 °C to form the coupled products in 68–86% yield (100% conversion) after 12 hours, whereas the ethyl acetoacetate complex 2 required over 22 hours at 60 °C for >95% conversion and formed the arylated product in 63% yield. The acetylacetone complex 1 did not react with PhI to form the arylated product; no reaction occurred at room temperature, and multiple unidentified species formed at 60 °C. Table 1: Reactions of copper enolate complexes with PhI.[a]

	⁺ R ¹ _C R ² ⁻ H 1-7	5 equiv Phl	R ¹ R ² + 1/2 [{(phen + pher)Cul}2] 1
Complex	R ¹	R ²	<i>T</i> [°C]	Yield [%] ^[b]
1	MeCO	MeCO	60	0
2	MeCO	EtCO ₂	60	63 ^[c]
3	EtCO ₂	EtCO ₂	25	79
4	MeCO ₂	MeCO ₂	25	80
5	CN	EtCO ₂	25	86
6	Ph	EtCO ₂	25	68
7	Ph	MeCO ₂	25	70

[a] Reaction conditions: 34.4 mM of complexes, [D₆]DMSO. [b] The yield was determined by ¹H NMR spectroscopy using trimethoxybenzene as an internal standard. [c] Reaction time: 22 h.

The reaction of diethyl malonate with 2 equivalents of PhI in the presence of 2 equivalents of Cs_2CO_3 catalyzed by the Cu^1 diethyl malonate complex **3** (10 mol%) gave the diethyl arylmalonate in 73% yield after 8 hours and in 95% yield after 20 hours at 25°C.^[11] The similarity of this rate to that for the stoichiometric reaction of **3** with PhI indicates that complex **3** is kinetically competent to be an intermediate in the catalytic process.

Kinetic data gained by ¹H NMR spectroscopy on the reactions of **3** with varying excess quantities of PhI and added phen at 60 °C showed that the reaction is first order in **3**, first order in PhI, and inverse first order in phen (see the Supporting Information for data). These data imply that the reaction occurs by reversible dissociation of one phen from **3** and that the reactive Cu¹ enolate species contains a 1:1:1 ratio of phen/Cu/enolate. This complex could be the ionic species [(phen)₂Cu][Cu{CH(CO₂Et)₂]₂] (**A**) or the neutral complex [(phen)Cu{CH(CO₂Et)₂]] (**B**) [Eq. (2)].

Scheme 2 summarizes the rate constants for analogous reactions of complexes 1–7 with 10 equivalents of PhI in the presence of 12 equivalents of phen. The relative reactivity follows the trend: 1, $2 < 3 < 4 < 5 < 6 \approx 7$. These data show



Scheme 2. Rate of reactions of enolate complexes 1-7 with PhI. [a] < 5 % conversion after 18 h at 60 °C.

that complexes of the more-electron-donating enolates react with the iodoarenes faster than the complexes of the lesselectron-donating enolates.

The catalytic reactions of diethyl malonate with PhI in the presence of 10 mol% of **3** were also inhibited by added phen. Without added phen, this reaction formed the product in 73% yield after 8 hours and in 95% yield after 20 hours at room temperature. The same reaction conducted with 30 mol% of added phen gave diethyl phenylmalonate in only 4% yield after 8 hours and 11% yield after 20 hours (11% conversion). The same reaction with a full equivalent of added phen gave no detectable arylated product after 8 hours. These results are consistent with the reversible dissociation of one phen ligand from **3** to generate the active intermediate (**A** or **B**) in the catalytic process.

To gain more direct information on the reactivity of the species generated by dissociation of phen, we generated this complex by treatment of the complex 3 (34.4 mM) with 1 equivalent of CuI in DMSO [Eq. (3)]. This reaction formed

$$[(phen)_2Cu][CH(CO_2Et)_2] + Cul \xrightarrow{} -1/2 [((phen)Cul)_2] A \longrightarrow B$$
(3)

[{(phen)CuI}₂] and the ionic or neutral complex **A** or **B** containing a 1:1 ratio of phen to enolate, as indicated by ¹H NMR spectroscopy. The species containing a single phen reacted with 2 equivalents of PhI to form diethyl phenyl-malonate in 37% yield in less than 10 minutes at 25°C and 62% yield with 100% conversion of the copper species after 1 hour,^[12] whereas complex **3** reacted with PhI under analogous reaction conditions to form the coupled product in only 23% yield after 78 minutes and 63% yield after 9 hours.

To assess further the potential of the species containing a 1:1:1 ratio of phen/Cu/enolate to be the active intermediates in the reactions of the isolated complexes with aryl halides, the ratio of products from reactions of sterically distinct o- and p-iodotoluenes with the Cu^I enolate complexes containing or lacking phen ligands was measured (Table 2). The

Table 2: Selectivity of Cu¹ enolate complexes toward *o*- and *p*-iodo-toluene.

		$CH(CO_2Et)_2 CH(CO_2Et)_2$					
c e	opper(I) + + + (•	\bigcirc			
		Þ	 para-substituted product	<i>ortho</i> -substitute product	ed		
Entry	Cu ^I enolate	t	Yield [%] Ratio		Ratio		
			para-	ortho-			
			substituted	substituted			
			product	product			
1	3	1 h	45	2	95:5		
I	3	3 h	73	3	95:5		
2	A or B	< 5 min	62	2	97:3		
2	K[CH(CO ₂ Et) ₂] +	1 h	6.2	2.4	72:28		
2	Cul	21 h	13	6	69:31		
4	2 K[CH(CO ₂ Et) ₂] +	2 h	23	10	70:30		
4	Cul	23 h	37	16	69:31		

reaction of a 1:1 mixture of o- and p-iodotoluene (10 equiv each) with complex **3** (34.4 mM) formed two diethyl arylmalonates in a 95:5 ratio (entry 1). The reaction of o- and piodotoluene with complex **A** or **B**, generated by treatment of complex **3** (34.4 mM) with 1 equivalent of CuI, gave the diethyl arylmalonates in a similar 97:3 ratio (entry 2). The similarity of this ratio is consistent with our conclusion that the species reacting with the iodoarene contains a 1:1:1 ratio of phen, enolate, and copper, as drawn from kinetic data on the reactions of isolated **3** in the presence of added phen.

Although attempts to generate enolate complexes lacking any dative ligand did not lead to pure, isolable species (see the Supporting Information for details), we did conduct stoichiometric reactions of the aryl iodides with the combination of copper and alkali metal enolates. The reaction of o- and piodotoluene with a 1:1 and 2:1 ratio of potassium diethyl malonate to CuI (34.4 mm each) without phen, formed the diethyl arylmalonates in an approximately 70:30 ratio and a total yield of only 19% after 21 hours (Table 2, entry 3) and 53% after 23 hours (entry 4), respectively. The low selectivity and the slow rate of these reactions in the absence of phen further imply that the iodoarenes react with a phen-ligated copper species during the reactions initiated with isolated phen-ligated complexes. Moreover, the lower selectivity of the enolate complex containing the putative [Cu{H- $(CO_2Et)_{2}^{-1}$ anion (entry 4) implies that the active intermediate in the reaction of 3 with iodoarenes is the neutral phen-ligated copper enolate species **B**, rather than the anionic $[Cu{CH(CO_2Et)_2}_2]^-$ portion of the ionic species **A**.

With an understanding of the structure of the starting enolate complexes and the composition of the reactive intermediate, we sought to distinguish between reaction mechanisms involving electron transfer to generate aryl radicals proposed previously^[7] and mechanisms involving the formation of Cu^{III} intermediates by nonradical pathways more akin to those deduced for reactions of complexes with reactive anionic oxygen and nitrogen ligands.^[13] To do so, we studied the reaction of complex **3** with *o*-(allyloxy)iodobenzene [**8**; Eq. (4)]. The aryl radical of **8** is known to undergo



cyclization with a rate constant of $9.6 \times 10^9 \text{ s}^{-1}$ in DMSO to form a [3-(2,3-dihydrobenzofuran)]methyl radical, which is eventually converted into 2-methyldihydrobenzofuran (11).

The reaction of **3** with **8** in DMSO at 25 °C produced the arylated molonate **9** in 59% yield and ethyl 2-aryl-2-oxoacetate **10** in 14% yield after two days [Eq. (4)].^[14] No detectable amount of cyclized product **11** was observed by GC/MS. The detection limit for product **11** by GC/MS was

1030 www.angewandte.org

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

determined to be less than 0.2% of the amount of **9**. Thus, any aryl radical formed from this process must react with the copper enolate complex to form the arylmalonate **9** in less than 2×10^{-13} s. Because this timescale is the lifetime of a transition state, and the organic product would be required to form by recombination of the free aryl radical with the copper enolate in less than 2×10^{-13} s, free radicals that could be formed by initial electron transfer are unlikely to be intermediates in the reaction of copper malonate complexes with aryl halides.

To assess the potential intermediacy of the arylcopper(III) complexes proposed to lie on the reaction pathway, we computed the energies for oxidative addition of PhI to the phen-ligated C-bound Cu^I complexes of acetylacetone, dimethyl malonate, and methyl phenylacetate anions. These three enolate complexes (**1**, **4**, and **7**) reacted with iodoarene with the most diverse rates (see Scheme 2). The free energies of activation (ΔG^{\pm}) for oxidative addition of PhI to the C-bound Cu^I complexes of the anions of acetylacetone, dimethyl malonate, and methyl phenylacetate to form an aryl–Cu^{III} intermediate were calculated to be 27.2, 21.9, and 22.7 kcal mol⁻¹, respectively, at 25 °C (see **1**-TS1, **4**-TS1, and **7**-TS1 in Figure S1 in the Supporting Information). These computational results imply that the Cu^{III} species lie at energies that are accessible under mild reaction conditions.

However, the copper enolate complexes containing a single phen ligand could exist in the O,O-bound (or O-bound) form **B1** or in the C-bound form **B2** (Scheme 3; also see



Scheme 3. Proposed catalytic cycle with complex **3**. Oxidative addition to the C-bound enolate is shown. Oxidative addition to an O-bound isomer, followed by rearrangement to the C-bound Cu^{III} intermediate is less likely, but not ruled out by our data.

Table S5 in the Supporting Information). Thus, we computed the relative free energies of the C-bound species and the Obound forms (Table S5). The O,O-bound structures containing the anions of acetylacetone and dimethyl malonate are computed to lie at energies that are lower than those of the Cbound Cu^I structures by 12.5 and 4.2 kcal mol⁻¹, respectively (see 1-O,O and 4-O,O in Figure S1). Thus, the transition state for oxidative addition to the C-bound isomers of the complexes of acetyl acetonate and dimethyl malonate anions are predicted to lie 39.7 kcal mol⁻¹ and 26.1 kcal mol⁻¹ above the energies of the most stable species containing a 1:1:1 ratio of enolate, phen, and copper. In contrast, the more stable Cu^I complex containing the anion of methyl phenylacetate is computed to be the C-bound form. Thus, the barrier for oxidative addition of PhI to the C-bond enolate containing a 1:1:1 ratio of enolate, phen, and copper is predicted to be only 22.7 kcal mol⁻¹ above the most stable species in this case (Figure S1). The prohibitively large barrier calculated for oxidative addition to the acetylacetonate complex is consistent with our observation of the lack of reactivity of **1** with PhI to form 3-phenyl acetylacetone. The 26.1 and 22.7 kcal mol⁻¹ barriers computed for reaction of the malonate and phenylacetate complexes, respectively, agree with the moderate rate observed for the reaction of PhI with complex **4** and the faster rate observed for the reaction of iodobenzene with complex **7** (see Scheme 2).^[15]

The calculated barriers for reductive elimination of α -aryl carbonyl compounds from the C-bound Cu^{III} acetylacetone, dimethyl malonate, and methyl phenylacetate complexes are 3.5, 4.1, and 6.3 kcal mol⁻¹, respectively (see **1**-TS2, **4**-TS2, and **7**-TS2 in Figure S1 in the Supporting Information). These data indicate that the reductive elimination to form the C–C bond from copper complexes of these stabilized anions is much faster than reductive elimination to form the same type of C–C bond from known arylpalladium(II) enolate complexes.^[16]

In summary, studies on isolated, phen-ligated enolate complexes of 1,3-dicarbonyl compounds and phenylacetates strongly indicate that the C-bound Cu^I enolate complex ligated by a single phen lies on the pathway for the reaction of copper enolates with iodoarenes (Scheme 3). The most stable Cu^I enolate complex containing two phen ligands reversibly dissociates one phen ligand to form the O,O-bound (or Obound) enolate species B1 containing one phen ligand. We propose that this complex equilibrates with the C-bound isomer B2 and that oxidative addition of the iodoarene occurs to the C-bound isomer to form a Cu^{III} enolate intermediate, which undergoes facile reductive elimination to release the arylated product and [{(phen)CuI}2]. Among the enolate complexes studied, the phenylacetate complex, for which the C-bound enolate is computed to be more stable than the alternative O-bound form, and for which the enolate is the most strongly electron donating, is the most reactive toward oxidative addition of iodoarenes. The acetylacetone complex, for which the O,O-chelating mode is calculated to be more favorable than the C-bound form and for which the enolate is the most weakly electron donating, is the least reactive toward addition of iodoarenes. The intermediacy of the phenligated enolate, the equilibration of species containing a free enolate anion and a C-bound enolate, and reaction by pathways lacking aryl radicals all contradict prior assertions about the mechanism of copper-catalyzed couplings of enolates^[3,7] and require revisions of previous mechanistic proposals.

Received: September 22, 2011 Revised: October 18, 2011 Published online: December 8, 2011

Keywords: copper · cross-coupling · kinetics · reaction mechanism · structure elucidation

^[1] W. R. H. Hurtley, J. Chem. Soc. 1929, 1870.

 ^[2] a) F. Monnier, M. Taillefer, Angew. Chem. 2009, 121, 7088;
 Angew. Chem. Int. Ed. 2009, 48, 6954; b) K. Okuro, M. Furuune,
 M. Miura, M. Nomura, J. Org. Chem. 1993, 58, 7606.

Angewandte Communications

- [3] E. J. Hennessy, S. L. Buchwald, Org. Lett. 2002, 4, 269.
- [4] H. J. Cristau, P. P. Cellier, J. F. Spindler, M. Taillefer, *Chem. Eur. J.* 2004, 10, 5607.
- [5] a) X. A. Xie, Y. Chen, D. W. Ma, J. Am. Chem. Soc. 2006, 128, 16050; b) X. Xie, G. Cai, D. Ma, Org. Lett. 2005, 7, 4693; c) Y. W. Jiang, N. Wu, H. H. Wu, M. Y. He, Synlett 2005, 2731.
- [6] S. F. Yip, H. Y. Cheung, Z. Zhou, F. Y. Kwong, Org. Lett. 2007, 9, 3469.
- [7] a) J.-i. Setsune, T. Ueda, K. Shikata, K. Matsukawa, T. Iida, T. Kitao, *Tetrahedron* **1986**, *42*, 2647; b) J.-i. Setsune, K. Matsukawa, T. Kitao, *Tetrahedron Lett.* **1982**, *23*, 663; c) J. Setsune, K. Matsukawa, H. Wakemoto, T. Kitao, *Chem. Lett.* **1981**, 367.
- [8] Complex 1 has been synthesized by a different method, but its structure remained unknown. R. Nast, W.-H. Lepel, *Chem. Ber.* 1969, 102, 3224.
- [9] CCDC 844995 (1), 844996 (4), 844997 (5), and 844998 (6) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [10] Related bis(phen)-ligated Cu^I nitrophenylmethane and nitromethane complexes in which the enolate moieties were proposed to be present as the free anions were reported previously, but their structures were not defined by X-ray diffraction. N. Marsich, A. Camus, J. Inorg. Nucl. Chem. **1977**, 39, 275.
- [11] The catalytic reaction with PhBr as the electrophile gave no product under the same reaction conditions, but formed the diethyl arylmalonate in 86% yield after 15 h at 80°C.
- [12] Full conversion of the species B into diethyl phenylmalonate and free diethyl malonate after 1 h. The latter was presumably formed by protonation of B with the trace amount of water in the DMSO.
- [13] a) L. M. Huffman, A. Casitas, M. Font, M. Canta, M. Costas, X. Ribas, S. S. Stahl, *Chem. Eur. J.* **2011**, *17*, 10643; b) L. M.

Huffman, S. S. Stahl, *Dalton Trans.* 2011, 40, 8959; c) A. Casitas,
N. Ioannidis, G. Mitrikas, M. Costas, X. Ribas, *Dalton Trans.* 2011, 40, 8796; d) H.-Z. Yu, Y.-Y. Jiang, Y. Fu, L. Liu, J. Am. Chem. Soc. 2010, 132, 18078; e) J. W. Tye, Z. Weng, R. Giri, J. F. Hartwig, Angew. Chem. 2010, 122, 2231; Angew. Chem. Int. Ed. 2010, 49, 2185; f) R. Giri, J. F. Hartwig, J. Am. Chem. Soc. 2010, 132, 15860; g) A. E. King, L. M. Huffman, A. Casitas, M. Costas, X. Ribas, S. S. Stahl, J. Am. Chem. Soc. 2010, 132, 12068; h) L. M. Huffman, S. S. Stahl, J. Am. Chem. Soc. 2008, 130, 9196; i) J. W. Tye, Z. Weng, A. M. Johns, C. D. Incarvito, J. F. Hartwig, J. Am. Chem. Soc. 2008, 130, 9971; j) S.-L. Zhang, L. Liu, Y. Fu, Q.-X. Guo, Organometallics 2007, 26, 4546.

- [14] Compound 10 was likely formed through the oxidation of 9. P. M. Allen, U. Hess, C. S. Foote, M. M. Baizer, *Synth. Commun.* 1982, *12*, 123.
- [15] These isolable species are the ionic complexes 1, 4 and 7, rather than the neutral complexes on which we are able to conduct meaningful calculations. The relative stability of these ionic species and the neutral species [(phen)Cu(enolate)] is difficult to evaluate by computational methods because the ions can exist as contact, solvent-separated, or fully solvated ion pairs. At the same time, the acidity of the carbonyl compounds in DMSO (acetylacetone = 13.3, dimethyl malonate = 15.9 and methylphenylacetate \approx 22) shows clearly that the enolate from acetylacetone is more stable than that from dimethyl malonate, which is more stable than that from methyl phenylacetate. Thus the energy difference between the observed ionic enolate complex and the neutral species should follow the trend, acetylacetone > dimethyl malonate > methyl phenylacetate. These energy differences would further increase the difference of free-energy barriers between reactions of PhI with the complexes 1, 4 and 7. Thus, the relative computed barriers would remain consistent with the observed reactivity of these enolate complexes with PhI.
- [16] D. A. Culkin, J. F. Hartwig, Acc. Chem. Res. 2003, 36, 234.