

A NEW METHOD FOR THE PREPARATION OF N-(NUCLEOSIDYL)- α -AMINO ACIDS
USING TRIOCTYLMETHYLAMMONIUM SALTS OF α -AMINO ACIDS

Frank Roland Schröder* and Friedrich Cramer

Max-Planck-Institut Für Experimentelle Medizin, Abteilung Chemie, D-3400 Göttingen, FRG


8-Bromoadenosine reacts with trioctylmethylammonium salts of α -amino acids (e.g. glycine) to yield N-(adenosin-8-yl)- α -amino acids which can be readily separated from the starting material by two subsequent extraction steps at different pH values.

N-(nucleosidyl)- α -amino acids play an important part in biological chemistry. N-(Inosin-2-yl)-L-alanine has been found in *Fusarium* species¹, while [N-(nebularin-6-yl)carbamoyl]-L-threonine is a constituent of the modified nucleosides in tRNA². N-(Adenosin-8-yl)-L-valine acts as an α -amino acid analogue which can inhibit the binding of valine to valyl-tRNA synthetase from yeast³.

N-(Nucleosidyl)- α -amino acids are prepared by nucleophilic attack of the α -amino group of an α -amino acid on the corresponding halogeno or methylthio substituted nucleoside⁴. The reaction product is usually isolated by anion-exchange chromatography because the recrystallization of these compounds is extremely difficult⁵. In the purine ring system a substituent with an unshared pair of electrons (electron donor) at C-6 (e.g. -NH₂, -NHR, -NR₂ with R = Alkyl) increases the electron density at C-8⁶ so the H-8 can very easily be substituted by a convenient electrophile e.g. bromine. On the other hand, the velocity of a S_{NAr} reaction at this carbon atom is decreased. Therefore it has been suggested that in such ring systems a bromine atom at C-8 cannot be substituted by an α -amino acid or a similar weak nucleophile⁷. Initially we attempted such a synthesis by reaction of 8-bromoadenosine (2) with a twofold excess of sodium glycinate at 120°C in dimethylsulfoxide, for 28 h (A). The main difficulty in this step is the low solubility of sodium glycinate in dimethylsulfoxide. N-(adenosin-8-yl)glycine (3)⁸ was formed in a 20% yield as estimated by thin layer chromatography⁹. Additionally unreacted 8-bromoadenosine (2) and 8-bromoadenine (6) were found, when the reaction mixture was chromatographed on a diethylaminoethyl-cellulose (DEAE A 25) column with a triethylammonium bicarbonate (TEA) gradient. N-(Adenosin-8-yl)-L-alanine (4) and N-(adenosin-8-yl)-L-valine (5) could be prepared and purified in an analogous way¹⁰.

Trioctylmethylammonium chloride (Adogen¹¹) is a potent phase transfer reagent¹². Because of this we prepared the trioctylmethylammonium salt of glycine by extraction of a mixture of Adogen and glycine in water/n-propanol with chloroform at pH 11. The product was a yellow oil which dissolved readily in toluene and other non polar solvents. The substance gave a positive Ninhydrin test and could be used without further purification. A five fold excess of trioctylmethylammonium glycinate over 8-bromoadenosine (2) gave after 12 h in dimethylsulfoxide at 105°C N-(adenosin-8-yl)glycine (3) in 60 % yield (as estimated by thin layer

chromatography)⁹ (B); no byproducts could be detected. Water was added and the trioctylmethylammonium salt of (3) could be extracted with chloroform from the aqueous phase at pH 6 when the heterocyclic amino groups are not protonated. 8-Bromoadenosine (2), the unreacted trioctylmethylammonium glycinate, which is protonated at this pH, and dimethylsulfoxide remained in the aqueous phase. The organic phase was shaken vigorously with 25% aqueous ammonia to convert the trioctylmethylammonium salt of (3) into the ammonium salt which is soluble in water. After evaporation of excess ammonia and water N-(adenosin-8-yl)glycine (3)⁸ could be purified by recrystallization.



With this method even the rather unreactive 8-bromoadenosine (2) does react with the amino groups of amino acids in satisfactory yield. Thus, this method may be extended to other convenient substituted nucleosides, providing a general synthesis for N-(nucleosidyl)-amino acids.

References and Notes

1. A. Ballio, C. Delfini and S. Russi, *Nature* **186**, 968-969 (1960).
2. J.A. McCloskey and S. Nishimura, *Accounts Chem. Res.* **10**, 403-410 (1977).
3. F.R. Schröder, unpublished results.
4. M. Kawana, P.C. Srivastava and R.K. Robins, *J. Carboh. Nucleosides, Nucleotides* **8**, 131-144 (1981) and references herein.
5. J.F. Gerster and R.K. Robins, *J. Amer. Chem. Soc.* **87**, 3752-3759 (1965).
6. H. Lönnberg, P. Lehtikoinen and K. Neuvonen, *Acta Chem. Scand.* **B 36**, 707-712 (1982).
7. T. Kato, S. Ogawa and I. Ito, *Tetrahedron Lett.* **22**, 3205-3208 (1981).
8. N-(adenosin-8-yl)glycine (3): UV spectrum $\lambda_{\max} = 278.5$ nm, $\epsilon = 18214$ (pH 1); elemental analysis - found: C 42.22 %, H 4.61 %, N 25.93 %, calculated for $C_{12}H_{16}N_6O_6$: C 42.20 %, H 4.7 %, N 24.6 %.
¹H NMR (100 MHz DMSO-*d*₆) δ 7.94, 1 H, s, 2-H; 6.52, 2 H, s, 6-NH₂; 5.9, 1 H, d, 1'-H; 4.76, 1 H, t, 2'-H; 4.14, 1 H, m, 3'-H; 4.04, 1 H, m, 4'-H; 3.66, 2 H, d, 5'-H; 3.18, 2 H, s (-CH₂- of glycine).
9. The compounds synthesized according to (A) and (B) were identical by thin layer chromatography, UV spectra and elemental analysis.
10. Thin layer chromatography was performed using DC-Mikroarten SI-F purchased from RIEDEL DE HAEN, Seelze (FRG) with chloroform : methanol : 25 % aqueous ammonia (7 : 3 : 1) (v/v/v) as an eluent.
11. F.R. Schröder, Diplomarbeit, Göttingen (FRG) (1981).
12. Adogen was purchased from SERVA, Heidelberg (FRG), another trade name is Aliquat 336.
13. J. Barry, G. Bram, G. Decodts, A. Loupy, P. Pigeon and J. Sansoulet, *Tetrahedron Lett.* **23**, 5407-5408 (1982).

(Received in Germany 18 March 1983)