



Pergamon

Tetrahedron Letters 40 (1999) 8439–8441

TETRAHEDRON
LETTERS

Facile cleavage of carbohydrate benzyl ethers and benzyldene acetals using the $\text{NaBrO}_3/\text{Na}_2\text{S}_2\text{O}_4$ reagent under two-phase conditions

Matteo Adinolfi, Gaspare Barone, Luigi Guariniello and Alfonso Iadonisi *

Dipartimento di Chimica Organica e Biologica, Università degli Studi di Napoli Federico II, Via Mezzocannone 16, I-80134 Napoli, Italy

Received 28 July 1999; accepted 14 September 1999

Abstract

Benzyl ether and benzyldene acetal carbohydrate protecting groups can be selectively cleaved by reaction with sodium bromate/sodium dithionite in ethyl acetate/water. Under the mild (room temperature, short reaction time) conditions needed, a variety of other protecting functionalities such as acetyl, chloroacetyl, benzoyl, pivaloyl, tosyl, *t*-butyldimethylsilyl, trityl, and isopropylidene groups remain unaffected. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords: cleavage reactions; carbohydrates.

Benzyl ethers are widely used in organic synthesis for the protection of hydroxyl groups because of their tolerance to a wide range of chemical conditions. These versatile *O*-protecting groups, among the most useful in carbohydrate chemistry, are typically removed with Pd/C (or $\text{Pd}(\text{OH})_2/\text{C}$) with hydrogen,^{1–3} by catalytic transfer hydrogenation using formic acid⁴ or other reagents^{3,5,6} as suitable sources of hydrogen, or in the presence of strong Lewis acids, such as $\text{BF}_3 \cdot \text{OEt}_2$ (in combination with sulfur nucleophiles),⁷ strictly anhydrous FeCl_3 ,^{8,9} TMSI¹⁰ etc. Furthermore, the deprotection can be performed under strongly basic conditions with sodium or lithium in ammonia.^{11,12} Debenzylation can also be performed under UV irradiation by using NBS/ H_2O through a reaction pathway presumably involving radical bromination at the benzylic position and subsequent hydrolysis of the resulting α -bromo ether.¹³

An interesting approach for the selective mono α -bromination of alkyl benzenes based on the combined use of NaBrO_3 and a suitable reducing reagent (such as NaHSO_3) under two-phase conditions has recently appeared in the literature.¹⁴ The proposed mechanism contemplates the gradual decomposition of the formed HBrO in aqueous solution with liberation of $\text{Br} \cdot$ which moves to the organic phase and brominates the side-chain of the dissolved arene. Due to our interest in carbohydrate chemistry, we envisaged that such a reaction could be usefully exploited for the removal of benzyl protecting groups

* Corresponding author. Fax: +39 0815521217; e-mail: iadonisi@unina.it

Table 1
Cleavage of benzyl and benzylidene carbohydrate protecting groups

Entry	Reagent	Products ^a	Time	Yield
1			3h	82%
2			6h	85%
3			3h	96%
4			3h	78%
5			1.5h	97% ^b
6			2h	73%
7			1.5h	88% 7b/7c=35/65
8			1h	92% ^c

^a Satisfactory spectral data were obtained for all products. ^b With NaHSO₃: time 2h, yield 95%. ^c With NaHSO₃: time 1h, yield 80%.

from sugar substrates. As a matter of fact we have found that the use of NaBrO₃ and sodium dithionite (Na₂S₂O₄) in ethyl acetate/water at room temperature is particularly effective for the fast and clean deprotection of several benzylated monosaccharides (Table 1, entries 1–6).¹⁵

Various functionalities such as esters, silyl, trityl ethers, acetanides and tosylates commonly employed for the derivatization of sugars, survived unaffected under the reaction conditions.

The propensity of cyclic benzylidenes to give benzylic bromination and subsequent oxidative opening with NBS/H₂O under UV or thermal activation is well known.¹⁶ Therefore, we tested the NaBrO₃/Na₂S₂O₄ procedure to perform that transformation. The six-membered cyclic 4,6-*O*-benzylidene derivative **7a** was converted to a mixture of the 6-*O*- and 4-*O*-benzoyl regioisomers **7b** and **7c**, in high yield but with low regioselectivity (35:65).¹⁷ On the other hand, the five-membered benzylidene **8a** was converted into the single axial 4-*O*-benzoyl isomer **8b**. Interestingly, the use of NaBrO₃/Na₂S₂O₄ required shorter reaction times and resulted in higher yields than the NBS/H₂O/UV procedure.¹⁶

In a typical procedure the benzylated (or benzylidene) sugar (0.3 mmol) was dissolved in ethyl acetate (4 mL) and then a solution of NaBrO₃ (136 mg, 0.9 mmol) in water (3 mL) was added. To the well stirred two-phase system an aqueous solution of Na₂S₂O₄ (85% pure, 157 mg, dissolved in 6 mL water) was added dropwise over 10 min at room temperature. After completion of the reaction (TLC) the mixture was diluted with EtOAc and the organic phase was washed with aq. sodium thiosulfate. The crude product was then purified by silica gel chromatography. With NaHSO₃ in place of Na₂S₂O₄ slightly lower yields were observed (entries 5 and 8).

In conclusion we have shown that the use of the NaBrO₃/Na₂S₂O₄ system can be a valuable alternative approach for the deprotection of benzyl ethers due to the mildness, cheapness, and experimental simplicity of the procedure.

Acknowledgements

Financial support by MURST and CNR is acknowledged.

References

- Freifelder, M. *Catalytic Hydrogenation in Organic Synthesis*; Wiley: New York, 1978; p. 109.
- Pearlman, W. M. *Tetrahedron Lett.* **1967**, 17, 1663–1664.
- Hanessian, S.; Liak, T. J.; Vanasse, B. *Synthesis* **1981**, 396–397.
- Rao, V. S.; Perlman, A. S. *Carbohydr. Res.* **1980**, 83, 175–177.
- Felix, A. M.; Heimer, E. P.; Lambros, T. J.; Tzougraki, C.; Meienhofer, J. *J. Org. Chem.* **1978**, 43, 4194–4196.
- Bieg, T.; Szeja, W. *Synthesis* **1985**, 76–77.
- Fuji, K.; Ichikawa, K.; Node, M.; Fujita, E. *J. Org. Chem.* **1979**, 44, 1661–1664.
- Kartha, K. P. R.; Dasgupta, F.; Singh, P. P.; Srivastava, H. C. *J. Carbohydr. Chem.* **1986**, 5, 437–444.
- Rodebaugh, R.; Debenham, J. S.; Fraser-Reid, B. *Tetrahedron Lett.* **1996**, 37, 5477–5478.
- Klemer, A.; Bieber, M.; Wilbers, H. *Justus Liebigs Ann. Chem.* **1983**, 1416–1421.
- Holick, S. A.; Anderson, L. *Carbohydr. Res.* **1974**, 34, 208–213.
- McCloskey, C. M. *Adv. Carbohydr. Chem.* **1957**, 12, 149.
- Bingley, R. W.; Hehemann, D. G. *J. Org. Chem.* **1990**, 55, 378–380.
- Kikuchi, D.; Sakaguchi, S.; Ishii, Y. *J. Org. Chem.* **1998**, 63, 6023–6026.
- Benzaldehyde (traces) and benzoic acid were detected by NMR in the crude reaction material as by-products.
- Bingley, R. W.; Goewey, G. S.; Johnston, J. C. *J. Org. Chem.* **1984**, 49, 992–996.
- The mechanistically related DDQ oxidative opening of 4,6-*O*-*p*-methoxybenzylidene sugar derivatives also proceeds with low regioselectivity (Zhang, Z.; Magnusson, G. *J. Org. Chem.* **1996**, 61, 2394–2400).