

prepared triphenyl phosphine methylene<sup>7</sup> in a pressure bottle using dry ether as solvent and nitrogen at 65° for three hours. From the mixture was obtained, after two chromatographic separations, a highly viscous liquid; yield, 14%. *Anal.* Calcd. for C<sub>16</sub>H<sub>24</sub>O (III): C, 82.69; H, 10.41. Found: C, 82.61; H, 10.34;  $\epsilon$  (265 m $\mu$ ), 23,200. The infrared absorption spectrum showed bands due to the presence of the =CH<sub>2</sub> group at 890, 1594, 1625 and 1646 cm.<sup>-1</sup>. The corresponding bands for vitamin D<sub>3</sub> examined at the same time were at 892, 1600, 1630 and 1650 cm.<sup>-1</sup>.

We wish to acknowledge financial support from the Research Corporation and to thank Dr. S. M. Nagy and his associates of this Institute for the analyses and the infrared spectra.

(7) G. Wittig and U. Schöllkopf, *Chem. Ber.*, **87**, 1318 (1954).

DEPARTMENT OF CHEMISTRY  
MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
CAMBRIDGE 39, MASSACHUSETTS

NICHOLAS A. MILAS  
Li-CHIN CHIANG  
CHARLES P. PRIESING  
ASHER A. HYATT  
JOHN PETERS

RECEIVED JUNE 16, 1955

SYNTHESIS AND BIOLOGICAL ACTIVITY OF 1- AND 6-DEHYDRO-9 $\alpha$ -HALOCORTICOIDS

Sir:

It has been demonstrated in our laboratories that substitution of a chlorine<sup>1,2</sup> or more effectively of a fluorine<sup>2,3</sup> atom in the 9 $\alpha$ -position of 11-oxygenated corticoids leads to considerable enhancement in glucocorticoid<sup>1,2,3,4</sup> anti-inflammatory<sup>5</sup> and sodium retaining activity.<sup>6</sup> More recently, others have shown that introduction of a double bond into the 1,2-position of cortisone and hydrocortisone leads to 3-4 fold increases in both glucocorticoid<sup>7</sup> and antirheumatic<sup>8</sup> activity. It was therefore of

- (1) J. Fried and E. F. Sabo, *THIS JOURNAL*, **75**, 2273 (1953).
- (2) J. Fried, J. E. Herz, E. F. Sabo, A. Borman, F. M. Singer and P. Numerof, *ibid.*, **77**, 1068 (1955).
- (3) J. Fried and E. F. Sabo, *ibid.*, **76**, 1455 (1954).
- (4) A. Borman and F. M. Singer, *Fed. Proc.*, **13**, 185 (1954).
- (5) F. M. Singer and A. Borman, *ibid.*, **14**, 281 (1955).
- (6) A. Borman, F. M. Singer and P. Numerof, *Proc. Soc. Exp. Biol. Med.*, **86**, 570 (1954).
- (7) H. L. Herzog, A. Nobile, S. Tolksdorf, W. Charney, E. B. Hershsberg, P. L. Perlman and M. M. Pechet, *Science*, **121**, 176 (1955).
- (8) J. J. Bunim, M. M. Pechet and A. J. Bollet, *J. Am. Med. Assoc.*, **167**, 311 (1955).

considerable interest to prepare steroids possessing both a 1,2-double bond and a 9 $\alpha$ -halogen atom for biological evaluation. A recent publication<sup>9</sup> describing one such steroid, 1-dehydro-9 $\alpha$ -fluorohydrocortisone acetate prompts us to report on our experiences with this and related compounds.

1-Dehydrohydrocortisone acetate<sup>7</sup> was dehydrated with methanesulfonyl chloride and pyridine in dimethylformamide to  $\Delta^{1,4,9(11)}$ -pregnatriene-17 $\alpha$ ,21-diol-3,20-dione 21-acetate (m.p. 222–223°,  $[\alpha]^{23D} + 52^\circ$  (CHCl<sub>3</sub>),  $\lambda_{\text{max}}^{\text{alc}}$  238 m $\mu$  ( $\epsilon = 16,100$ )<sup>10</sup>; found: C, 71.81; H, 7.10), which on treatment with N-bromoacetamide and perchloric acid in dioxane furnished 1-dehydro-9 $\alpha$ -bromohydrocortisone acetate I (m.p. 180–185° (dec.),  $[\alpha]^{23D} + 123^\circ$  (dioxane),  $\lambda_{\text{max}}^{\text{alc}}$  241 m $\mu$  (13,400); found: C, 57.04; H, 6.29; Br, 16.66; L.G.<sup>11</sup> 4, Na<sup>11</sup> 10). Treatment of I with potassium acetate in boiling alcohol furnished 9 $\beta$ ,11 $\beta$ -oxido- $\Delta^{1,4}$ -pregnadiene-17 $\alpha$ ,21-diol-3,20-dione 21-acetate (m.p. 213–215°,  $[\alpha]^{23D} + 64^\circ$  (CHCl<sub>3</sub>),  $\lambda_{\text{max}}^{\text{alc}}$  249 m $\mu$  (15,800); found: C, 69.06; H, 7.07), which with HBr in chloroform (0°) reverted to I, and with HCl and HF afforded the most potent glucocorticoids presently known, 1-dehydro-9 $\alpha$ -chlorohydrocortisone acetate II (m.p. 242–243° (dec.),  $[\alpha]^{23D} + 145^\circ$  (alc.),  $\lambda_{\text{max}}^{\text{alc}}$  238 m $\mu$  (15,000); found: C, 63.13; H, 6.62; Cl, 8.38; L.G. 13, Na 20–30) and 1-dehydro-9 $\alpha$ -fluorohydrocortisone acetate III (m.p. 243–245°,  $[\alpha]^{23D} + 99^\circ$  (acetone),  $\lambda_{\text{max}}^{\text{alc}}$  238 m $\mu$  (14,500); found: C, 65.88; H, 7.20; F, 4.64; L.G. 28, Na 2–3), respectively. Oxidation of II and III with CrO<sub>3</sub> in acetic acid produced 1-dehydro-9 $\alpha$ -chlorocortisone acetate (m.p. 262–264° (dec.),  $[\alpha]^{23D} + 244^\circ$  (CHCl<sub>3</sub>),  $\lambda_{\text{max}}^{\text{alc}}$  236 m $\mu$  (15,500); found: C, 63.60; H, 6.38; L.G. 12–15) and 1-dehydro-9 $\alpha$ -fluorocortisone acetate (m.p. 274–277°,  $[\alpha]^{23D} + 158^\circ$  (alc.),  $\lambda_{\text{max}}^{\text{alc}}$  235 m $\mu$  (15,600); found: C, 65.91; H, 6.40; L.G. 25–30).

Alternatively, we have independently prepared III from 9 $\alpha$ -fluorohydrocortisone acetate<sup>8</sup> as described by Hirschmann, *et al.*<sup>9</sup> Catalytic reduction (Pd-BaSO<sub>4</sub> in ethyl acetate or alcohol) furnished 9 $\alpha$ -fluoroallopregnane-11 $\beta$ ,17 $\alpha$ ,21-triol-3,20-dione 21-acetate<sup>12</sup> (m.p. 234–235°; [ $\alpha$ ]<sub>D</sub><sup>20</sup> +67° (acetone); found: C, 65.34; H, 7.69; L.G. < 0.1, Na < 0.1), which with two moles of bromine in acetic acid afforded an amorphous mixture of dibromides. The latter upon dehydrobromination with boiling collidine and chromatography on acid-washed alumina yielded successively 2-bromo-9 $\alpha$ -fluoro- $\Delta^1$ -pregnene-11 $\beta$ ,17 $\alpha$ -21-triol-3,20-dione 21-

- (9) R. F. Hirschman, R. Miller, R. E. Beyler, L. H. Sarett and M. Tishler, *Tetrahedron Journal*, **77**, 3166 (1955).
- (10) The infrared spectra of this and other  $\Delta^{1,4}$ -3-keto steroids possess bands in the ranges characteristic of the acetylated dihydroxy-acetone side chain (5.72–5.76  $\mu$  and 5.80–5.85  $\mu$ ) and of the 1,4-diene-3-one system (6.01–6.04  $\mu$ , 6.14–6.20  $\mu$  and 6.20–6.26  $\mu$ ).
- (11) The rat liver glycogen assay (L. G.) represents an excellent measure of glucocorticoid, and in our experience also of antiinflammatory, activity. The figures given are expressed in terms of cortisone acetate = 1. The sodium retention test (Na) employed here has been described in ref. 6. The standard of comparison is DCA = 1.
- (12) The reduction product possesses the *allo*-configuration since on monobromination it yielded mainly the 2-bromo derivative, which on dehydrobromination with collidine afforded 9 $\alpha$ -fluoro- $\Delta^1$ -*allo*pregnene-11 $\beta$ ,17 $\alpha$ ,21-tri-*allo*-2,3-dione 21-acetate<sup>8</sup> (m.p. 237–239°, [ $\alpha$ ]<sub>D</sub><sup>20</sup> +90° (CHCl<sub>3</sub>),  $n_D^{20}$  1.4828  $\mu$  (6, 100); found: C, 65.56%; H, 7.19).

acetate (monoalcoholate: m.p. 184–185° (dec.) after melting and resolidification at 126–139°,  $[\alpha]^{23D} +79^\circ$  (CHCl<sub>3</sub>),  $\lambda_{\max}^{\text{alc}}$  250 m $\mu$  (8,000),  $\lambda_{\max}^{\text{Nujol}}$  2.85  $\mu$ , 2.98  $\mu$ , 5.80  $\mu$ , 5.90  $\mu$ , 5.98  $\mu$ , 6.24  $\mu$ ; found: C, 54.48; H, 6.78; Br, 14.50; OC<sub>2</sub>H<sub>5</sub>, 7.57; L.G. < 0.3), 2-bromo-9 $\alpha$ -fluoro- $\Delta^4$ -pregnene-11 $\beta$ ,17 $\alpha$ ,21-triol-3,20-dione 21-acetate<sup>13</sup> (m.p. 174–175° (dec.);  $[\alpha]^{23D} +136^\circ$  (CHCl<sub>3</sub>);  $\lambda_{\max}^{\text{alc}}$  242 m $\mu$  (12,200),  $\lambda_{\max}^{\text{CHCl}_3}$  2.85–2.95  $\mu$ , 5.78  $\mu$ , 5.85  $\mu$ , 5.94  $\mu$ , 6.15  $\mu$ ; found: C, 55.70; H, 6.30; Br, 15.16; L.G. 1.0, Na 10), 6-dehydro-9 $\alpha$ -fluorohydrocortisone acetate<sup>9,14</sup> (IV) (m.p. 216–217,  $[\alpha]^{23D} +123^\circ$  (alc.),  $+135^\circ$  (CHCl<sub>3</sub>),  $\lambda_{\max}^{\text{alc}}$  281 m $\mu$  (23,000),  $\lambda_{\max}^{\text{Nujol}}$  3.00  $\mu$  (OH), 5.76  $\mu$ , 5.81  $\mu$  (acetylated side chain); 6.10  $\mu$ , 6.16  $\mu$ , 6.22  $\mu$  ( $\Delta^{4,6,3}$ -ketone); found: C, 65.75; H, 7.04; L.G. 5, Na 20–30), the desired III, and an isomer of III (m.p. 271–272°,  $[\alpha]^{23D} +73^\circ$  (alc.),  $\lambda_{\max}^{\text{alc}}$  237 m $\mu$  (15,200),  $\lambda_{\max}^{\text{Nujol}}$  3.00  $\mu$ , 5.75  $\mu$ , 5.92  $\mu$ , 6.04  $\mu$ , 6.18  $\mu$ , 6.24  $\mu$ ; found: C, 65.96; H, 6.84; L.G. < 1).

Substitution of a hydrogen atom for a hydroxyl group at C-21 results in a greater decrease of salt retaining than of glucocorticoid activity.<sup>15</sup> We have therefore prepared the 21-desoxy derivatives of III and IV as follows. Saponification of III and IV with potassium carbonate in aqueous methanol yielded the respective dehydro-9 $\alpha$ -fluorohydrocortisones ( $\Delta^1$ : m.p. 274–275° (dec.),  $[\alpha]^{23D} +94^\circ$  (alc.),  $\lambda_{\max}^{\text{alc}}$  238 m $\mu$  (15,500); found: C, 66.68; H, 7.16) and ( $\Delta^6$ : m.p. 257–259°,  $[\alpha]^{23D} +101^\circ$  (alc.),  $\lambda_{\max}^{\text{alc}}$  281 m $\mu$  (25,600); found: C, 66.30; H, 7.00), which were converted into the 21-mesylates in pyridine at 0° ( $\Delta^1$ : m.p. 220° (dec.),  $[\alpha]^{23D} +98^\circ$  (alc.),  $\lambda_{\max}^{\text{alc}}$  238 m $\mu$  (15,000); found: C, 58.04; H, 6.36; S, 7.52) and ( $\Delta^6$ : m.p. 237–238° (dec.),  $[\alpha]^{23D} +94^\circ$  (alc.),  $\lambda_{\max}^{\text{alc}}$  281 m $\mu$  (27,500); found: C, 58.19; H, 6.05; S, 7.54). The latter were converted into 9 $\alpha$ -fluoro- $\Delta^{1,4}$ -pregnadiene-11 $\beta$ ,17 $\alpha$ -diol-3,20-dione (m.p. 313–314° (dec.),  $[\alpha]^{23D} +47^\circ$  (pyridine),  $\lambda_{\max}^{\text{alc}}$  238 m $\mu$  (15,500); found: C, 69.47; H, 7.66; L.G. 4, Na < 0.1) and 9 $\alpha$ -fluoro- $\Delta^{4,6}$ -pregnadiene-11 $\beta$ ,17 $\alpha$ -diol-3,20-dione (m.p. 294–296°,  $[\alpha]^{23D} +112^\circ$  (dioxane),  $\lambda_{\max}^{\text{alc}}$  281 m $\mu$  (26,000); found: C, 69.66; H, 7.48; L.G. 0.3, Na < 0.1) either directly with sodium iodide in boiling acetic acid or via the 21-iodo derivatives (sodium iodide in acetone) and reduction of the latter with sodium bisulfite in aqueous dioxane.

Our present data may be summarized by stating that introduction of a double bond in the 1,2-position of a 9 $\alpha$ -halocorticoid leads to increases in both glucocorticoid and sodium retaining activity in the rat ranging from about 2.5-fold in the case of the

fluoro to 10-fold in the case of the bromo derivatives. Dehydrogenation in the 6,7-position, on the other hand, effects in the two cases examined a two-fold decrease in glucocorticoid and a 20-fold increase in salt-retaining activity.

JOSEF FRIED  
KLAUS FLOREY  
EMILY F. SABO  
JOSEF E. HERZ  
ALBERT R. RESTIVO  
ALECK BORMAN  
FRANK M. SINGER

THE SQUIBB INSTITUTE FOR  
MEDICAL RESEARCH  
NEW BRUNSWICK, N. J.

RECEIVED JUNE 27, 1955


# THE REACTION OF THIONYL CHLORIDE WITH ALLYLIC ALCOHOLS<sup>1a</sup>

Sir:

Several mechanisms are available for the reaction of allylic alcohols with thionyl chloride.<sup>1b</sup> Without a solvent, mixtures of isomeric chlorides are always obtained. However, we have found that the S<sub>N</sub>i' mechanism<sup>2</sup> may be made very dominant by the use of dilute ether solution, where the liberated hydrogen chloride is rendered quite inactive.<sup>3</sup> Under these conditions, crotyl alcohol yields 99%  $\alpha$ -methylallyl chloride and  $\alpha$ -methylallyl alcohol yields 100% crotyl chloride.

The ether technique has now been found successful even in some more reactive systems.

With optically active *trans*- $\alpha,\gamma$ -dimethylallyl alcohol (I) the more likely conformation of the transition state of the S<sub>N</sub>i' process would lead to active *trans*-chloride (II) and the configuration of the new asymmetric center would be opposite to that of the original one. The less likely conformation would give optically active *cis*-chloride. We have found that *trans*-alcohol (I) is converted to *trans*-chloride (II) (100% *trans*-isomer) which has the opposite configuration of the alcohols as illustrated below. In fact, the optical purity of the chloride is higher than that which results under conditions favorable for direct displacement of chlorosulfinate ion by providing a soluble hydrochloride.<sup>3</sup>



Another  $\alpha,\gamma$ -dialkylsubstituted allyl system for which the ether technique is successful involves the isomeric 5-methyl-2-cyclohexenols.<sup>4</sup> We find it is also successful even with cinnamyl alcohol in ether solution 0.1M in each reagent. Under these conditions, the reaction, slow enough to be followed kinetically, is approximately first order in both alcohol and thionyl chloride. Ultraviolet spectra

(13) This compound was formed in good yield when the reaction temperature was lowered to 100°. It was reduced to 9 $\alpha$ -fluorohydrocortisone acetate with zinc and acetic acid.

(14) It is noteworthy that in contrast to the experience with 9-unsubstituted steroids (cf. A. L. Wilds and C. Djerassi, *THIS JOURNAL*, **68**, 2125 (1946)) the yield of IV exceeded that of III. IV is more satisfactorily prepared, however, by treatment of the dibromide with lithium chloride in dimethylformamide (cf. Holysz, *ibid.*, **75**, 4432 (1953)).

(15) J. Fried, in Conference on Hydrocortisone, its Newer Analogues and Aldosterone as Therapeutic Agents, N. Y. Academy of Sciences, **61**, 573 (1955).

(1a) Acknowledgment is made of the partial support of this research by a National Science Foundation grant.

(1b) W. G. Young, Abstracts of Twelfth National Organic Chemistry Symposium, pp. 23–26 (1951).

(2) J. D. Roberts, W. G. Young and S. Winstein, *THIS JOURNAL*, **64**, 2157 (1942).

(3) W. G. Young, F. Caserio and D. Brandon, *Science*, **117**, 473 (1953).

(4) H. L. Goering, R. D. Nevitt and E. F. Silversmith, *THIS JOURNAL*, **77**, 4042 (1955).