J.C.S. Снем. Сомм., 1972

## Synthesis of Oximino-macrolides

By J. R. MAHAJAN,\* G. A. L. FERREIRA, and H. C. ARAŬJO (Universidade de Brasīlia, Dept. de Quīmica, Brasīlia, Brasil)

Summary A number of oximino-macrolides have been prepared by the action of n-butyl nitrite on the corresponding enol ethers in the presence of aqueous acetic or hydrochloric acid.

WE report a simple synthesis of oximino-macrolides which can be hydrolysed to the corresponding keto-macrolides. Several 10- to 12-membered keto-macrolides have previously been prepared by Borowitz et al.1

A solution of the cyclic enol ether (1 equiv.)<sup>†</sup> and n-butyl nitrite (1 equiv.) t in 50% ethanol was treated with acetic acid or 10% hydrochloric acid. The oximino-macrolide was extracted with ether in almost quantitative yield.§ With the exception of 5-oximino-octanolide, these oximes are smoothly converted into the corresponding semicarbazones and 2,4-dinitrophenylhydrazones, under usual conditions, and are hydrolysed to the corresponding keto-macrolides according to the procedure described by Pines et al. (Scheme).<sup>2</sup>



The reaction proceeds very vigorously with dihydropyran, affording a number of products which are under investigation. However, the reaction is comparatively slow and incomplete with the enol ether derived from cholestan-3-one.

Although both nitrosyl chloride and alkyl nitrites react with acyclic enol ethers, under anhydrous conditions, to afford the corresponding nitroso-adducts,3 the present hydrolytic nitrosation of the cyclic enol ethers with the subsequent fission of the C-C bond represents a novel reaction of possible general applicability.

(Received, 19th July 1972; Com. 1250.)

 $\dagger$  Enol ethers were prepared by slight modifications of the enamine + acrylate/reduction sequence used by Borowitz et al.<sup>1a</sup> ‡ Any alkyl nitrite or even sodium nitrite may be used.

§ All the compounds described here gave satisfactory elemental analyses and expected n.m.r. and i.r. spectra.

<sup>1</sup> (a) I. J. Borowitz, G. J. Williams, L. Gross, H. Beller, D. Kurland, N. Suciu, V. Bandurco, and R. D. G. Rigby, *J. Org. Chem.*, 1972, **37**, 581; I. J. Borowitz, G. J. Williams, L. Gross, and R. Rapp, *ibid.*, 1968, **33**, 2013; (b) I. J. Borowitz, G. Gonis, R. Kelsey, R. Rapp, and G. J. Williams, *ibid.*, 1966, **31**, 3062.

<sup>2</sup> S. H. Pines, J. M. Chemerda, and M. A. Kozlowski, J. Org. Chem., 1966, 31, 3446.
<sup>3</sup> S. Tchelitcheff, U.S.P. 267425-6/1954; Chem. Abs., 1955, 49, 6301h.