## A New Synthesis of Pteridines

By RAYMOND D. YOUSSEFYEH\* and A. KALMUS

(Faculty of Medicine, The School of Pharmacy, The Hebrew University, Jerusalem, Israel)

Summary Reaction of 4-amino-1,3-dimethyl-5-nitrosouracil with various amine salts led to the formation of the corresponding pteridines as well as purines.

THE condensation of 4-amino-5-nitrosopyrimidine with compounds which possess an active methylene group adjacent to a functional group was first developed by Timmis<sup>1</sup> as a route to pteridine formation. Further



pteridines as well as purines when a 4-amino-5-nitrosopyrimidine is fused with a suitable amine salt. The yield of pteridine increases in parallel with the activity of  $\beta$ -methylene group in the amine salt. Thus, fusion of 4-amino-1,3-dimethyl-5-nitrosouracil (I) with 2-phenylethylamine hydrochloride (IIa) at 180° led to 41% of 1,3-dimethyl-6-phenyl-lumazine (IIIa)† [m.p. 257-258° (lit.3 m.p. 258-259°)] and 11% of 8-benzyltheophylline (IVa) [m.p. 298-299° (lit.4 m.p. 298-300°)]; (I) and n-propylamine hydrochloride (IIb) yielded 19% of 6-methyl-1,3dimethyl-lumazine (IIIb),<sup>5</sup> 37% of 8-ethyltheophylline (IVb),<sup>6</sup> and 25% of pyrimido-pteridine (V).<sup>7</sup>

We now describe the ready formation of the corresponding

The procedure involves the fusion of a mixture of thoroughly mixed pyrimidine and an excess of amine salt at 170-180°, followed by dilution with water and extraction with dichloromethane. After drying and removal of the solvent, the residue is chromatographed over neutral alumina. By this method, (I) with 2-methoxyethylamine hydrochloride (IIc) led to 1,3-dimethyl-lumazine (IIIc)\* and pyrimido-pteridine (V).7 These same products in addition to 8-methyltheophylline (IVc)<sup>6</sup> were obtained from the fusion of (I) and ethylamine hydrochloride (IId).

The formation of (V) from (I) by thermal or acid induced reaction has been reported in recent years.<sup>7</sup>

We thank Mr. M. Weisz for technical assistance.

(Received, October 13th, 1969; Com. 1542.)

studies<sup>2</sup> demonstrated that purines are the sole products if the methylene component is not activated and the condensation with nitroso-group is intramolecular.

† Satisfactory analysis, n.m.r., u.v., and i.r. spectral data are obtained for all the products.

<sup>1</sup>G. M. Timmis, Nature, 1949, 164, 139; U.S.P. 2,581,889/1952.
<sup>2</sup>H. Goldner, G. Dietz, and E. Carstens, Naturviss., 1964, 51, 137; Tetrahedron Letters, 1965, 2701; Annalen, 1966, 691, 142; ibid., 1966, 692, 134; G.P. 1,245,969 and 1,249,871/1967; E. Buhler and W. Pfleiderer, Angew. Chem. Internat. Edn., 1965, 77, 146.
<sup>3</sup> R. B. Angier, J. Org. Chem., 1963, 28, 1398.
<sup>4</sup> G. P. Hager, J. C. Krantz, and J. B. Harmon, J. Amer. Pharmaceut. Assoc. (Sci. Edn.), 1954, 43, 148.
<sup>5</sup> H. Zondler, H. S. Forest, and J. M. Logowski, J. Heterocyclic Chem., 1967, 4, 124.
<sup>6</sup> J. H. Speer and A. L. Raymond, J. Amer. Chem. Soc., 1953, 75, 114.
<sup>7</sup> E. C. Taylor and E. E. Garcia, J. Amer. Chem. Soc., 1964, 86, 4721 and references therein.
<sup>8</sup> W. Pfleiderer, Chem. Ber., 1957, 90, 2582.