

• Operationally simple

• 2° and 3° Boronic Esters

Chan–Lam Amination of Secondary and Tertiary Benzylic Boronic Esters

James D. Grayson, Francesca M. Dennis, Craig C. Robertson, and Benjamin M. Partridge*



metalation from B to Cu occurs through a single-electron, rather than a two-electron process.

A dvances in catalysis have enabled chemists to prepare aryl amines in a reliable and predictable manner, using methods such as the Buchwald–Hartwig,¹ Chan–Lam,² and Ullmann reactions.^{1c,2b,3} These reactions allow amine formation from common chemical building blocks (e.g., aryl halides or arylboronic acids) and are typically operationally simple to carry out. As a result, they have found application in a range of disciplines from drug discovery and manufacture to materials science.^{1b}

While highly desirable, the corresponding coupling methods that form alkyl C–N bonds are underdeveloped.⁴ For Pd catalysis, the reductive elimination step to form alkyl–N bonds is challenging.⁵ More progress has been made using Cu catalysis, through the coupling of amines with alkyl radicals, generated under photoredox catalysis from alkyl halides⁶ and carboxylic acids.⁷

Because of the versatility of organoboron reagents, we are interested in developing an alkyl variant of the Chan–Lam reaction (Scheme 1). This reaction has been studied in detail for





the coupling of cyclopropyl boronic acid,⁸ but the reaction of alkylboronic esters has only been rarely reported.⁹ Advances in this area would complement traditional methods to form alkyl amines, such as nucleophilic substitution and reductive amination.¹⁰ These often suffer from the necessity of protecting group strategies, over-alkylation, and, in the case of nucleophilic substitution, the need for toxic alkylating agents.

C(sp³)-N coupling

Herein, we report a Chan–Lam coupling of secondary and tertiary benzylic boronic esters with anilines. This complements a recent report of Cu-catalyzed decarboxylative amination of benzylic carboxylic acids which was proposed to occur through a Chan–Lam type mechanism.^{11,12}

As part of the ongoing investigation into Cu-catalyzed oxidative transformations of boronic esters,¹³ we explored the reaction of boronic ester 1 with aniline 2a in the presence of $Cu(OAc)_2$ and Cs_2CO_3 (see Table 1).¹⁴ Heating these reagents in a mixture of MeOH and pyridine at 50 $^\circ\text{C}$ under air gave amine 3a, alongside oxidation products 4 and 5 which are consistent with our previous observations.¹³ The remainder of the mass is likely to be accounted for by protodeboronation of boronic ester 1 to give ethylbenzene, whose volatility means it was unable to be detected after workup. Reducing the loading of base led to an increase in amine 3a and a decrease in the amount of alcohol 5 formed. However, in the absence of Cs_2CO_3 , formation of ketone 4 dominated. Pleasingly, oxidation side products could be minimized by conducting the reaction under an inert atmosphere. However, this requires the use of stoichiometric loadings of $Cu(OAc)_2$, to act as both catalyst

 Received:
 April 26, 2021

 Published:
 June 25, 2021





© 2021 The Authors. Published by American Chemical Society

Table 1. Evaluation of Reaction Conditions on the Yield of 3a⁴



	equiv [Cu]	equiv 2a	MeOH:Pyr	equiv Cs ₂ CO ₃	yield (%) ⁶			
entry					1	3a	4	5
1 ^c	2	2	3:2	2	0	16	34	48
2 ^{<i>c</i>}	2	2	3:2	0.5	0	30	32	5
3 ^c	2	2	3:2		0	6	60	<5
4	2	2	3:2	0.5	0	39	6	6
5	2	4	3:2	0.5	23	48	<5	6
6	2	4	3:1	0.5	51	38	<5	<5
$7^{d,e}$	2	4	3:1	0.5		73 ^f		
8 ^{<i>d</i>,<i>e</i>}	1	2	3:1	0.5		36 ^f		
9 ^{<i>d</i>,<i>e</i>}	1.5	3	3:1	0.5		67 ^f		
10 ^{<i>d</i>,<i>e</i>}	2	1	3:1	0.5		46 ^f		
11^d	0	4	3:1	0.5	n.d.	0	<5	33
12^d	2	4	1:0	0.5	45	37	<5	0

^{*a*}Reactions performed using 0.05 mmol of 1a. ^{*b*}Determined using ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. ^{*c*}Reaction conducted under air. ^{*d*}16 h reaction time. ^{*e*}Reactions performed using 0.50 mmol of 1a. ^{*f*}Yield is of isolated material. n.d. = not determined.

and oxidant in the reaction. Because of the relatively low cost and benign safety profile of $Cu(OAc)_2$, we chose this compromise rather than using other chemical oxidants that have been used previously in Chan-Lam reactions.^{9a,c,15} Increasing the amount of aniline 2a led to an improved yield of 3a, with a ratio of $2a:Cu(OAc)_2$ of 2:1 found to be best. Lowering the amount of pyridine in the solvent mixture gave a better mass balance, presumably due to lower levels of protodeboronation. Finally, extending the reaction time led to complete conversion of boronic ester 1, giving amine 3a in excellent yield. A series of control experiments showed the necessity of $Cu(OAc)_2$ for amination to occur (entry 11). In the absence of pyridine, the efficiency of amination is reduced (entry 12). This suggests that pyridine acts to break up aggregates of $Cu(OAc)_{2}$, providing a higher concentration of active catalyst in solution.¹⁶ Switching the stoichiometry to make the amine the limiting reagent did lead to coupling, albeit in moderate yield (entry 10). However, these conditions would allow an isolable amount of coupling product to be obtained if the aniline substrate is prohibitively expensive.

Scheme 2 details the scope of the reaction of boronic ester 1 with a variety of anilines. In all cases, only mono-substitution of the aniline was observed. There is a clear trend that electron rich anilines react in higher yield over a 16 h time frame. Anilines substituted with electron withdrawing groups did react successfully but often required higher temperatures and prolonged reaction times for acceptable yields. Functional groups tolerated include alkenes (3j), amides (3g), aryl halides (3e, 3k, 3n), esters (3h), sulfides (3d), and trifluoromethyl groups (3f, 3i). Importantly, no Ullmann coupling was observed when using substrates containing an aryl bromide or chloride.³ C-N coupling also occurred exclusively at the aniline in the presence of a primary amide, which complements reports of amide coupling from Watson and co-workers.^{9a,17} Heteroaromatic anilines were successfully coupled in modest to good yield (3p, 3q). Secondary anilines can also be coupled successfully, with examples including methyl- and isopropylsubstituted anilines (3r, 3s), and tetrahydroquinoline (3t).

However, amine 3b, when resubjected to the coupling conditions, reacted in low conversion after 48 h (3u). This suggests why only the product of monoalkylation is typically observed as the rate of the second alkylation is much slower by comparison. We have briefly explored the generality of our conditions with alkylamines. Morpholine did undergo coupling to form amine 3v, though with low efficiency. However, primary alkyl amines 6 and 7 were found to be unreactive.

We next explored the scope of the boronic ester using panisidine as the coupling partner (Scheme 3). Benzylic boronic esters containing both electron donating and electron withdrawing groups on the arene are tolerated. Trifluoromethylsubstituted boronic ester 8f, however, reacted slowly. Aryl halides and ortho substituents on the arene are also tolerated. Extension of the alkyl chain was possible, with groups containing ethers, azides, and alcohols tolerated (91–90). Primary benzylic boronic ester 8p reacted smoothly and only gave the product of monoalkylation. The reaction conditions were successfully applied to allylic boronic esters 8q and 8r to form allylic amines in good yield. The reaction of 2-methylallylboronic ester 8r with p-anisidine did give a mixture of mono- and dialkylated products, but this is the only boronic ester for which we have observed dialkylation. In comparison, aliphatic boronic ester 8s showed low reactivity under these conditions.

Methods using tertiary boronic esters to form C–N bonds are currently limited to the preparation of primary amines.¹⁸ Using our conditions, we were pleased to observe the successful coupling of boronic ester **10a** with a range of primary anilines (Scheme 4). This is a rare example of a tertiary boronic ester reacting under transition metal catalysis.¹⁹ Unlike secondary boronic esters, no reaction occurred between *N*-methylaniline and boronic ester **10a**, presumably due to the increased steric congestion from these substrates. Interestingly, amination of dibenzylboronic esters proceeded at room temperature, except CF₃-containing boronic ester (**11f**). Monobenzylic tertiary boronic ester, however, required a reaction temperature of 50 °C to give **11g** in reasonable yield within 16 h. In contrast, *tert*-butyl pinacol boronic ester was unreactive under the reaction Scheme 2. Scope of Anilines in the Amination of Boronic Ester 1^a



^aReactions were conducted on a 0.5 mmol scale. Yields reported are of isolated material of the corresponding product. ^bOxidation of remaining boronic ester using H₂O₂/NaOH was performed prior to isolation of the amine product. "Reaction conducted at 65 °C. ^dReaction time of 48 h. ^eReaction time of 36 h.

conditions. While the yields of coupling are generally lower than with secondary boronic esters, the method can produce useful amounts of these otherwise hard to make C-tertiary amines.

Interestingly, when *p*-anisidine was reacted with tertiary boronic ester 13, alongside C-N coupling product 14, we also observed a product of C-C coupling (15, isolated as thiourea 16, Scheme 5). While this reactivity was not seen between panisidine and secondary boronic esters, a reaction of 4aminophenol with 1 did give a mixture of products from C-N (17), C-O (18), and C-C (19) coupling. Single isomers of both 15 and 19 were observed, with X-ray crystallography of the corresponding thiourea of 19 confirming arylation occurs ortho to nitrogen.²⁰ We have assigned 16 through analogy. The selectivity is consistent with electrophilic aromatic substitution reactions, suggesting the formation of 15 and 19 may involve a carbocation intermediate.



^{*a*}Yields reported are of isolated material. ^{*b*}Reaction conducted at 65 °C. CReaction time of 48 h. Reaction time of 32 h. Reaction conducted with N-methylaniline instead of p-anisidine.







^aReactions were conducted on a 0.5 mmol scale. Yields reported are of isolated material. ^bReaction conducted at 50 °C.

We wanted to learn more about the reaction mechanism, as little information is available about the reaction of alkylboronic

pubs.acs.org/joc Scheme 3. Scope of Boronic Esters in the Amination with p-

Note

pubs.acs.org/joc

Scheme 5. Observation of C-C Bond Formation^a



^{*a*}Reaction conditions for amination reactions: $Cu(OAc)_2$ (2 equiv), Cs_2CO_3 (0.5 equiv), MeOH/Pyr (3:1), 25 °C (reaction with 13) or 50 °C (reaction with 1). Yields reported are of isolated material. ^{*b*}Phenyl isothiocyanate (1.1 equiv), hexane:MeCN:Et₃N (1:0.1:0.1).

esters with Cu salts. We started by performing a radical clock experiment, using cyclopropane containing boronic ester **20** (eq 1). This reacted to give amine **21**, with ring opening of the cyclopropane, without observation of the corresponding benzyl amine **22**. This suggests that the boronic ester serves as a radical precursor,²¹ with a two-electron transmetalation to Cu unlikely. This is in contrast to our previous investigation into the Cucatalyzed oxidation of benzylic boronic esters, where oxidation of **20** occurred without ring opening of the cyclopropane.¹³ While we cannot rule out further oxidation of the radical to a carbocation, the observed product distribution indicates this is likely to be a minor pathway, due to stability of cyclopropylcarbinyl cations which would presumably promote formation of **22**.



Consistent with the formation of a benzyl radical is the reaction of enantiomerically enriched boronic ester (*S*)-1a (eq 2). This reacted smoothly to form amine 3a, but with complete loss of stereochemical information. However, we cannot rule out whether configurational instability of an alkyl-Cu intermediate is the cause of stereoablation.²²

In conclusion, we have developed an operationally simple Chan-Lam coupling of secondary and tertiary boronic esters with anilines. The amination reaction is a rare example of a transition-metal-mediated reaction using a tertiary alkylboron coupling partner. Monoalkylation of the amine occurs selectively, and the conditions tolerate a diverse range of functional groups. Initial investigation into the reaction mechanism suggests that transmetalation from boron to copper is likely to occur through a single-electron process. Further Cheteroatom bond forming processes from alkylboronic esters are under development and will be reported in due course.

EXPERIMENTAL SECTION

General. All reagents and solvents used were supplied by commercial sources without further purification unless specified. Aniline (distillation) and *p*-anisidine (recrystallization from ethanol) were purified before use. All air-sensitive reactions were carried out under a nitrogen or argon atmosphere using oven-dried apparatus. Anhydrous Et₂O, THF, and toluene were dried and purified by passage through activated alumina columns using a solvent purification system. All petroleum ether used was 40-60 °C petroleum ether. All reactions which required elevated temperatures were heated using a stirrer hot plate and oil bath with an external probe to control the temperature. Thin layer chromatography (TLC) was performed on aluminumbacked plates precoated with silica. Compounds were visualized by exposure to UV light or by dipping the plates into solutions of vanillin or KMnO₄, followed by heating. All flash chromatography was carried out using silica gel mesh 40-63. It should be noted that the time taken for chromatography of boronic esters should be kept to a minimum to avoid extensive decomposition and reduced yields.

Infrared spectra were recorded on a PerkinElmer 100 FT instrument on the neat compound. NMR spectra were recorded on Bruker Advance 400 and 500 instruments at the indicated 101, 128, 126, 377, 400, and 500 MHz as dilute solutions in the indicated deuterated solvent. NMR spectra were recorded at ambient temperature unless otherwise stated. All chemical shifts (δ) are reported in parts per million (ppm) relative to the residual protio solvent $(\delta H: CHCl_3 = 7.27 \text{ ppm})$, DMSO = 2.50 ppm, or CH₃CN = 1.94 ppm) or the solvent itself (δ C: CDCl₃ = 77.0 ppm, DMSO = 39.5 ppm, or CH₃CN = 1.32, 118.3 ppm). All multiplets are designated by the following abbreviations: s = singlet, br s = broad singlet, d = doublet, dt = doublet triplet, td = triplet doublet, ddd = doublet of doublets of doublets, q = quartet, br q = broad quartet, m = multiplet. All coupling constants (I) are reported in Hertz (Hz). Assignments of ¹H NMR signals were made using 2D NMR spectra, typically HSQC and HMBC experiments. ¹³C NMR data were acquired as DEPT-Q experiments as standard. For samples where quaternary carbons were not observed by DEPT-Q₁¹³C NMR spectra were acquired as decoupled spectra. ¹⁹F NMR spectra were acquired as decoupled spectra. High-resolution mass spectra were recorded using either electrospray ionization (ESI) or electron ionization (EI) by the Chemistry Mass Spectrometry Facility in the Faculty of Science, University of Sheffield. HPLC analysis was carried out using the Chromatography Facility at the Department of Chemistry, University of Sheffield. Melting points were measured using a Linkam HFs91 heating stage, used in conjunction with a TC92 controller, and are uncorrected.

Single crystal X-ray intensity data were collected at 100 K on a Bruker D8 Venture diffractometer equipped with a Photon 100 CMOS detector using a Cu K α microfocus X-ray source from crystals mounted in fomblin oil on a MiTiGen microloop and cooled in a stream of cold N₂. Data were corrected for absorption using empirical methods (SADABS)²³ based upon symmetry equivalent reflections combined with measurements at different azimuthal angles.²⁴ The crystal structures were solved and refined against F^2 values using ShelXT²⁵ for solution and ShelXL²⁶ for refinement accessed via the Olex2 program.²⁷ Non-hydrogen atoms were refined anisotropically. Hydrogen atoms were placed in calculated positions with idealized geometries and then refined by employing a riding model and isotropic displacement parameters.

Synthesis of Substrates. Boronic esters $(1, 8a-p, 20)^{13}$ and boronic esters $(8q, 8r)^{28}$ were prepared by literature methods. *3-Methoxypropyl N,N-Bis(propan-2-yl)carbamate (S1).*



N,*N*-Diisopropylcarbamoyl chloride (3.96 g, 24.2 mmol) was added to a solution of 3-methoxy propanol (2.2 mL, 23 mmol) and triethylamine (3.4 mL, 24 mmol) in CH₂Cl₂ (77 mL). The mixture was heated at reflux for 18 h, cooled to room temperature, and diluted with H₂O (30 mL). The mixture was extracted with DCM (3 × 20 mL). The combined organic layers were washed with brine (20 mL), dried (MgSO₄), filtered, and concentrated *in vacuo*. The crude material was purified by flash chromatography (10% EtOAc/petroleum ether) to give *carbamate* **S1** (3.88 g, 77%) as a yellow oil. **IR** 2969, 2932, 1687, 1435, 1289 cm⁻¹. ¹**H NMR** (400 MHz, CDCl₃) δ 4.16 (2H, t, *J* = 6.4 Hz, C(O)OCH₂), 4.13–3.55 (2H, m, 2 × CH₃CH), 3.47 (2H, t, *J* = 6.4 Hz, CH₂OCH₃), 3.34 (3H, s, OCH₃), 1.92 (2H, p, *J* = 6.4 Hz, OCH₂CH₂), 1.25–1.15 (12H, m, 4 × CHCH₃). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 155.7 (C), 69.6 (CH₂), 61.7 (CH₂), 58.6 (CH₃), 46.3 (br, 2 × CH), 29.4 (CH₂), 20.9 (4 × CH₃). **HRMS** (Q-TOF) *m/z*: [M + H]⁺ Calcd for C₁₁H₂₄NO₃⁺ 218.1756; found 218.1751.

(±)-tert-Butyl 3-[3-Methoxy-1-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)propyl]-1H-indole-1-carboxylate (81).



Using a modification of the procedure by Aggarwal and co-workers,²⁹ a Schlenck flask containing carbamate S1 (0.576 g, 2.65 mmol) was backfilled with nitrogen three times. TMEDA (0.55 mL, 3.7 mmol) and anhydrous Et₂O (10.2 mL) were added, and the mixture was cooled to -78 °C. s-BuLi (1.3 M in cyclohexane, 2.8 mL, 3.7 mmol) was added dropwise, and the mixture was stirred at -78 °C for 4 h. t-Butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-1-carboxylate (0.700 g, 2.04 mmol) was added dropwise. A solution of MgBr₂ in Et₂O (0.378 g, 2.04 mmol, 1 M) was added dropwise, and the mixture was stirred at 34 °C for 18 h. Toluene (10 mL) was added, and the mixture was heated to 90 °C for 18 h. H₂O (20 mL) and Et₂O (15 mL) were added, and the mixture was extracted with Et_2O (3 × 15 mL). The combined organic layers were dried (MgSO₄), filtered, and concentrated in vacuo. The crude material was purified by column chromatography (5% EtOAc/petroleum ether) to give boronic ester 81 (0.422 g, 50%) as a colorless oil. IR 2977, 2930, 1729, 1452, 1368, 1142 cm^{-1} . ¹**H** NMR (400 MHz, CDCl₃) δ 8.11 (s, 1H), 7.66–7.61 (1H, m, ArH), 7.41 (1H, s, ArH), 7.31–7.27 (1H, m, ArH), 7.24–7.17 (1H, m, ArH), 3.48-3.36 (2H, m, CH2OCH3), 3.33 (3H, s, OCH3), 2.66-2.57 (1H, m, BCH), 2.25–2.14 (1H, m, BCHCH_ACH_B), 2.05–1.93 (1H, m, BCHCH_ACH_B), 1.66 (9H, s, $3 \times C(O)OCCH_3$), 1.22 (6H, s, $2 \times$ BOCCH₃), 1.21 (6H, s, 2 × BOCCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 149.8 (C), 135.6 (C), 130.7 (C), 124.0 (CH), 122.3 (C), 122.0 (CH), 121.5 (CH), 119.6 (CH), 115.0 (CH), 83.5 (2 × C), 83.0 (C), 72.0 (CH₂), 58.5 (CH₃), 31.0 (CH₂), 28.2 (3 × CH₃), 24.7 (4 × CH₃), 18.4 (BCH). ¹¹B NMR (128 MHz, CDCl₃) δ 33.7. HRMS (Q-TOF) m/z: [M + Na]⁺ Calcd for C₂₃H₃₄BNNaO₅⁺ 438.2428; found 438.2422.

(*S*)-4,4,5,5-*Tetramethyl*-2-(1-*phenylethyl*)-1,3,2-*dioxaborolane* ((*S*)-1). Following the procedure by Yun and co-workers,³⁰ a mixture of CuCl (20.8 mg, 0.211 mmol), KOtBu (56.7 mg, 0.506 mmol), and (*R*)-DTBM-Segphos (250.0 mg, 0.211 mmol) in anhydrous toluene (5.0 mL) was stirred for 30 min under an atmosphere of nitrogen. Pinacolborane (10.1 mmol, 1.46 mL) was added to the reaction mixture and stirred for 10 min. Styrene (8.44 mmol, 0.97 mL) was added, and the mixture was stirred at 60 °C for 18 h. The reaction mixture was cooled, filtered through a pad of Celite eluting with EtOAc, and concentrated *in vacuo*. The product was purified by column chromatography (5% Et₂O/petroleum ether) to give boronic ester (*S*)-1 (1.10 g, 54%) as a colorless oil. The data were consistent with the literature.³¹ $[\alpha]_D^{21}$ +12.0 (*c* 1.00, CHCl₃); lit. $[\alpha]_D^{23}$ +11.9 (*c* 1.08, CHCl₃) 97:3 er. (*S*).^{32 1}H NMR (400 MHz, CDCl₃) δ 7.33–7.19 (4H, m, ArH), 7.19–7.08 (1H, m, ArH), 2.45 (1H, q, *J* = 7.5 Hz, CH), 1.34 (3H, d, J = 7.5 Hz, CHCH₃), 1.22 (6H, s, $2 \times CCH_3$), 1.21 (6H, s, $2 \times CCH_3$). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 144.9 (C), 128.3 ($2 \times CH$), 127.8 ($2 \times CH$), 125.1 (CH), 83.3 ($2 \times C$), 24.6 ($2 \times CH_3$), 24.6 ($2 \times CH_3$), 17.0 (CH₃). ¹¹B NMR (128 MHz, CDCl₃) δ 33.5. e.r. = 97:3, measured through chiral HPLC analysis of the corresponding alcohol obtained after oxidation. HPLC (Phenomenex Cellulose-1 column (250×4.6 mm), IPA/hexane 10/90, flow rate = 1.0 mL/min, l = 254 nm), $t_R = 6.2$ min (minor), 7.1 min (major).

General Procedure 1: Lithiation-Borylation. Using a modification of the procedure by Aggarwal and co-workers,²⁹ a Schlenk flask containing the corresponding carbamate (1.2 equiv) was backfilled with nitrogen three times. Anhydrous Et₂O (0.2 M) was added, and the mixture was cooled to -78 °C. *s*-BuLi (1.3 equiv) was added dropwise, and the mixture was stirred at -78 °C for 1 h. The boronic ester (1 equiv) was added dropwise, and the mixture was stirred at -78 °C for 1 h and then at room temperature for 2 h. H₂O (20 mL) and Et₂O (15 mL) were added, and the mixture was extracted with Et₂O (3 × 15 mL). The combined organic layers were dried (MgSO₄), filtered, and concentrated *in vacuo*.

(±)-2-[1-(4-Methoxyphenyl)-1-phenylethyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**10a**).



The title compound was prepared according to *General Procedure 1* using 1-phenylethyl diisopropylcarbamate³³ (1.60 g, 6.42 mmol), s-BuLi (1.3 M in cyclohexane, 5.6 mL, 7.3 mmol), and 4-methoxyphenylboronic acid pinacol ester (1.00 g, 4.27 mmol) added in Et₂O (2 mL). Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave boronic ester **10a** (1.25 g, 87%) as a white solid. The data were consistent with the literature.²⁹ m.p. 64–66 °C (petroleum ether); no literature value available. ¹H NMR (400 MHz, CDCl₃) δ 7.31–7.24 (2H, m, ArH), 7.25–7.20 (2H, m, ArH), 7.20–7.15 (3H, m, ArH), 6.86–6.80 (2H, m, ArH), 3.81 (3H, s, OCH₃), 1.68 (3H, s, CCH₃), 1.22 (12H, s, 4 × OCCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 157.3 (C), 148.1 (C), 139.4 (C), 129.4 (2 × CH), 128.4 (2 × CH), 127.9 (2 × CH), 125.3 (CH), 113.3 (2 × CH), 83.7 (2 × C), 55.1 (CH₃), 25.9 (CH₃), 24.4 (4 × CH₃). ¹¹B NMR (128 MHz, CDCl₃) δ 33.6.

(±)-2-[1-(4-Chlorophenyl)-1-phenylethyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**10b**).



The title compound was prepared according to *General Procedure 1* using 1-phenylethyl diisopropylcarbamate³³ (1.25 g, 5.03 mmol), *s*-BuLi (1.3 M in cyclohexane, 4.2 mL, 5.5 mmol), and 4-chlorophenylboronic acid pinacol ester (1.00 g, 4.19 mmol) added in Et₂O (2 mL). Flash chromatography (4% EtOAc/petroleum ether) of the crude material gave boronic ester **10b** (1.38 g, 96%) as an off-white solid. The data were consistent with the literature.²⁹ m.p. 74–75 °C (petroleum ether); no literature value available. ¹H NMR (400 MHz, CDCl₃) δ 7.30–7.27 (1H, m, ArH), 7.26–7.15 (8H, m, ArH), 1.67 (3H, s, CCH₃), 1.21 (6H, s, 2 × OCCH₃), 1.21 (6H, s, 2 × OCCH₃), 1.21 (6H, s, 2 × OCCH₃), 1.3C{¹H} NMR (101 MHz, CDCl₃) δ 147.2 (C), 146.2 (C), 131.1 (C), 129.9 (2 × CH), 128.4 (2 × CH), 128.1 (2 × CH), 128.0 (2 × CH), 125.6 (CH), 83.9 (2 × C), 25.7 (CH₃), 24.4 (4 × CH₃). ¹¹B NMR (128 MHz, CDCl₃) δ 33.1.

(±)-2-[1-(4-Trifluoromethylphenyl)-1-phenylethyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**10c**).



The title compound was prepared according to General Procedure 1 using 1-phenylethyl diisopropylcarbamate³³ (0.551 g, 2.21 mmol), s-BuLi (1.3 M in cyclohexane, 1.8 mL, 2.4 mmol), and 4trifluoromethlphenylboronic acid pinacol ester (0.50 g, 1.84 mmol) added in Et₂O (2 mL). Flash chromatography (4% EtOAc/petroleum ether) of the crude material gave boronic ester 10c (0.575 g, 83%) as an off white solid. m.p. 65–66 °C (petroleum ether). IR 2977, 2932, 1616, 1317, 1113 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.53–7.49 (2H, m, ArH), 7.36-7.28 (4H, m, ArH), 7.22-7.18 (3H, m, ArH), 1.71 (3H, s, CCH₃), 1.22 (6H, s, 2 × OCCH₃), 1.21 (6H, s, 2 × OCCH₃). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 152.2 (C), 146.6 (C), 128.8 (2 × CH), 128.4 (2 × CH), 128.2 (2 × CH), 127.6 (q, $J_{\rm F}$ = 32.2 Hz, C), 125.8 (CH), 124.8 (q, $J_F = 3.8$ Hz, 2 × CH), 124.4 (q, $J_F = 271.6$ Hz, CF₃), 84.0 (2 × C), 25.6 (CH₃), 24.5 (2 × CH₃), 24.4 (2 × CH₃). ¹¹B NMR (128 MHz, CDCl₃) δ 33.3. ¹⁹F NMR (377 MHz, CDCl₃) δ -62.2. **HRMS** (Q-TOF) m/z: $[M + H]^+$ Calcd for $C_{21}H_{25}BF_3O_2^+$ 377.1900; found 377.1894.

(±)-2-(2-Phenylbut-2-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (**10d**).



The title compound was prepared according to *General Procedure 1* using 1-phenylethyl diisopropylcarbamate³³ (1.92 g, 7.69 mmol), *s*-BuLi (1.3 M in cyclohexane, 6.4 mL, 8.3 mmol), and ethylboronic acid pinacol ester (1.00 g, 6.41 mmol). Flash chromatography (4% EtOAc/petroleum ether) of the crude material gave boronic ester **10d** (1.51 g, 90%) as a yellow oil. The data were consistent with the literature.^{29 1}H **NMR** (400 MHz, CDCl₃) δ 7.34–7.26 (4H, m, ArH), 7.17–7.11 (1H, m, ArH)), 1.89 (1H, dq, *J* = 14.7, 7.4 Hz, CCH_AH_B), 1.72 (1H, dq, *J* = 14.7, 7.4 Hz, CCH₄H_B), 1.22 (6H, s, 2 × OCCH₃), 0.84 (3H, t, *J* = 7.4 Hz, CH₂CH₃). ¹³C{¹H</sup> **NMR** (101 MHz, CDCl₃) δ 147.3 (C), 128.0 (2 × CH), 126.9 (2 × CH), 125.0 (CH), 83.2 (2 × C), 31.9 (CH₂), 24.6 (4 × CH₃), 21.0 (CH₃), 10.0 (CH₃). ¹¹B **NMR** (128 MHz, CDCl₃) δ 33.7.

(±)-2-(5-Azido-2-phenylpentan-2-yl)-4,4,5,5-tetramethyl-1,3,2dioxaborolane (13).



The title compound was prepared according to *General Procedure 1* using 1-phenylethyl diisopropylcarbamate³³ (1.19 g, 4.77 mmol), s-BuLi (1.3 M in cyclohexane, 4.2 mL, 5.4 mmol), and 2-(3-bromopropyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane³⁴ (0.700 g, 3.32 mmol). Flash chromatography (1% EtOAc/petroleum ether) of the crude material gave *boronic ester* **13** (0.540 g, 52%) as a colorless oil. **IR** 2977, 2932, 2092, 1350, 1312, 1145 cm⁻¹. ¹H **NMR** (400 MHz, CDCl₃) δ 7.33–7.28 (4H, m, ArH), 7.18–7.12 (1H, m, ArH), 3.25–3.17 (2H, m, CH₂N), 1.88–1.72 (2H, m, CH₂CH₂N), 1.55–1.42 (2H,

m, CCH₂), 1.36 (3H, s, BCCH₃), 1.22 (6H, s, $2 \times OCCH_3$), 1.21 (6H, s, $2 \times OCCH_3$). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 146.4 (C), 128.2 (2 × CH), 126.8 (2 × CH), 125.3 (CH), 83.5 (2 × C), 52.1 (CH₂), 36.5 (CH₂), 24.9 (CH₂), 24.6 (2 × CH₃), 24.5 (2 × CH₃), 21.3 (CH₃). ¹¹B NMR (128 MHz, CDCl₃) δ 33.7. HRMS (Q-TOF) *m/z*: [M + Na]⁺ Calcd for C₁₇H₂₆BN₃O₃Na⁺ 338.2016; found 338.2017.

Experimental Procedure for the Cu-Promoted Amination of Alkylboronic Esters. General Procedure 2: Preparative Scale Cu-Catalyzed Amination of Alkylboronic Esters. A flask containing the corresponding boronic ester (0.50 mmol, 1 equiv), aniline (2.00 mmol, 4 equiv), Cu(OAc)₂ (0.182 g, 1.00 mmol), and Cs₂CO₃ (0.0820 g, 0.252 mmol) was purged with argon. Methanol (1.0 mL) and pyridine (0.33 mL) were added, and the mixture was stirred at 50 or 65 °C until the reaction was complete (as determined by TLC). The mixture was cooled to room temperature, NH₄OH (10 mL) was added, and the mixture was extracted with Et₂O (3 × 10 mL), dried (MgSO₄), filtered, and concentrated *in vacuo*. The crude material was purified by column chromatography.

General Procedure 3: Preparative Scale Cu-Catalyzed Amination of Alkylboronic Esters with Oxidative Workup. A flask containing the corresponding boronic ester (0.50 mmol, 1 equiv), aniline (2.00 mmol, 4 equiv), $Cu(OAc)_2$ (0.182 g, 1.00 mmol), and Cs_2CO_3 (0.082 g, 0.25 mmol) was purged with argon. Methanol (1.0 mL) and pyridine (0.33 mL) were added, and the mixture was stirred at 50 or 65 °C until the reaction was complete (as determined by TLC). The mixture was cooled to room temperature, NH₄OH (10 mL) was added, and the mixture was extracted with Et₂O (3 × 10 mL), dried (MgSO₄), filtered, and concentrated *in vacuo*. THF (1 mL), H₂O (1 mL), and sodium perborate (0.382 g, 2.50 mmol) were added, and the mixture was stirred at RT for 1 h. The mixture was diluted with H₂O (10 mL) and extracted with EtOAc (3 × 10 mL). The combined organic phases were dried (MgSO₄), filtered, and concentrated *in vacuo*. The crude material was purified by column chromatography.

General Procedure 4: Preparative Scale Tertiary Alkyl Chan–Lam Coupling. A flask containing the corresponding boronic ester (0.50 mmol, 1 equiv), aniline (2.00 mmol, 4 equiv), $Cu(OAc)_2$ (0.182 g, 1.00 mmol), and Cs_2CO_3 (0.0820 g, 0.252 mmol) was purged with argon. Methanol (1.0 mL) and pyridine (0.33 mL) were added, and the mixture was stirred at RT for 16 h. NH₄OH (10 mL) was added, and the mixture was extracted with Et₂O (3 × 10 mL), dried (MgSO₄), filtered, and concentrated *in vacuo*. The crude material was purified by column chromatography.

(±)-*N*-(1-Phenylethyl)aniline (**3***a*). The title compound was prepared according to General Procedure 3 using boronic ester 1 (0.116 g, 0.501 mmol) and aniline (0.18 mL, 2.0 mmol), heating at 50 °C for 16 h. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave amine **3a** (0.0719 g, 73%) as a brown oil. The data were consistent with the literature.³⁵ ¹H NMR (400 MHz, CDCl₃) δ 7.41–7.36 (2H, m, ArH), 7.36–7.30 (2H, m, ArH), 7.26–7.21 (1H, m, ArH), 7.14–7.06 (2H, m, ArH), 6.69–6.61 (1H, m, ArH), 6.56–6.48 (2H, m, ArH), 4.50 (1H, q, J = 6.7 Hz, CH), 4.05 (1H, s, NH), 1.53 (3H, d, J = 6.7 Hz, CHCH₃). ¹³C{¹H}</sup> NMR (101 MHz, CDCl₃) δ 147.2 (C), 145.2 (C), 129.1 (2 × CH), 128.6 (2 × CH), 126.8 (CH), 125.8 (2 × CH), 117.2 (CH), 113.2 (2 × CH), 53.4 (CH), 25.0 (CH₃).

(±)-4-Methoxy-N-(1-phenylethyl)aniline (**3b**). The title compound was prepared according to *General Procedure* 2 using boronic ester **1** (0.116 g, 0.501 mmol) and *p*-anisidine (0.246 g, 2.00 mmol), heating at 50 °C for 16 h. Flash chromatography (4% EtOAc/petroleum ether) of the crude material gave amine **3b** (0.0823 g, 72%) as a yellow oil. The data were consistent with the literature.^{35 1}H NMR (400 MHz, CDCl₃) δ 7.45–7.39 (2H, m, ArH), 7.39–7.33 (2H, m, ArH), 7.39–7.33 (1H, m, ArH), 6.75 (2H, d, *J* = 8.9 Hz, ArH), 6.52 (2H, d, *J* = 8.9 Hz, ArH), 4.46 (1H, q, *J* = 6.7 Hz, CH), 3.82 (1H, s, NH), 3.74 (3H, s, OCH₃), 1.54 (3H, d, *J* = 6.7 Hz, CHCH₃).¹³C{¹H} NMR (101 MHz, CDCl₃) δ 151.8 (C), 145.4 (C), 141.5 (C), 128.5 (2 × CH), 126.8 (CH), 125.8 (2 × CH), 114.7 (2 × CH), 114.5 (2 × CH), 55.7 (CH₃), 54.2 (CH), 25.1 (CH₂).

(±)-4-Methyl-N-(1-phenylethyl)aniline (3c). The title compound was prepared according to *General Procedure 3* using boronic ester 1 (0.116 g, 0.499 mmol) and *p*-toluidine (0.216 g, 2.00 mmol), heating at

50 °C for 16 h. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave amine **3c** (0.0729 g, 69%) as a brown oil. The data were consistent with the literature.^{35 1}H NMR (400 MHz, CDCl₃) δ 7.41–7.34 (2H, m, ArH), 7.35–7.29 (2H, m, ArH), 7.25–7.18 (1H, m, ArH), 6.91 (2H, d, *J* = 8.3 Hz, ArH), 6.45 (2H, d, *J* = 8.3 Hz, ArH), 4.47 (1H, q, *J* = 6.7 Hz, NCH), 3.92 (1H, s, NH), 2.20 (3H, s, CCH₃), 1.51 (3H, d, *J* = 6.7 Hz, CHCH₃).¹³C¹H} NMR (101 MHz, CDCl₃) δ 145.4 (C), 145.0 (C), 129.6 (2 × CH), 128.6 (2 × CH), 126.8 (CH), 126.3 (C), 125.8 (2 × CH), 113.3 (2 × CH), 53.6 (CH), 25.1 (CH₃), 20.3 (CH₃).

(±)-4-(*Methylthio*)-*N*-(1-*phenylethyl*)*aniline* (**3***d*). The title compound was prepared according to *General Procedure* 2 using boronic ester 1 (0.115 g, 0.495 mmol) and 4-(methylthio)aniline (0.25 mL, 2.0 mmol), heating at 50 °C for 16 h. Flash chromatography (5% EtOAc/ petroleum ether) of the crude material gave amine **3d** (0.0743 g, 62%) as a brown oil. The data were consistent with the literature.³⁶ ¹H NMR (400 MHz, CDCl₃) δ 7.28–7.20 (4H, m, ArH), 7.17–7.12 (1H, m, ArH), 7.04 (2H, d, *J* = 8.7 Hz, ArH), 6.37 (2H, d, *J* = 8.7 Hz, ArH), 4.37 (1H, q, *J* = 6.7 Hz, NCH), 3.96 (1H, s, NH), 2.27 (3H, s, SCH₃), 1.42 (3H, d, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 146.1 (C), 144.9 (C), 131.3 (2 × CH), 128.7 (2 × CH), 126.9 (CH), 125.8 (2 × CH), 124.0 (C), 113.8 (2 × CH), 53.5 (CH), 25.0 (CH₃), 19.1 (CH₃).

(±)-4-Chloro-N-(1-phenylethyl)aniline (3e). The title compound was prepared according to *General Procedure* 3 using boronic ester 1 (0.116 g, 0.501 mmol) and 4-chloroaniline (0.256 g, 2.01 mmol), heating at 65 °C for 16 h. Flash chromatography (2% EtOAc/ petroleum ether) of the crude material gave amine 3e (0.0672g, 58%) as a brown oil. The data were consistent with the literature.^{35 1}H NMR (400 MHz, CDCl₃) δ 7.40–7.30 (4H, m, ArH), 7.27–7.21 (1H, m, ArH), 7.10–7.01 (2H, m, ArH), 6.52–6.32 (2H, m, ArH), 4.45 (1H, q, J = 6.7 Hz, NCH), 4.06 (1H, s, NH), 1.52 (2H, d, J = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 145.8 (C), 144.7 (C), 128.9 (2 × CH), 128.7 (2 × CH), 127.0 (CH), 125.8 (2 × CH), 121.8 (C), 114.4 (2 × CH), 53.6 (CH), 25.0 (CH₃).

(±)-α-Methyl-N-[4-(trifluoromethoxy)phenyl]-benzenemethanamine (**3f**). The title compound was prepared according to *General Procedure* 2 using boronic ester **1** (0.116 g, 0.499 mmol) and 4-(trifluoromethoxy)aniline (0.27 mL, 2.01 mmol), heating at 65 °C for 48 h. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave *amine* **3f** (0.0853g, 61%) as a yellow oil. **IR** 3420, 2968, 1613, 1512, 1248, 1153 cm⁻¹. ¹**H NMR** (400 MHz, CDCl₃) δ 7.38– 7.31 (4H, m, ArH), 7.27–7.24 (1H, m, ArH), 6.95 (2H, d, *J* = 8.9 Hz, ArH), 6.46 (2H, d, *J* = 8.9 Hz, ArH), 4.45 (1H, q, *J* = 6.9 Hz, NCH), 4.10 (1H, s, NH), 1.53 (3H, d, *J* = 6.9 Hz, CH₃). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 146.0 (C), 144.7 (C), 140.3 (C, q, *J_F* = 1.9 Hz), 128.7 (2 × CH), 127.1 (CH), 125.7 (2 × CH), 122.2 (2 × CH), 120.64 (OCF₃, q, *J_F* = 255.2 Hz), 113.4 (2 × CH), 53.7 (CH), 25.0 (CH₃). ¹⁹**F NMR** (377 MHz, CDCl₃) δ –58.5. **HRMS** (Q-TOF) *m*/*z*: [M + H]⁺ Calcd for C₁₅H₁₅F₃NO⁺ 282.1106; found 282.1095.

(±)-4-(1-Phenylethylamino)benzamide (**3***g*). The title compound was prepared according to a modification of General Procedure 3 using boronic ester 1 (0.117 g, 0.502 mmol) and 4-aminobenzamide (0.272 g, 2.01 mmol), heating at 65 °C for 36 h. The mixture was cooled to room temperature, Et₂O (5 mL) was added, and the mixture was passed through a plug of Celite. Flash chromatography (3% MeOH/DCM) of the crude material gave a mixture of amine 3g and pinacol (1:0.3, 0.0623 g). THF (1 mL), H₂O (1 mL), and sodium periodate (0.060 g, 0.281 mmol) were added, and the mixture was stirred at RT for 16 h. The mixture was diluted with H₂O (10 mL) and extracted with EtOAc $(3 \times 10 \text{ mL})$. The combined organic phases were dried (MgSO₄), filtered, and concentrated in vacuo. Flash chromatography (3% MeOH/ DCM) of the crude material gave amine 3g (0.0526 g, 44%) as an offwhite solid. The data were consistent with the literature.³⁷ m.p 153-155 °C (CH₂Cl₂). No literature value. ¹H NMR (400 MHz, DMSO) δ 7.53 (2H, d, J = 8.6 Hz, ArH), 7.46 (1H, s, NH), 7.39-7.33 (2H, m, ArH), 7.33-7.25 (2H, m, ArH), 7.22-7.14 (1H, m, ArH), 6.81 (1H, s, NH), 6.72–6.66 (1H, m, NH), 6.47 (2H, d, J = 8.6 Hz, ArH), 4.54 (1H, q, J = 6.8 Hz, NCH), 1.42 (3H, d, J = 6.8 Hz, CH₃). ¹³C{¹H} NMR (101 MHz, DMSO) δ 168.4 (C), 150.8 (C), 146.1 (C), 129.3 (2 ×

CH), 128.8 (2 × CH), 127.0 (CH), 126.3 (2 × CH), 121.4 (C), 111.9 (2 × CH), 52.2 (CH), 24.9 (CH₃).

(±)-Methyl 4-(1-Phenylethylamino)benzoate (3h). The title compound was prepared according to General Procedure 2 using boronic ester 1 (0.117 g, 0.503 mmol) and 4-methoxycarbonylaniline (0.302 g, 2.01 mmol), heating at 65 °C for 48 h. Flash chromatography (10% EtOAc/petroleum ether) of the crude material gave amine 3h (0.0648 g, 53%) as a white solid. The data were consistent with the literature.³⁸ m.p. 101–103 °C (petroleum ether); no literature value available. ¹H NMR (400 MHz, CDCl₃) δ 7.79 (2H, d, *J* = 8.8 Hz, ArH), 7.35–7.33 (4H, m, ArH), 7.27–7.22 (1H, m, ArH), 6.48 (2H, d, *J* = 8.8 Hz, ArH), 4.57 (1H, q, *J* = 6.6, NCH), 4.48 (1H, s, NH), 4.48 (1H, s, OCH₃), 1.56 (3H, d, *J* = 6.6 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 167.2 (C), 150.9 (C), 144.1 (C), 131.4 (2 × CH), 128.8 (2 × CH), 127.1 (CH), 125.7 (2 × CH), 118.3 (C), 112.1 (2 × CH), 53.0 (CH), 51.5 (CH₃), 24.7 (CH₃).

(±)-*N*-(1-Phenylethyl)-4-(trifluoromethyl)aniline (3i). The title compound was prepared according to *General Procedure 3* using boronic ester 1 (0.116 g, 0.501 mmol) and 4-trifluoromethylaniline (0.25 mL, 2.0 mmol), heating at 65 °C for 48 h. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave amine 3i (0.0479 g, 36%) as a colorless oil. The data were consistent with the literature.^{39 1}H NMR (400 MHz, CDCl₃) δ 7.39–7.34 (6H, m, ArH), 7.32–7.27 (1H, m, ArH), 6.57–6.53 (2H, m, ArH), 4.55 (1H, q, J = 6.7 Hz, NCH), 4.42 (1H, s, NH), 1.58 (3H, d, J = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 149.6 (C), 144.2 (C), 128.8 (2 × CH), 127.2 (CH), 126.4 (2 × CH, q, J_F = 3.8 Hz), 125.7 (2 × CH), 125.0 (C, q, J_F = 270.4 Hz), 118.6 (C, q, J_F = 32.5 Hz), 112.4 (2 × CH), 53.2 (CH), 24.8 (CH₃). ¹⁹F NMR (377 MHz, CDCl₃) δ –61.1.

(±)-N-(1-Phenylethyl)-4-(prop-2-en-1-yloxy)aniline (3j). The title compound was prepared according to General Procedure 2 using boronic ester 1 (0.116 g, 0.500 mmol) and 4-[(prop-2-en-1yl)oxy]aniline (0.299 g, 2.00 mmol), heating at 50 °C for 48 h. Flash chromatography (6% EtOAc/petroleum ether) of the crude material gave amine 3j (0.0863 g, 68%) as a yellow oil. IR 3404, 3025, 2866, 1508, 1229 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.36 (2H, m, ArH), 7.36–7.30 (2H, m, ArH), 7.26–7.21 (1H, m, ArH), 6.72 (2H, d, *J* = 8.9 Hz, ArH), 6.47 (2H, d, *J* = 8.9 Hz, ArH), 6.03 (1H, ddt, *J* = 17.2, 10.6, 5.4 Hz, CH=CH₂), 5.37 (1H, dd, J = 17.2, 1.5 Hz, CH= CH_AH_B), 5.24 (1H, dd, J = 10.6, 1.5 Hz, $CH=CH_AH_B$), 4.46–4.39 (3H, m, NCH, OCH₂), 3.80 (1H, s, NH), 1.51 (3H, d, J = 6.7 Hz, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 150.8 (C), 145.4 (C), 141.7 (C), 133.9 (CH), 128.6 (2 × CH), 126.8 (CH), 125.9 (2 × CH), 117.3 (CH₂), 115.8 (2 × CH), 114.4 (2 × CH), 69.6 (CH₂), 54.2 (CH), 25.1 (CH₃). HRMS (Q-TOF) m/z: [M + H]⁺ Calcd for C₁₇H₂₀NO⁺ 254.1545; found 254.1539.

(±)-3-Bromo-N-(1-phenylethyl)aniline (3k). The title compound was prepared according to *General Procedure* 3 using boronic ester 1 (0.116 g, 0.501 mmol) and 3-bromoaniline (0.344 g, 2.00 mmol), heating at 65 °C for 16 h. Flash chromatography (2% EtOAc/ petroleum ether) of the crude material gave amine 3k (0.0613 g, 44%) as a brown oil. The data were consistent with the literature.⁴⁰ ¹H NMR (400 MHz, CDCl₃) δ 7.38–7.31 (4H, m, ArH), 7.29–7.23 (1H, m, ArH), 6.96–6.90 (1H, m, ArH), 6.78–6.74 (1H, m, ArH), 6.70–6.67 (1H, m, ArH), 6.44–6.37 (1H, m, ArH), 4.47 (1H, q, J = 6.6 Hz, CH), 4.10 (1H, s, NH), 1.53 (3H, d, J = 6.6 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 148.4 (C), 144.4 (C), 130.4 (CH), 128.7 (2 × CH), 127.1 (CH), 125.7 (2 × CH), 123.0 (C), 120.0 (CH), 116.0 (CH), 111.8 (CH), 53.3 (CH), 24.8 (CH₃).

(±)-3-Methoxy-N-(1-phenylethyl)aniline (3I). The title compound was prepared according to *General Procedure 3* using boronic ester 1 (0.120 g, 0.519 mmol) and *m*-methoxyaniline (0.251 g, 2.04 mmol), heating at 50 °C for 48 h. Flash chromatography (5% EtOAc/ petroleum ether) of the crude material gave amine 3I (0.0552 g, 47%) as a yellow oil. The data were consistent with the literature.⁴¹ ¹H NMR (400 MHz, CDCl₃) δ 7.39–7.31 (4H, m, ArH), 7.24 (1H, t, *J* = 7.1 Hz, ArH), 7.01 (1H, t, *J* = 8.1 Hz, ArH), 6.23 (1H, dd, *J* = 8.1, 2.1 Hz, ArH), 6.16 (1H, dd, *J* = 8.1, 1.8 Hz, ArH), 6.08 (1H, dd, *J* = 2.1, 1.8 Hz, ArH), 4.49 (1H, q, *J* = 6.7 Hz, CH₃). ¹³C{¹H}</sup> NMR (101 MHz, CDCl₃) δ

160.6 (C), 148.6 (C), 145.1 (C), 129.8 (CH), 128.6 ($2 \times CH$), 126.9 (CH), 125.8 ($2 \times CH$), 106.4 (CH), 102.4 (CH), 99.3 (CH), 54.9 (CH₃), 53.5 (CH), 24.9 (CH₃).

(±)-2-Methoxy-N-(1-phenylethyl)aniline (**3m**). The title compound was prepared according to *General Procedure 3* using boronic ester **1** (0.116 g, 0.499 mmol) and *o*-methoxyaniline (0.23 mL, 2.0 mmol), heating at 50 °C for 16 h. Flash chromatography (4% EtOAc/petroleum ether) of the crude material gave amine **3m** (0.0801 g, 71%) as a colorless oil. The data were consistent with the literature.⁴² ¹H **NMR** (400 MHz, CDCl₃) δ 7.41–7.37 (2H, m, ArH), 7.36–7.30 (2H, m, ArH), 7.26–7.21 (1H, m, ArH), 6.81–6.76 (1H, m, ArH), 6.75–6.69 (1H, m, ArH), 6.66–6.59 (1H, m, ArH), 6.39–6.33 (1H, m, ArH), 4.65 (1H, s, NH), 4.50 (1H, q, *J* = 6.6 Hz, CH), 3.91 (3H, s, OCH₃), 1.57 (3H, d, *J* = 6.6 Hz, CHCH₃). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 146.5 (C), 145.4 (C), 137.2 (C), 128.6 (2 × CH), 126.7 (CH), 125.8 (2 × CH), 121.1 (CH), 116.3 (CH), 111.0 (CH), 109.2 (CH), 55.4 (CH₃), 53.3 (CH), 25.2 (CH₃).

(±)-4-Bromo-2-methoxy-N-(1-phenylethyl)aniline (3n). The title compound was prepared according to General Procedure 3 using boronic ester 1 (0.119 g, 0.514 mmol) and 4-bromo-2-methoxyaniline (0.409 g, 2.02 mmol), heating at 50 °C for 16 h. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave amine 3n (0.0843 g, 54%) as a light brown solid. m.p. 95-97 °C (petroleum ether). IR 3425, 2963, 2933, 1590, 1506, 1223 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.36-7.29 (4H, m, ArH), 7.26-7.20 (1H, m, ArH), 6.85 (1H, d, J = 2.1 Hz, ArH), 6.79 (1H, dd, J = 8.4, 2.1 Hz, ArH), 6.17 (1H, d, J = 8.4 Hz, ArH), 4.60 (1H, s, NH), 4.49-4.39 (1H, m, CH), 3.88 (3H, s, OCH₃), 1.55 (3H, d, J = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 147.1 (C), 144.8 (C), 136.2 (C), 128.6 (2 × CH), 126.9 (CH), 125.7 (2 × CH), 123.6 (CH), 112.6 (CH), 111.8 (CH), 107.7 (C), 55.7 (CH₃), 53.2 (CH), 25.1 (CH₃). HRMS (Q-TOF) m/z: $[M + H]^+$ Calcd for $C_{15}H_{18}^{79}BrNO^+$ 306.0494; found 306.0488.

(±)-*N*-(1-Phenylethyl)-2,6-dimethylbenzenamine (**3o**). The title compound was prepared according to *General Procedure* 3 using boronic ester **1** (0.116 g, 0.501 mmol) and 2,6-dimethylaniline (0.25 mL, 2.0 mmol), heating at 65 °C for 48 h. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave amine **3o** (0.0339 g, 30%) as a red oil. The data were consistent with the literature.⁴³ ¹H NMR (400 MHz, CDCl₃) δ 7.33–7.28 (4H, m, ArH), 7.26–7.21 (1H, m, ArH), 6.96 (2H, d, *J* = 7.4 Hz, ArH), 6.79 (1H, t, *J* = 7.4 Hz, ArH), 4.33 (1H, q, *J* = 6.7 Hz, NCH), 3.21 (1H, s, NH), 2.28 (6H, s, 2 × CCH₃), 1.52 (3H, d, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 145.3 (C), 144.9 (C), 129.4 (C), 128.8 (2 × CH), 128.4 (2 × CH), 126.9 (CH), 126.1 (2 × CH), 121.5 (CH), 56.7 (CH₃), 22.6 (CH), 18.9 (2 × CH₃).

(±)-1-Methyl-N-(1-phenylethyl)-1H-pyrazol-5-amine (**3p**). The title compound was prepared according to *General Procedure* 2 using boronic ester **1** (0.116 g, 0.499 mmol) and 1H-pyrazol-5-amine (0.194 g, 2.01 mmol), heating at 65 °C for 16 h. Flash chromatography (1:1 → 2:1 EtOAc/petroleum ether) of the crude material gave *amine* **3p** (0.0589 g, 59%) as a colorless oil. **IR** 3293, 2967, 2926, 1550, 1501 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.42–7.37 (2H, m, ArH), 7.35–7.30 (2H, m, ArH), 7.26–7.20 (1H, m, ArH), 7.00 (1H, d, *J* = 2.3 Hz, ArH), 5.29 (1H, d, *J* = 2.3 Hz, ArH), 4.46 (1H, q, *J* = 6.7 Hz, NCH), 4.07 (1H, s, NH), 3.68 (3H, s, NCH₃), 1.50 (3H, d, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 156.6 (C), 145.6 (C), 130.8 (CH), 128.4 (2 × CH), 126.8 (CH), 126.0 (2 × CH), 91.3 (CH), 54.7 (CH), 38.4 (CH₃), 24.5 (CH₃). HRMS (Q-TOF) *m*/*z*: [M + H]⁺ Calcd for C₁₂H₁₆N₃⁺ 202.1344; found 202.1339.

(±)-Methyl 3-[(1-Phenylethyl)amino]thiophene-2-carboxylate (**3q**). The title compound was prepared according to *General Procedure* 3 using boronic ester 1 (0.117 g, 0.504 mmol) and methyl 3-amino-2thiophenecarboxylate (0.316 g, 2.01 mmol), heating at 65 °C for 48 h. Flash chromatography (6% EtOAc/petroleum ether) of the crude material gave *amine* **3q** (0.0595 g, 45%) as a yellow oil. IR 3364, 2950, 1663, 1566, 1246 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.34–7.32 (4H, m, ArH), 7.25–7.23 (1H, m, ArH), 7.19–7.18 (1H, d, J = 5.4 Hz, ArH), 6.36 (1H, d, J = 5.4 Hz, ArH), 4.60 (1H, p, J = 6.8 Hz, NHCH), 3.86 (3H, s, OCH₃), 1.58 (3H, d, J = 6.8 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 165.5 (C), 155.4 (C), 144.9 (C), 132.0 (CH), 128.7 (2 × CH), 127.1 (CH), 125.7 (2 × CH), 117.2 (CH), 98.9 (C), 54.8 (CH₃), 51.1 (CH), 24.9 (CH₃). **HRMS** (Q-TOF) *m/z*: [M + H]⁺ Calcd for C₁₄H₁₆NO₂S⁺ 262.0896; found 262.0905.

(±)-*N*-*Methyl*-*N*-(1-phenylethyl)aniline (3r). The title compound was prepared according to *General Procedure* 2 using boronic ester 1 (0.116 g, 0.501 mmol) and *N*-methylaniline (0.22 mL, 2.0 mmol), heating at 65 °C for 16 h. Flash chromatography (2% EtOAc/ petroleum ether) of the crude material gave amine 3r (0.0877 g, 83%) as a yellow oil. The data were consistent with the literature.⁴⁴ ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.35 (4H, m, ArH), 7.33–7.27 (3H, m, ArH), 6.92–6.87 (2H, m, ArH), 6.81–6.75 (1H, m, ArH), 5.18 (1H, q, J = 6.9 Hz, CH), 2.73 (3H, s, NCH₃), 1.60 (3H, d, J = 6.9 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 150.2 (C), 142.8 (C), 129.2 (2 × CH), 128.4 (2 × CH), 126.9 (2 × CH), 126.8 (CH), 116.6 (CH), 113.0 (2 × CH), 56.5 (CH), 31.8 (CH₃), 16.3 (CH₃).

(±)-N-(1-Phenylethyl)-N-(propan-2-yl)aniline (3s). The title compound was prepared according to General Procedure 3 using boronic ester 1 (0.116 g, 0.498 mmol) and N-isopropylaniline (0.29 mL, 2.0 mmol), heating at 65 °C for 48 h. Flash chromatography (100% pentane) of the crude material gave amine 3s (0.0525 g, 44%) as a colorless oil. IR 2670, 2931, 1595, 1500, 1263 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.47–7.40 (2H, m, ArH), 7.38–7.30 (2H, m, ArH), 7.26-7.21 (1H, m, ArH), 7.20-7.11 (2H, m, ArH), 6.85-6.74 (3H, m, ArH), 4.79 (1H, q, J = 6.9 Hz, NCHAr), 3.89 (1H, sept, J = 6.7 Hz, CH $(CH_3)_2$, 1.53 (3H, d, J = 6.9 Hz, NCHCH₃), 1.20 (3H, d, J = 6.7 Hz, $CH(CH_3CH_3)$, 1.16 (3H, d, J = 6.7 Hz, $CH(CH_3CH_3)$). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 147.0 (C), 144.9 (C), 128.3 (2 × CH), 128.3 (2 × CH), 126.8 (CH), 126.3 (2 × CH), 119.5 (2 × CH), 118.4 (CH), 54.1 (CH), 48.7 (CH), 20.4 (CH₃), 20.3 (CH₃), 19.4 (CH₃). **HRMS** (Q-TOF) m/z: $[M + H]^+$ Calcd for $C_{17}H_{22}N^+$ 240.1752; found 240.1747

(±)-1-(1-Phenylethyl)-1,2,3,4-tetrahydroquinoline (**3t**). The title compound was prepared according to *General Procedure* 3 using boronic ester 1 (0.116 g, 0.499 mmol) and 1,2,3,4-tetrahydroquinoline (0.266 mg, 2.00 mmol), heating at 65 °C for 16 h. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave amine **3t** (0.0789 g, 67%) as a clear oil. The data were consistent with the literature.^{45 1}H NMR (400 MHz, CDCl₃) δ 7.40–7.33 (4H, m, ArH), 7.29–7.24 (1H, m, ArH), 7.08–7.03 (1H, m, ArH), 7.03–6.98 (1H, m, ArH), 6.75–6.69 (1H, m, ArH), 6.63–6.57 (1H, m, ArH), 5.16 (1H, q, *J* = 6.9 Hz, NCH), 3.21–3.14 (1H, m, NCH_ACH_B), 3.10–3.02 (1H, m, NCH_ACH_B), 2.86–2.74 (2H, m, ArCH₂), 1.94–1.85 (2H, m, NCH₂CH₂), 1.61 (3H, d, *J* = 6.9 Hz, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 145.6 (C), 142.7 (C), 129.2 (CH), 128.4 (2 × CH), 127.1 (CH), 126.9 (2 × CH), 126.7 (CH), 122.8 (C), 115.4 (CH), 110.6 (CH), \$4.6 (CH), 42.5 (CH₂), 28.5 (CH₂), 22.2 (CH₂), 16.0 (CH₃).

(±)-4-Methoxy-N,N-bis(1-phenylethyl)aniline (3u). The title compound was prepared according to General Procedure 2 using boronic ester 1 (0.0603 g, 0.260 mmol) and (\pm) -4-methoxy-N-(1-phenylethyl)aniline (3b) (0.226 g, 1.00 mmol), heating at 50 $^\circ C$ for 48 h. Flash chromatography (5% EtOAc/petroleum ether) of the crude material (d.r. 1:1) gave amine **3u** (0.0016 g, 18%, isolated d.r = 1.8:1 A:B) as a yellow oil, and amine 3b was recovered (182.8 mg, 80%). Upon further purification, diastereomer B was isolated as a single diastereomer. IR 3028, 2971, 1507, 1243, 1036, 699 cm⁻¹. ¹H NMR (diastereomer A) (400 MHz, CDCl₃) 7.33-7.29 (4H, m, ArH), 7.25-7.19 (6H, m, ArH), 6.73–6.71 (4H, m, ArH), 4.26 (2H, q, J = 6.8 Hz, 2 × CH), 3.80 $(3H, s, OCH_3)$, 1.18 (6H, d, J = 6.8 Hz, 2 × CH₃). ¹H NMR (diastereomer B) (400 MHz, $CDCl_3$) δ 7.30–7.29 (8H, m, ArH), 7.24-7.20 (2H, m, ArH), 6.71 (2H, d, J = 9.0 Hz, ArH), 6.65 (2H, d, J =9.0 Hz, ArH), 4.45 (2H, q, J = 6.7 Hz, 2 × CH), 3.74 (3H, s, CH₃), 1.27 (6H, d, J = 6.7 Hz, $2 \times CH_3$). ¹³C{¹H} NMR (mixed diastereomer A +B) (101 MHz, CDCl₃) δ 156.4 (C), 155.5 (C), 145.0 (2 × C), 143.7 (2 × C), 138.9 (C), 138.0 (C), 130.8 (CH), 128.7 (2 × CH), 128.0 (4 × CH), 128.0 (2 × CH), 127.9 (4 × CH), 127.8 (2 × CH), 126.7 (CH), 126.6 (CH), 112.8 (2 × CH), 112.5 (2 × CH), 58.3 (CH), 57.9 (CH), 55.2 (CH₃), 22.0 (CH₃), 18.2 (CH₃). HRMS (Q-TOF) m/z: $[M + H]^+$ Calcd for C₂₃H₂₅NO⁺ 332.2009; found 332.2007.

(±)-4-(1-Phenylethyl)-morpholine (**3v**). The title compound was prepared according to *General Procedure 2* using boronic ester 1 (0.116 g, 0.502 mmol) and morpholine (0.18 mL, 2.0 mmol), heating at 65 °C for 48 h. Flash chromatography (5–10% EtOAc/petroleum ether) of the crude material gave amine **3v** (0.0301 g, 31%) as a yellow oil. The data were consistent with the literature.^{46 1}H NMR (400 MHz, CDCl₃) δ 7.34–7.29 (4H, m, ArH), 7.26–7.21 (1H, m, ArH), 3.73–3.66 (4H, m, 2 × OCH₂), 3.31 (1H, q, *J* = 6.6 Hz, NCH), 2.56–2.44 (2H, m, 2 × NCH_AH_B), 2.41–2.34 (2H, m, 2 × NCH_AH_B), 1.36 (3H, d, *J* = 6.6 Hz, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 143.9 (C), 128.3 (2 × CH), 127.6 (2 × CH), 127.0 (CH), 67.2 (2 × CH₂), 65.4 (H), 51.3 (2 × CH₂), 19.8 (CH₃).

(±)-4-Methoxy-N-(1-(4-methoxyphenyl)ethyl)aniline (**9a**). The title compound was prepared according to *General Procedure* 2 using boronic ester **8a** (0.143 g, 0.507 mmol) and *p*-anisidine (0.249 g, 2.02 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine **9a** (0.0899 g, 69%) as a yellow oil. The data were consistent with the literature.³⁵ ¹H NMR (400 MHz, CDCl₃) δ 7.29 (2H, d, *J* = 8.7 Hz, ArH), 6.87 (2H, d, *J* = 8.7 Hz, ArH), 6.71 (2H, d, *J* = 8.9 Hz, ArH), 6.49 (2H, d, *J* = 8.9 Hz, ArH), 4.39 (1H, q, *J* = 6.7 Hz, CH), 3.80 (3H, s, OCH₃), 3.71 (3H, s, OCH₃), 1.48 (3H, d, *J* = 6.7 Hz, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 158.4 (C), 151.8 (C), 141.6 (C), 137.5 (C), 126.9 (2 × CH), 114.7 (2 × CH), 114.6 (2 × CH), 113.9 (2 × CH), 55.7 (CH₃), 55.2 (CH₃), 53.6 (CH), 25.1 (CH₃).

(±)-4-Methoxy-N-(1-p-tolylethyl)aniline (**9b**). The title compound was prepared according to *General Procedure* 2 using boronic ester **8b** (0.123 g, 0.500 mmol) and *p*-anisidine (0.246 g, 2.00 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/petroleum ether) of the crude material gave amine **9b** (0.0853 g, 71%) as an orange oil. The data were consistent with the literature.^{35 1}H NMR (400 MHz, CDCl₃) δ 7.26 (2H, d, *J* = 7.9 Hz, ArH), 7.14 (2H, d, *J* = 7.9 Hz, ArH), 6.70 (2H, d, *J* = 8.9 Hz, ArH), 6.49 (2H, d, *J* = 8.9 Hz, ArH), 4.40 (1H, q, *J* = 6.7 Hz, NCH), 3.75 (1H, br. s, NH), 3.71 (3H, s, OCH₃), 2.33 (3H, s, CCH₃), 1.49 (3H, d, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 151.8 (C), 142.4 (C), 141.6 (C), 136.3 (C), 129.3 (2 × CH), 125.8 (2 × CH), 114.7 (2 × CH), 114.5 (2 × CH), 55.7 (CH₃), 53.9 (CH), 25.2 (CH₃), 21.1 (CH₃).

(±)-N-(1-(Biphenyl-4-yl)ethyl)-4-methoxyaniline (9c). The title compound was prepared according to General Procedure 2 using boronic ester 8c (0.156 g, 0.507 mmol) and p-anisidine (0.247 g, 2.01 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine 9c (0.0975 g, 63%) as an orange solid. The data were consistent with the literature.⁴ m.p. 85-86 °C (petroleum ether); no literature value available. ¹H NMR (400 MHz, CDCl₃) δ 7.64-7.50 (4H, m, ArH), 7.48-7.38 (4H, m, ArH), 7.37–7.31 (1H, m, ArH), 6.72 (2H, d, J = 8.9 Hz, ArH), 6.51 (2H, d, J = 8.9 Hz, ArH), 4.47 (1H, q, J = 6.7 Hz, NCH), 3.79 (1H, s, NH), 3.71 (3H, s, OCH₃), 1.55 (3H, d, J = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.0 (C), 144.6 (C), 141.6 (C), 141.0 (C), 139.7 (C), 128.7 (2 × CH), 127.4 (2 × CH), 127.1 (CH), 127.0 (2 \times CH), 126.3 (2 \times CH), 114.8 (2 \times CH), 114.6 (2 \times CH), 55.7 (CH₃), 54.0 (CH), 25.1 (CH₃).

(±)-*N*-(1-(4-Fluorophenyl)ethyl)-4-methoxyaniline (**9d**). The title compound was prepared according to *General Procedure* 2 using boronic ester **8d** (0.122 g, 0.488 mmol) and *p*-anisidine (0.246 g, 2.00 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine **9d** (0.0839 g, 70%) as an orange oil. The data were consistent with the literature.^{35 1}H **NMR** (400 MHz, CDCl₃) δ 7.36–7.30 (2H, m, ArH), 7.04–6.97 (2H, m, ArH), 6.70 (2H, d, *J* = 8.7 Hz, ArH), 6.45 (2H, d, *J* = 8.7 Hz, ArH), 4.40 (1H, q, *J* = 6.7 Hz, CH), 3.77 (1H, s, NH), 3.71 (3H, s, OCH₃), 1.48 (3H, d, *J* = 6.7 Hz, CHCH₃).¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 161.7 (C, d, *J_F* = 244.1 Hz), 151.9 (C), 141.1 (C, d, *J_F* = 3.0 Hz), 127.3 (2 × CH, d, *J_F* = 8.0 Hz), 115.4 (2 × CH, *J_F* = 21.3 Hz), 114.7 (2 × CH), 114.5 (2 × CH), 55.7 (CH₃), 53.6 (CH), 25.3 (CH₃). ¹⁹F **NMR** (377 MHz, CDCl₃) δ –116.3.

(±)-4-Methoxy-N-(1-4-chlorophenylethyl)aniline (9e). The title compound was prepared according to *General Procedure 2* using boronic ester 8e (0.133 g, 0.499 mmol) and *p*-anisidine (0.246 g, 2.00

pubs.acs.org/joc

Note

mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine **9e** (0.0729 g, 56%) as an orange oil. The data were consistent with the literature.⁴⁸ ¹H **NMR** (400 MHz, CDCl₃) δ 7.33–7.27 (4H, m, ArH), 6.70 (2H, d, *J* = 8.9 Hz, ArH), 6.44 (2H, d, *J* = 8.9 Hz, ArH), 4.39 (1H, q, *J* = 6.7 Hz, NCH), 3.70 (3H, s, OCH₃), 1.48 (3H, d, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 152.1 (C), 144.1 (C), 141.3 (C), 132.4 (C), 128.8 (2 × CH), 127.3 (2 × CH), 114.8 (2 × CH), 114.6 (2 × CH), 55.7 (CH₃), 53.8 (CH), 25.1 (CH₃).

(±)-4-Methoxy-N-(1-(4-(trifluoromethyl)phenyl)ethyl)aniline (**9f**). The title compound was prepared according to *General Procedure* 2 using boronic ester **8f** (0.148 g, 0.493 mmol) and *p*-anisidine (0.249 g, 2.02 mmol), heating at 65 °C for 48 h. Flash chromatography (6% EtOAc/petroleum ether) of the crude material gave amine **9f** (0.0609 g, 42%) as a yellow oil. The data were consistent with the literature.^{35 1}H **NMR** (400 MHz, CDCl₃) δ 7.58 (2H, d, *J* = 8.2 Hz, ArH), 7.49 (d, *J* = 8.2 Hz, 2H), 6.70 (d, *J* = 8.9 Hz, 2H), 6.43 (d, *J* = 8.9 Hz, 2H), 4.46 (q, *J* = 6.8 Hz, 1H), 3.70 (s, 3H), 1.51 (d, *J* = 6.8 Hz, 3H). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 152.2 (C), 149.8 (C), 141.0 9 (C), 129.1 (C, q, *J*_F = 32.3 Hz), 126.2 (2 × CH), 125.6 (2 × CH q, *J*_F = 3.7 Hz), 124.2 (C, q, *J*_F = 270.4 Hz), 114.8 (2 × CH), 114.5 (2 × CH), 55.7 (CH₃), 54.0 (CH), 25.1 (CH₃). ¹⁹F **NMR** (377 MHz, CDCl₃) δ -62.4.

(±)-*N*-[1-(3-Chlorophenyl)ethyl]-4-methoxyaniline (**9g**). The title compound was prepared according to *General Procedure* 2 using boronic ester **8g** (0.133 g, 0.497 mmol) and *p*-anisidine (0.247 g, 2.00 mmol), heating at 65 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine **9g** (0.0792 g, 61%) as an orange oil. The data were consistent with the literature.⁴⁸ ¹H **NMR** (400 MHz, CDCl₃) δ 7.38 (1H, s, ArH), 7.27–7.24 (2H, m, ArH), 7.24–7.19 (1H, m, ArH), 6.71 (2H, d, *J* = 8.8 Hz, ArH), 6.46 (2H, d, *J* = 8.8 Hz, ArH), 4.38 (1H, q, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 152.1 (C), 147.9 (C), 141.2 (C), 134.5 (C), 129.9 (CH), 127.0 (CH), 126.1 (CH), 124.1 (CH), 114.8 (2 × CH), 114.5 (2 × CH), 55.7 (CH₃), 54.0 (CH), 25.1 (CH₃).

(±)-*N*-[1-(3-Bromophenyl)ethyl]-4-methoxyaniline (9h). The title compound was prepared according to *General Procedure* 2 using boronic ester 8h (0.153 g, 0.490 mmol) and *p*-anisidine (0.252 g, 2.04 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine 9h (0.0738 g, 49%) as a brown oil. The data were consistent with the literature.⁴⁹ ¹H NMR (400 MHz, CDCl₃) δ 7.53 (1H, s, ArH), 7.36 (1H, d, *J* = 7.7 Hz, ArH), 7.30 (1H, d, *J* = 7.7 Hz, ArH), 7.19 (t, *J* = 7.7 Hz, ArH), 6.71 (2H, d, *J* = 8.9 Hz, ArH), 6.45 (2H, d, *J* = 8.9 Hz, ArH), 4.37 (1H, q, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.1 (C), 148.2 (C), 141.1 (C), 130.2 (CH), 130.0 (CH), 129.0 (CH), 124.5 (CH), 122.8 (C), 114.8 (2 × CH), 114.5 (2 × CH), 55.7 (CH₃), 54.0 (CH), 25.2 (CH₃).

(±)-4-Methoxy-N-(1-(3-methoxyphenyl)ethyl)aniline (9i). The title compound was prepared according to *General Procedure* 2 using boronic ester 8i (0.129 g, 0.493 mmol) and *p*-anisidine (0.246 g, 2.03 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine 9i (0.0998 g, 79%) as a yellow oil. The data were consistent with the literature.⁴² ¹H NMR (400 MHz, CDCl₃) δ 7.27–7.22 (1H, m, ArH), 7.00–6.96 (1H, m, ArH), 6.94 (1H, s, ArH), 6.80–6.75 (1H, m, ArH), 6.71 (2H, d, *J* = 8.8 Hz, ArH), 6.49 (2H, d, *J* = 8.8 Hz, ArH), 4.39 (1H, q, *J* = 6.7 Hz, NCH), 3.80 (3H, s, CCHCOCH₃), 3.71 (3H, s, CHCHCOCH₃), 1.50 (3H, d, *J* = 6.7 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 159.9 (C), 151.8 (C), 147.4 (C), 141.5 (C), 129.6 (CH), 118.2 (CH), 114.7 (2 × CH), 114.5 (2 × CH), 111.9 (CH), 111.6 (CH), 55.7 (CH₃), 55.1 (CH₃), 54.3 (CH), 25.1 (CH₃).

(±)-4-Methoxy-N-(1-o-toly/lethyl)aniline (9j). The title compound was prepared according to *General Procedure 2* using boronic ester 8j (0.120 g, 0.487 mmol) and *p*-anisidine (0.251 g, 2.04 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/petroleum ether) of the crude material gave amine 9j (0.0809 g, 69%) as a brown oil. The data were consistent with the literature.^{50 1}H NMR (400 MHz, CDCl₃) δ 7.46–7.40 (1H, m, ArH), 7.20–7.10 (3H, m, ArH), 6.70 (2H, d, J =

8.9 Hz, ArH), 6.42 (2H, d, J = 8.9 Hz, ArH), 4.62 (1H, q, J = 6.6 Hz, NCH), 3.77 (1H, s, NH), 3.70 (3H, s, OCH₃), 2.44 (3H, s, CCH₃), 1.47 (3H, d, J = 6.6 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 151.8 (C), 143.0 (C), 141.6 (C), 134.6 (C), 130.6 (CH), 126.6 (CH), 126.5 (CH), 124.6 (CH), 114.8 (2 × CH), 114.1 (2 × CH), 55.7 (CH₃), 50.4 (CH), 23.1 (CH₃), 19.0 (CH₃).

 (\pm) -N-(4-Methoxyphenyl)- α -methyl-1-naphthalenemethanamine (9k). The title compound was prepared according to General Procedure 2 using boronic ester 8k (0.141 g, 0.499 mmol) and panisidine (0.249 g, 2.02 mmol), heating at 65 °C for 16 h. Flash chromatography (6% EtOAc/petroleum ether) of the crude material gave amine 9k (0.0623 g, 45%) as a yellow solid. The data were consistent with the literature.⁵¹ ¹H NMR (400 MHz, CDCl₃) δ 8.20– 8.18 (1H, m, ArH), 7.93-7.91 (1H, m, ArH), 7.77-7.75 (1H, m, ArH), 7.68-7.66 (1H, m, ArH), 7.60-7.51 (2H, m, ArH), 7.44-7.41 (1H, m, ArH), 6.67 (2H, d, J = 8.8 Hz, ArH), 6.45 (2H, d, J = 8.8 Hz ArH), 5.23 $(1H, q, I = 6.6 \text{ Hz}, \text{NCH}), 3.96 (1H, \text{ br s}, \text{NH}), 3.69 (3H, s, \text{OCH}_2),$ 1.66 (3H, d, J = 6.6 Hz, CHCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 151.8 (C), 141.3 (C), 140.1 (C), 134.1 (C), 130.7 (C), 129.1 (CH), 127.3 (CH), 126.0 (CH), 125.9 (CH), 125.4 (CH), 122.6 (CH), 122.3 (CH), 114.7 (2 × CH), 114.2 (2 × CH), 55.7 (CH₃), 50.1 (CH), 23.8 (CH₃).

(±)-tert-Butyl 3-(3-Methoxy-1-[(4-methoxyphenyl)amino]propyl)-1H-indole-1-carboxylate (91). The title compound was prepared according to General Procedure 2 using boronic ester 81 (0.208 g, 0.500 mmol) and *p*-anisidine (0.249 g, 2.02 mmol), heating at 65 °C for 48 h. Flash chromatography (10% EtOAc/petroleum ether) of the crude material gave amine 91 (0.151 g, 73%) as a dark yellow oil. IR 2932, 2832, 2248, 1726, 1510, 906 cm⁻¹. ¹H NMR (400 MHz, $CDCl_3$) $\delta 8.15$ (1H, s, ArH), 7.65 (1H, d, J = 7.7 Hz, ArH), 7.53 (1H, s, ArH), 7.36-7.32 (1H, m, ArH), 7.26-7.24 (1H, m, ArH), 6.73 (2H, d, *J* = 8.9 Hz, ArH), 6.60 (2H, d, *J* = 8.9 Hz, ArH), 4.77 (1H, t, *J* = 6.3 Hz, NCH), 3.72 (3H, s, ArOCH₃), 3.60-3.55 (1H, m, CH_AH_BO), 3.51-3.46 (1H, m, CH_AH_BO), 3.37 (3H, s, CH₂OCH₃), 2.26-2.18 (2H, m, CHCH₂), 1.67 (9H, s, 3 × CCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.0 (C), 149.7 (C), 141.7 (C), 135.8 (C), 129.0 (C), 124.4 (2 \times CH), 123.0 (CH), 122.4 (2 × CH), 119.3 (CH), 115.4 (CH), 114.7 (2 × CH), 83.6 (C), 70.3 (CH₂), 58.8 (CH₃), 55.7 (CH₃), 50.4 (CH), 36.3 (CH₂), 28.2 (3 × CH₃). HRMS (Q-TOF) m/z: [M + Na]⁺ Calcd for C₂₄H₃₀N₂O₄⁺ 433.2106; found 433.2098.

(±)-N-[3-(Benzyloxy)-1-phenylpropyl]-4-methoxyaniline (9m). The title compound was prepared according to General Procedure 2 using boronic ester 8m (0.177 g, 0.501 mmol) and *p*-anisidine (0.247 g, 2.02 mmol), heating at 65 °C for 32 h. Flash chromatography (8% EtOAc/petroleum ether) of the crude material gave amine 9m (0.103 g, 59%) as an orange oil. IR 3385, 3028, 2857, 1510, 1235 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.38-7.28 (9H, m, ArH), 7.25-7.19 (1H m, ArH), 6.67 (2H, d, J = 8.9 Hz, ArH), 6.42 (2H, d, J = 8.9 Hz, ArH), 4.61 (1H, s, NH), 4.51 (2H, s, ArCH₂), 4.49-4.45 (1H, m, NCH), 3.69 (3H, s, OCH₃), 3.63-3.52 (2H, m, CH₂CH₂O), 2.12-2.05 (2H, m, CHCH₂). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 151.6 (C), 144.0 (C), 141.9 (C), 138.1 (C), 128.5 (2 × CH), 128.4 (2 × CH), 127.7 (2 × CH), 127.7 (CH), 126.8 (CH), 126.4 (2 × CH), 114.6 (2 × CH), 114.3 (2 × CH), 73.2 (CH₂), 68.1 (CH₂), 57.8 (CH₃), 55.7 (CH), 38.3 (CH₂). HRMS (Q-TOF) m/z: [M + H]⁺ Calcd for C₂₃H₂₆NO₂⁺ 348.1964; found 348.1958.

(±)-*N*-(4-Azido-1-phenylbutyl)-4-methoxyaniline (9n). The title compound was prepared according to *General Procedure* 2 using boronic ester 8n (0.151 g, 0.500 mmol)and *p*-anisidine (0.247 g, 2.01 mmol), heating at 65 °C for 16 h. Flash chromatography (5% EtOAc/ petroleum ether) of the crude material gave *amine* 9n (0.0869 g, 59%) as a brown oil. IR 3396, 2930, 2093, 1509, 1451, 1233 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.36–7.29 (4H, m, ArH), 7.26–7.21 (1H, m, ArH), 6.70 (2H, d, *J* = 8.9 Hz, ArH), 6.49 (2H, d, *J* = 8.9 Hz, ArH), 4.32–4.23 (1H, m, NCH), 3.70 (3H, s, OCH₃), 3.34–3.25 (2H, m, CH₂), 1.96–1.81 (2H, m, CH₂), 1.77–1.69 (1H, m, CH₂), 1.66–1.59 (1H, m, CH₂). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.0 (C), 143.6 (C), 141.3 (C), 128.7 (2 × CH), 127.1 (CH), 126.4 (2 × CH), 114.8 (2 × CH), 114.6 (2 × CH), 58.6 (CH₃), 55.7 (CH), 51.2 (CH₂), 35.7

(CH₂), 25.8 (CH₂). **HRMS** (Q-TOF) m/z: $[M + H]^+$ Calcd for $C_{17}H_{21}N_4O^+$ 297.1715; found 297.1717.

(±)-(1-{4-[(4-Methoxyphenyl)amino]-4-phenylbutyl}-1H-1,2,3triazol-4-yl)methanol (90). The title compound was prepared according to a modification of General Procedure 2 using boronic ester 80 (0.0899 g, 0.252 mmol), p-anisidine (0.124 g, 1.00 mmol), Cu(OAc)₂ (0.0910 g, 0.501 mmol), Cs₂CO₃ (0.0410 g, 0.126 mmol), methanol (0.5 mL), and pyridine (0.17 mL). The mixture was stirred at 65 °C for 48 h. Flash chromatography (2%-5% EtOAc/petroleum ether) of the crude material gave amine 90 (0.0457 g, 52%) as an off white solid. m.p. 135-136 °C (CHCl₃). IR 3333, 3207, 2928, 2858, 1737, 1513, 1234 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.45 (1H, s, ArH), 7.35-7.28 (4H, m, ArH), 7.26-7.20 (1H, m, ArH), 6.68 (2H, d, *J* = 8.8 Hz, ArH), 6.48 (2H, d, *J* = 8.8 Hz, ArH), 4.78 (2H, s, CH₂OH), 4.38-4.32 (2H, m, CH₂N), 4.29-4.24 (1H, m, CHN), 3.69 (3H, s, OCH₃), 2.10–2.02 (1H, m, CHCH_ACH_B), 1.96–1.89 (1H, m, $CHCH_{A}CH_{B}$), 1.85–1.75 (2H, m, $CH_{2}CH_{2}N$). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.4 (C), 147.7 (C), 142.9 (C), 140.6 (C), 128.7 (2 × CH), 127.3 (CH), 126.4 (2 × CH), 121.5 (CH), 115.1 (2 × CH), 114.8 (2 × CH), 58.8 (CH), 56.6 (CH₂), 55.7 (CH₃), 50.0 (CH₂), 35.0 (CH₂), 27.1 (CH₂). HRMS (Q-TOF) m/z: [M + H]⁺ Calcd for C20H25N4O2+ 353.1978; found 353.1972.

4-Methoxy-N-(naphthalen-2-ylmethyl)aniline (**9p**). The title compound was prepared according to *General Procedure* 2 using boronic ester **8p** (0.135 g, 0.505 mmol) and *p*-anisidine (0.249 g, 2.02 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine **9p** (0.0825 g, 62%) as a yellow solid. The data were consistent with the literature.⁵² m.p. 106–108 °C (petroleum ether); no literature value available. ¹H NMR (400 MHz, CDCl₃) δ 7.88–7.78 (4H, m, ArH), 7.54–7.42 (3H, m, ArH), 6.79 (2H, d, *J* = 8.9 Hz, ArH), 6.66 (2H, d, *J* = 8.9 Hz, ArH), 4.46 (2H, s, CH₂), 3.75 (3H, s, OCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.2 (C), 142.4 (C), 137.2 (C), 133.5 (C), 132.7 (C), 128.3 (CH), 127.7 (CH), 126.1 (CH), 125.9 (CH), 125.8 (CH₃), 19.4 (CH₂).

(±)-*N*-(4-Methoxybenzyl)cyclohex-2-enamine (9q). The title compound was prepared according to *General Procedure* 2 using boronic ester 8q (0.104 g, 0.501 mmol) and *p*-anisidine (0.247 g, 2.00 mmol), heating at 65 °C for 48 h. Flash chromatography (3% EtOAc/ petroleum ether) of the crude material gave amine 9q (0.0498 g, 49%) as a yellow oil. The data were consistent with the literature.⁵³ ¹H NMR (400 MHz, CDCl₃) δ 6.79 (2H, d, *J* = 9.0 Hz, ArH), 6.61 (2H, d, *J* = 9.0 Hz, ArH), 5.88–5.79 (1H, m, CHCH=CH), 5.80–5.72 (1H, m, CHCH=CH), 3.92 (1H, s, NCH), 3.76 (3H, s, OCH₃), 3.34 (1H, s, NH), 2.13–1.99 (2H, m, CH=CHCH₂), 1.95–1.85 (1H, m, NCHCH₄H_B), 1.81–1.69 (1H, m, NCHCH₂CH₄H_B), 1.67–1.52 (2H, m, NCHCH₂CH₄H_B + NCHCH₄H_B). ¹³C¹H NMR (101 MHz, CDCl₃) δ 152.0 (C), 141.3 (C), 129.9 (CH), 128.9 (CH), 115.0 (2 × CH), 114.9 (2 × CH), 55.8 (CH₃), 49.0 (CH), 29.0 (CH₂), 25.2 (CH₂), 19.8 (CH₂).

N-Methyl-N-(3-methylbut-2-en-1-yl)aniline (*9r*). The title compound was prepared according to *General Procedure 2* using boronic ester **8r** (0.093 g, 0.509 mmol) and *N*-methyl aniline (0.216 g, 2.01 mmol), heating at 50 °C for 16 h. Flash chromatography (6% EtOAc/ petroleum ether) of the crude material gave amine **9r** (0.0464 g, 57%) as a yellow oil. The data were consistent with the literature.^{54 1}H NMR (400 MHz, CDCl₃) δ 7.25–7.21 (2H, m, ArH), 6.71–6.68 (3H, m, ArH), 4.86 (1H, s, C=CH_AH_B), 4.81 (1H, s, C=CH_AH_B) 3.81 (2H, s, NCH₂), 2.97 (3H, s, NCH₃), 1.74 (3H, s, CCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 149.6 (C), 141.5 (C), 129.0 (2 × CH), 116.1 (CH), 111.9 (2 × CH), 110.7 (CH₂), 58.8 (CH₂), 38.2 (CH₃), 2.00 (CH₃).

N-Cyclohexyl-4-methoxyaniline (9s). Methanol (1.0 mL) and pyridine (0.33 mL) were added to a flask containing cyclohexylboronic acid pinacol ester (0.106 g, 0.503 mmol), *p*-anisidine (0.248 g, 2.03 mmol), Cu(OAc)₂ (0.182 g, 1.00 mmol), and Cs₂CO₃ (0.0820 g, 0.252 mmol), and the mixture was stirred at 80 °C for 16 h. The mixture was cooled to room temperature, NH₄OH (10 mL) was added, and the mixture was extracted with Et₂O (3×10 mL). The combined organic phases were dried (MgSO₄), filtered, and concentrated *in vacuo*. Flash chromatography (5% EtOAc/petroleum ether) of the crude material

gave amine **9s** (0.0114 g, 11%) as an orange oil. The data were consistent with the literature.⁵⁵ ¹H NMR (400 MHz, CDCl₃) δ 6.77 (2H, d, *J* = 8.2 Hz, ArH), 6.58 (2H, d, *J* = 8.2 Hz, ArH), 3.75 (3H, s, OCH₃), 3.21–3.13 (1H, m, NCH), 2.08–2.02 (2H, m, CHCH₄H_B), 1.80–1.69 (2H, m, CHCH₂CH₄H_B), 1.69–1.63 (1H, m, CHCH₂CH₂CH₄H_B), 1.40–1.31 (2H, m, CHCH₂CH₂CH₄H_B), 1.27 (1H, s, NH), 1.25–1.18 (1H, m, CHCH₂CH₂CH₄H_B), 1.18–1.08 (2H, m, CHCH₄H_B). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 151.9 (C), 141.6 (C), 114.9 (2 × CH), 114.8 (2 × CH), 55.8 (CH₃), 52.8 (CH), 33.6 (2 × CH₂), 26.0 (CH₂), 25.1 (2 × CH₂).

(±)-4-Methoxy-N-[1-(4-methoxyphenyl)-1-phenylethyl]aniline (11a). The title compound was prepared according to General Procedure 4 using boronic ester 10a (0.168 g, 0.498 mmol) and p-anisidine (0.246 g, 2.00 mmol). Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave amine 11a (0.0977 g, 59%) as a yellow oil. IR 3396, 2991, 2932, 2833, 1608, 1507, 1237 cm⁻¹. ¹H NMR (400 MHz, $CDCl_3$) δ 7.49–7.43 (2H, m, ArH), 7.37 (2H, d, J = 8.8 Hz, ArH), 7.35-7.29 (2H, m, ArH), 7.26-7.20 (1H, m, ArH), 6.85 (2H, d, J = 8.8 Hz, ArH), 6.64 (2H, d, J = 8.9 Hz, ArH), 6.41 (2H, d, J = 8.9 Hz, ArHH), 4.03 (1H, s, NH), 3.80 (3H, s, OCH₃), 3.70 (3H, s, OCH₃), 2.03 (3H, s, CCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 158.2 (C), 152.4 (C), 147.7 (C), 139.8 (C), 139.7 (C), 128.3 (2 × CH), 128.0 (2 × CH), 126.8 (2 × CH), 126.6 (CH), 118.2 (2 × CH), 114.2 (2 × CH), 113.6 (2 × CH), 62.2 (C), 55.6 (CH₃), 55.2 (CH₃), 26.7 (CH₃). HRMS (Q-TOF) *m/z*: [M]⁺ Calcd for C₂₂H₂₃NO₂⁺ 333.1723; found 333.1719.

(±)-4-Chloro-N-[1-(4-methoxyphenyl)-1-phenylethyl]aniline (11b). The title compound was prepared according to General Procedure 4 using boronic ester 10a (0.170 g, 0.501 mmol) and 4-chloroaniline (0.257 g, 2.01 mmol). Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave *amine* 11b (0.0710 g, 42%) as a yellow oil. IR 3473, 3383, 3006, 1601, 1493, 1247 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.46–7.40 (2H, m, J = 8.9 Hz, ArH), 7.37–7.31 (4H, m, ArH), 7.27–7.23 (1H, m, ArH), 6.97 (2H, d, J = 8.9 Hz, ArH), 6.86 (2H, d, J = 8.9 Hz, ArH), 6.35 (2H, d, J = 8.9 Hz, ArH), 4.39 (1H, s, NH), 3.81 (3H, s, OCH₃), 2.07 (3H, s, CCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 158.4 (C), 146.8 (C), 144.5 (C), 138.7 (C), 128.5 (2 × CH), 128.5 (2 × CH), 128.0 (2 × CH), 126.9 (CH), 126.7 (2 × CH), 122.2 (C), 117.0 (2 × CH), 113.7 (2 × CH), 62.0 (C), 55.2 (CH₃), 26.9 (CH₃). HRMS (Q-TOF) *m/z*: [M + Na]⁺ Calcd for C₂₁H₂₀³⁵ClNNaO⁺ 360.1131; found 360.1135.

(±)-*N*-[1-(4-Methoxyphenyl)-1-phenylethyl]-1-methyl-1H-pyrazol-5-amine (11c). The title compound was prepared according to General Procedure 4 using boronic ester 10a (0.169 g, 0.501 mmol) and 1-methyl-1H-pyrazol-3-amine (0.194 g, 2.00 mmol). Flash chromatography (1:5 EtOAc/petroleum ether) of the crude material gave amine 11c (0.0522 g, 34%) as a yellow oil. IR 3267, 2933, 2835, 1547, 1509, 1249 cm^{-1.} ¹H NMR (400 MHz, CDCl₃) δ 7.50–7.45 (2H, m, ArH), 7.39 (2H, d, *J* = 8.9 Hz, ArH), 7.33–7.28 (2H, m, ArH), 7.25–7.19 (1H, m, ArH), 6.92 (1H, d, *J* = 2.3 Hz, ArH), 6.84 (2H, d, *J* = 8.9 Hz, ArH), 4.84 (1H, d, *J* = 2.3 Hz, ArH), 4.58 (1H, s, NH), 3.79 (3H, s, OCH₃), 3.68 (3H, s, NCH₃), 2.00 (3H, s, CCH₃). ¹³C{¹H} NMR (126 MHz, CDCl₃) δ 158.1 (C), 155.0 (C), 147.7 (C), 139.6 (C), 130.3 (CH), 128.2 (2 × CH), 128.0 (2 × CH), 126.7 (2 × CH), 126.5 (CH), 113.4 (2 × CH), 94.0 (CH), 61.8 (C), 55.2 (CH₃), 38.3 (CH₃), 27.7 (CH₃). HRMS (Q-TOF) *m*/*z*: [M + H]⁺ Calcd for C₁₉H₂₂N₃O⁺ 308.1763; found 308.1757.

(±)-*N*-[1-(4-Methoxyphenyl)-1-phenylethyl]aniline (11d). The title compound was prepared according to *General Procedure* 4 using boronic ester 10a (0.169 g, 0.499 mmol) and aniline (0.18 mL, 2.0 mmol). Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave *amine* 11d (0.0642 g, 42%) as a yellow oil. IR 3409, 2989,2834, 1599, 1497, 1249 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.46 (2H, d, *J* = 8.8 Hz, ArH), 7.38–7.30 (4H, m, ArH), 7.26–7.21 (1H, m, ArH), 7.07–7.00 (2H, m, ArH), 6.86 (2H, d, *J* = 8.8 Hz, ArH), 6.68–6.63 (1H, m, ArH), 6.47–6.41 (2H, m, ArH), 4.36 (1H, s, NH), 3.80 (3H, s, OCH₃), 2.10 (3H, s, CCH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 158.3 (C), 147.3 (C), 146.0 (C), 139.3 (C), 128.6 (2 × CH), 128.4 (2 × CH), 128.0 (2 × CH), 126.8 (2 × CH), 126.7 (CH), 117.4 (CH), 116.0 (2 × CH), 113.6 (2 × CH), 61.9 (C), 55.2 (CH₃), 26.6

(CH₃). **HRMS** (Q-TOF) m/z: [M + Na]⁺ Calcd for C₂₁H₂₁NNaO⁺ 326.1521; found 326.1536.

(±)-*N*-[1-(4-Chlorophenyl)-1-phenylethyl]aniline (11e). The title compound was prepared according to *General Procedure 4* using boronic ester 10b (0.171 g, 0.499 mmol) and aniline (0.18 mL, 2.0 mmol), heating at 25 °C. Flash chromatography (2% EtOAc/ petroleum ether) of the crude material gave *amine* 11e (0.0681 g, 44%) as a yellow oil. IR 3150, 1492, 1097, 1012, 750 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.46–7.41 (4H, m, ArH), 7.37–7.28 (5H, m, ArH), 7.08–7.03 (2H, m, ArH), 6.73–6.66 (1H, m, ArH), 6.47–6.42 (2H, m, ArH), 4.34 (1H, s, NH), 2.11 (3H, s, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 146.8 (C), 145.6 (C), 145.4 (C), 132.6 (C), 128.7 (2 × CH), 128.5 (6 × CH), 127.1 (CH), 126.6 (2 × CH), 117.8 (CH), 116.1 (2 × CH), 62.1 (C), 26.7 (CH₃). HRMS (ESI) *m/z*: [M + H]⁺ Calcd for C₂₀H₁₈³⁵ClN⁺ 308.1201; found 308.1194.

(±)-*N*-{1-*Phenyl*-1-[4-(trifluoromethyl)phenyl]ethyl}aniline (11f). The title compound was prepared according to *General Procedure* 4 using boronic ester 10c (0.189 g, 0.502 mmol) and aniline (0.18 mL, 2.0 mmol), heating at 50 °C. Flash chromatography (2% EtOAc/petroleum ether) of the crude material gave *amine* 11f (0.0432 g, 25%) as a yellow oil. IR 3450, 2928, 1600, 1497, 1325 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.56–7.48 (4H, m, ArH), 7.34 (2H, d, *J* = 7.7 Hz, ArH), 7.28–7.23 (2H, m, ArH), 7.20–7.16 (1H, m, ArH), 7.00–6.93 (2H, m, ArH), 6.64–6.57 (1H, m, ArH), 6.34 (2H, d, *J* = 7.7 Hz, ArH), 4.27 (1H, s, NH), 2.05 (3H, s, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 150.7 (C), 146.5 (C), 145.5 (C), 129.0 (C, q, *J_F* = 32.6 Hz), 128.7 (2 × CH), 128.6 (2 × CH), 127.4 (2 × CH), 127.2 (CH), 126.6 (2 × CH), 125.4 (2 × CH, q, *J_F* = 3.7 Hz), 124.2 (CF₃, q, *J_F* = 271.8 Hz), 118.0 (CH), 116.2 (2 × CH), 62.4 (C), 26.6 (CH₃). ¹⁹F NMR (377 MHz, CDCl₃) δ –62.3. HRMS (EI) *m*/*z*: [M]⁺ Calcd for C₂₁H₁₈F₃N⁺ 341.1386; found 341.1399.

(±)-N-(2-Phenylbutan-2-yl)aniline (11g). The title compound was prepared according to *General Procedure* 4 using boronic ester 10d (0.130 g, 0.500 mmol) and aniline (0.18 mL, 2.0 mmol), heating at 50 °C. Flash chromatography (1% EtOAc/petroleum ether) of the crude material gave amine 11g (0.0466 g, 41%) as a yellow oil. The data were consistent with the literature.⁵⁶ IR 3359, 2918, 2850, 1600, 1468, 1263. ¹H NMR (400 MHz, CDCl₃) δ 7.52–7.45 (2H, m, ArH), 7.36–7.30 (2H, m, ArH), 7.26–7.21 (1H, m, ArH), 7.04–6.95 (2H, m, ArH), 6.65–6.56 (1H, m, ArH), 6.38–6.30 (2H, m, ArH), 4.01 (1H, s, NH), 1.92 (2H, q, *J* = 7.4, CH₂), 1.63 (3H, s, CCH₃), 0.83 (3H, t, *J* = 7.4 Hz, CH₂CH₃). ¹³C{¹H} NMR (400 MHz, CDCl3) δ 146.3 (C), 146.0 (C), 128.7 (2 × CH), 128.3 (2 × CH), 126.2 (3 × CH), 116.9 (CH), 115.1 (2 × CH), 58.4 (C), 36.4 (CH₂), 25.3 (CH₃), 8.2 (CH₃). HRMS (ESI) *m/z*: [M + H]⁺ Calcd for C₁₆H₁₉N⁺ 226.1590; found 226.1597.

Reaction of Boronic Ester 13 with *p***-Anisidine.** (See Scheme S1 in the Supporting Information.)

According to *General Procedure 4*, boronic ester **13** (0.159 g, 0.503 mmol) and *p*-anisidine (0.246 g, 2.00 mmol) were heated at 25 °C. Flash chromatography (5% EtOAc/petroleum ether) of the crude material gave *amine* **14** (0.0365 g, 23%) as a yellow oil, and an inseparable mixture of *alcohol* **S2** and *amine* **15** (40.9 mg).

Phenyl isothiocyanate (1.1 equiv) was added to a solution of a mixture of S2 and 15 in hexane:Et₃N:MeCN (1:0.1:0.1, 0.1 M), and the mixture was stirred for 16 h at room temperature. The precipitate was collected and washed with hexane (2×5 mL) to give *thiourea* 16 (16.0 mg, 7%). Flash chromatography (petroleum ether \rightarrow 20% EtOAc/ petroleum ether) of the mother liquor gave *alcohol* S2 (18.1 mg, 18%).

(±)-N-(5-Azido-2-phenylpentan-2-yl)-4-methoxyaniline (14). IR 3390, 2932, 2831, 2093, 1508, 1235, cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.53–7.45 (2H, m, ArH), 7.39–7.32 (2H, m, ArH), 7.27– 7.23 (1H, m, ArH), 6.62 (2H, d, *J* = 8.8 Hz, ArH), 6.31 (2H, d, *J* = 8.8 Hz, ArH), 3.69 (4H, s, NH, OCH₃), 3.26–3.12 (2H, m, CH₂N), 2.01– 1.88 (2H, m, CH₂CH₂N), 1.61 (3H, s, CCH₃), 1.55–1.47 (2H, m, CCH₂). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.0 (C), 146.2 (C), 139.7 (C), 128.5 (2 × CH), 126.5 (CH), 126.2 (2 × CH), 117.0 (2 × CH), 114.3 (2 × CH), 58.3 (C), 55.6 (CH₃), 51.6 (CH₂), 40.6 (CH₂), 26.3 (CH₃), 23.5 (CH₂). HRMS (Q-TOF) *m*/*z*: [M + H]⁺ Calcd for C₁₈H₂₃N₄O⁺ 311.1872; found 311.1866.

(±)-1-[2-(5-Azido-2-phenylpentan-2-yl)-4-methoxyphenyl]-3-phenylthiourea (**16**). m.p. 150–151 °C (hexane). IR 3334, 3170, 2959, 2096, 1508, 1214 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.57 (1H, s, NH), 7.35–7.29 (1H, m, ArH), 7.26–7.21 (2H, m, ArH), 7.18–7.13 (2H, m, ArH), 7.04 (6H, m, ArH), 6.90–6.87 (1H, m, ArH), 6.38 (1H, s, ArH), 3.90 (3H, s, OCH₃), 3.38–3.28 (1H, m, CH₄H_BN), 3.28–3.18 (1H, m, CH₄H_BN), 2.23–2.03 (2H, m, CH₂CH₂N₃), 1.57 (3H, s, CCH₃), 1.44–1.31 (2H, m, CCH₂). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 179.7 (C), 158.5 (C), 148.2 (C), 145.1 (C), 136.8 (C), 131.9 (CH), 129.6 (CH), 128.8 (4×CH), 126.6 (2×CH), 126.3 (2×CH), 125.4 (CH), 123.8 (C), 115.4 (CH), 110.8 (CH), 55.5 (CH₃), 51.9 (CH₂), 45.5 (C), 36.1 (CH₂), 27.8 (CH₃), 24.8 (CH₂). HRMS (Q-TOF) *m/z*: [M + H]⁺ Calcd for C₂₃H₂₈N₅OS⁺ 446.2015; found 446.2009.

(±)-5-Azido-2-phenylpentan-2-ol (**S2**). IR 3420, 2930, 2092, 1602, 1146, 1260 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.48–7.39 (2H, m, ArH), 7.40–7.33 (2H, m, ArH), 7.29–7.24 (1H, m, ArH), 3.26–3.19 (2H, m, CH₂N₃), 1.95–1.83 (2H, m, CH₂CH₂N₃), 1.80 (1H, s, OH), 1.64–1.57 (4H, m, CH₃, CCH₄H_B), 1.53–1.38 (1H, m, CCH₄H_B). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 147.2 (C), 128.3 (2 × CH), 126.7 (CH), 124.6 (2 × CH), 74.3 (C), 51.6 (CH₂), 41.1 (CH₂), 30.5 (CH₃), 23.7 (CH₂). HRMS (Q-TOF) *m*/*z*: [M + Na]⁺ Calcd for C₁₁H₁₅N₃ONa⁺ 228.1113; found 228.1107.

Reaction of Boronic Ester 1 with *p***-Aminophenol.** The title compounds were prepared according to *General Procedure 2* using boronic ester 1 (0.116 g, 0.500 mmol) and 4-aminophenol (0.218 g, 2.00 mmol), heating at 65 °C for 48 h. Flash chromatography (20% EtOAc/petroleum ether) of the crude material gave amine 17 (0.0499 g, 47%), *amine* **18** (0.0023 g, 2%), and *amine* **19** (0.0338 g, 32%) as red oils.

(±)-4-[(1-Phenylethyl)amino]phenol (17). The data were consistent with the literature.⁵⁷ ¹H NMR (400 MHz, CDCl₃) δ 7.38–7.30 (4H, m, ArH), 7.25–7.21 (1H, m, ArH), 6.62 (2H, d, *J* = 8.8 Hz, ArH), 6.43 (2H, d, *J* = 8.8 Hz, ArH), 4.41 (1H, q, *J* = 6.7 Hz, CH), 2.19 (1H, s, NH), 1.50 (3H, d, *J* = 6.7 Hz, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 147.4 (C), 145.4 (C), 141.6 (C), 128.6 (2 × CH), 126.8 (CH), 125.9 (2 × CH), 116.0 (2 × CH), 114.7 (2 × CH), 54.3 (CH), 25.1 (CH₃).

(±)-4-(1-Phenylethoxy)aniline (18). IR 3446, 3360, 2920, 2850, 1624, 1508, 1230 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.39–7.30 (4H, m, ArH), 7.27–7.21 (1H, m, ArH), 6.70 (2H, d, *J* = 8.7 Hz, ArH), 6.56 (2H, d, *J* = 8.7 Hz, ArH), 5.17 (1H, q, *J* = 6.5 Hz, CH), 3.44 (2H, s, NH₂), 1.60 (3H, d, *J* = 6.5 Hz, CH₃). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 151.1 (C), 143.6 (C), 139.9 (C), 128.5 (2 × CH), 127.3 (CH), 125.7 (2 × CH), 117.4 (2 × CH), 116.3 (2 × CH), 76.9 (CH), 24.3 (CH₃). HRMS (Q-TOF) *m*/*z*: [M + H]⁺ Calcd for C₁₄H₁₆NO⁺ 214.1232; found 214.1226.

(±)-4-Åmino-3-(1-phenylethyl)phenol (**19**). **IR** 3363, 2966, 2929, 1600, 1498, 1450, 1214 cm⁻¹. ¹**H NMR** (400 MHz, CDCl₃) δ 7.32–7.27 (2H, m, ArH), 7.23–7.18 (3H, m, ArH), 6.82 (1H, d, *J* = 2.7 Hz, ArH), 6.60 (1H, dd, *J* = 8.4, 2.7 Hz, ArH), 6.55 (1H, d, *J* = 8.4 Hz, ArH), 4.09 (1H, q, *J* = 7.2 Hz, CH), 1.59 (3H, d, *J* = 7.2 Hz, CH₃). ¹³C{¹H} **NMR** (101 MHz, CDCl₃) δ 148.8 (C), 145.3 (C), 137.5 (C), 132.1 (C), 128.7 (2 × CH), 127.5 (2 × CH), 126.4 (CH), 117.6 (CH), 114.6 (CH), 113.8 (CH), 40.1 (CH), 21.7 (CH₃). **HRMS** (Q-TOF) *m/z*: [M + H]⁺ Calcd for C₁₄H₁₆NO⁺ 214.1232; found 214.1226.

(±)-3-[4-Hydroxy-2-(1-phenylethyl)phenyl]-1-phenylthiourea (**53**). (See Scheme S2 in the Supporting Information.)

Phenyl isothiocyanate (23.8 mg, 0.176 mmol) was added to a solution of amine **19** (33.8 mg, 0.158 mmol) in hexane (1.6 mL), Et₃N (0.16 mL), and MeCN (0.16 mL), and the mixture was stirred for 16 h at room temperature. The precipitate was collected and washed with hexane (2 × 5 mL) to give *thiourea* **S3** (37.2 mg, 68%). **m.p.** 155–157 °C (hexane). **IR** 3339, 3152, 2967, 1747, 1533, 1275 cm⁻¹. ¹H **NMR** (400 MHz, CD₃CN) δ 7.73 (1H, s, NH), 7.33–7.23 (8H, m, ArH), 7.21–7.15 (2H, m, ArH), 7.09–7.06 (1H, m, ArH), 6.79 (1H, s, ArH), 6.73–6.68 (1H m, ArH), 4.33 (1H, q, *J* = 7.2 Hz, CH), 1.55 (3H, d, *J* = 7.2 Hz, CHCH₃). ¹³C{¹H} **NMR** (101 MHz, CD₃CN) δ 182.8 (C), 157.9 (C), 147.0 (C), 146.4 (C), 131.6 (CH), 129.6 (C), 129.4 (4 × CH), 128.5 (4 × CH), 127.1 (CH), 126.9 (C), 126.8 (CH), 115.5

(CH), 114.8 (CH), 40.4 (CH), 21.6 (CH₃). **HRMS** (Q-TOF) m/z: [M + H]⁺ Calcd for C₂₁H₂₁N₂OS⁺ 349.1375; found 349.1369. Crystals suitable for X-ray diffraction were obtained through recrystallization of **S3** from hexane and slow evaporation.

Reaction of Cyclopropane-Containing Boronic Ester 12. A flask containing boronic ester 20 (0.131 g, 0.507 mmol), p-anisidine (0.251 g, 2.04 mmol), Cu(OAc)₂ (0.184 g, 1.01 mmol), and Cs₂CO₃ (0.0858 g, 0.263 mmol) was purged with argon. Methanol (1.0 mL) and pyridine (0.33 mL) were added, and the mixture was stirred at 65 °C for 48 h. The mixture was cooled to room temperature, NH₄OH (10 mL) was added, and the mixture was extracted with $Et_2O(3 \times 10 \text{ mL})$, dried (MgSO₄), filtered, and concentrated in vacuo. Flash chromatography (6% EtOAc/petroleum ether) of the crude material gave amine 21 (0.0501g, 39%) as an orange oil. IR 3390, 2929, 2831, 1510, 1233 cm⁻¹. ¹H NMR (400 MHz, CDCl₃) δ 7.41–7.35 (2H, m, ArH), 7.36–7.29 (2H, m, ArH), 7.27–7.20 (1H, m, ArH), 6.81 (2H, d, J = 8.9 Hz, ArH), 6.62 (2H, d, J = 8.9 Hz, ArH), 6.51 (1H, d, J = 15.9 Hz, PhCH), 6.24 (1H, dt, J = 15.97.1 Hz, PhCH=CH), 3.77 (3H, s, OCH₃), 3.47 (1H, s, NH), 3.25 (2H, t, J = 6.7 Hz, CH₂N), 2.55 (2H, dtd, J = 7.1, 6.7, 1.1 Hz, CH₂CH₂N). ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 152.2 (C), 142.5 (Č), 137.3 (C), 132.2 (CH), 128.5 (2 × CH), 127.5 (CH), 127.2 (CH), 126.1 (2 × CH), 115.0 (2 × CH), 114.3 (2 × CH), 55.8 (CH₃), 44.3 (CH₂), 33.0 (CH₂). **HRMS** (Q-TOF) m/z: [M + H]⁺ Calcd for C₁₇H₂₀NO⁺ 254.1545; found 254.1539.

Reaction of Enantiomerically Enriched Boronic Ester (5)-1. According to *General Procedure 2*, boronic ester (*S*)-1 (0.0502 g, 0.216 mmol) and aniline (0.0803 g, 0.862 mmol) were reacted, heating at 50 °C for 16 h. Flash chromatography (4% EtOAc/petroleum ether) of the crude material gave amine **3a** (0.0311 g, 73%) as a yellow oil. The data were consistent with the literature.³⁵ See above for data. e.r. = 50:50. HPLC (Phenomenex Cellulose-1 column (250 × 4.6 mm), IPA/ hexane 6/94, flow rate = 1.0 mL/min, *l* = 254 nm), *t*_R = 6.5 min (minor), 8.0 min (major).

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.joc.1c00976.

Experimental details and characterization data (PDF)

Accession Codes

CCDC 2073100 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Benjamin M. Partridge – Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom; • orcid.org/ 0000-0002-8550-9994; Email: b.m.partridge@ sheffield.ac.uk

Authors

- James D. Grayson Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom
- Francesca M. Dennis Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom
- **Craig C. Robertson** Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom

Complete contact information is available at: https://pubs.acs.org/10.1021/acs.joc.1c00976

The manuscript was written through contributions of all authors.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the EPSRC (Grant EP/T009292/1), the Royal Society (Grant RSGR1180065), and the University of Sheffield for financial support.

REFERENCES

(1) (a) Heravi, M. M.; Kheilkordi, Z.; Zadsirjan, V.; Heydari, M.; Malmir, M. Buchwald-Hartwig Reaction: An Overview. J. Organomet. Chem. 2018, 861, 17–104. (b) Ruiz-Castillo, P.; Buchwald, S. L. Applications of Palladium-Catalyzed C-N Cross-Coupling Reactions. Chem. Rev. 2016, 116 (19), 12564–12649. (c) Beletskaya, I. P.; Cheprakov, A. V. The Complementary Competitors: Palladium and Copper in C-N Cross-Coupling Reactions. Organometallics 2012, 31 (22), 7753–7808.

(2) (a) Qiao, J. X.; Lam, P. Y. S. Copper-Promoted Carbon-Heteroatom Bond Cross-Coupling with Boronic Acids and Derivatives. *Synthesis* **2011**, *2011* (6), 829–856. (b) Monnier, F.; Taillefer, M. Copper-Catalyzed C(Aryl)-N Bond Formation. In *Topics in Organometallic Chemistry*; Taillefer, M., Ma, D., Eds.; Springer: Berlin, 2013; pp 173–204. (c) Munir, I.; Zahoor, A. F.; Rasool, N.; Naqvi, S. A. R.; Zia, K. M.; Ahmad, R. Synthetic Applications and Methodology Development of Chan-Lam Coupling: A Review. *Mol. Diversity* **2019**, *23* (1), 215–259. (d) West, M. J.; Fyfe, J. W. B.; Vantourout, J. C.; Watson, A. J. B. Mechanistic Development and Recent Applications of the Chan-Lam Amination. *Chem. Rev.* **2019**, *119* (24), 12491–12523.

(3) (a) Sambiagio, C.; Marsden, S. P.; Blacker, A. J.; McGowan, P. C. Copper Catalysed Ullmann Type Chemistry: From Mechanistic Aspects to Modern Development. *Chem. Soc. Rev.* **2014**, *43* (10), 3525–3550. (b) Monnier, F.; Taillefer, M. Catalytic C-C, C-N, and C-O Ullmann-Type Coupling Reactions. *Angew. Chem., Int. Ed.* **2009**, *48* (38), 6954–6971.

(4) Campos, K. R.; Coleman, P. J.; Alvarez, J. C.; Dreher, S. D.; Garbaccio, R. M.; Terrett, N. K.; Tillyer, R. D.; Truppo, M. D.; Parmee, E. R. The Importance of Synthetic Chemistry in the Pharmaceutical Industry. *Science* **2019**, *363* (6424), No. eaat0805.

(5) For stoichiometric reactions, see: (a) Peacock, D. M.; Jiang, Q.; Hanley, P. S.; Cundari, T. R.; Hartwig, J. F. Reductive Elimination from Phosphine-Ligated Alkylpalladium(II) Amido Complexes To Form Sp3 Carbon-Nitrogen Bonds. J. Am. Chem. Soc. 2018, 140 (14), 4893-4904. (b) Hanley, P. S.; Marquard, S. L.; Cundari, T. R.; Hartwig, J. F. Reductive Elimination of Alkylamines from Low-Valent, Alkylpalladium(II) Amido Complexes. J. Am. Chem. Soc. 2012, 134 (37), 15281-15284. (c) Peacock, D. M.; Jiang, Q.; Cundari, T. R.; Hartwig, J. F. Reductive Elimination to Form C(Sp3)-N Bonds from Palladium(II) Primary Alkyl Complexes. Organometallics 2018, 37 (19), 3243-3247. For catalytic reactions, see: (d) Pan, J.; Su, M.; Buchwald, S. L. Palladium(0)-Catalyzed Intermolecular Amination of Unactivated C Sp3-H Bonds. Angew. Chem., Int. Ed. 2011, 50 (37), 8647-8651. (e) Peacock, D. M.; Roos, C. B.; Hartwig, J. F. Palladium-Catalyzed Cross Coupling of Secondary and Tertiary Alkyl Bromides with a Nitrogen Nucleophile. ACS Cent. Sci. 2016, 2 (9), 647-652.

(6) (a) Bissember, A. C.; Lundgren, R. J.; Creutz, S. E.; Peters, J. C.; Fu, G. C. Transition-Metal-Catalyzed Alkylations of Amines with Alkyl Halides: Photoinduced, Copper-Catalyzed Couplings of Carbazoles. *Angew. Chem., Int. Ed.* **2013**, *52* (19), 5129–5133. (b) Ahn, J. M.; Peters, J. C.; Fu, G. C. Design of a Photoredox Catalyst That Enables the Direct Synthesis of Carbamate-Protected Primary Amines via Photoinduced, Copper-Catalyzed N-Alkylation Reactions of Unactivated Secondary Halides. *J. Am. Chem. Soc.* **2017**, *139* (49), 18101– 18106. (c) Matier, C. D.; Schwaben, J.; Peters, J. C.; Fu, G. C. Copper-Catalyzed Alkylation of Aliphatic Amines Induced by Visible Light. *J.* Am. Chem. Soc. 2017, 139 (49), 17707–17710. (d) Kainz, Q. M.; Matier, C. D.; Bartoszewicz, A.; Zultanski, S. L.; Peters, J. C.; Fu, G. C. Asymmetric Copper-Catalyzed C-N Cross-Couplings Induced by Visible Light. Science 2016, 351 (6274), 681–684. (e) Ahn, J. M.; Ratani, T. S.; Hannoun, K. I.; Fu, G. C.; Peters, J. C. Photoinduced, Copper-Catalyzed Alkylation of Amines: A Mechanistic Study of the Cross-Coupling of Carbazole with Alkyl Bromides. J. Am. Chem. Soc. 2017, 139 (36), 12716–12723.

(7) Liang, Y.; Zhang, X.; MacMillan, D. W. C. Decarboxylative Sp3 C-N Coupling via Dual Copper and Photoredox Catalysis. *Nature* **2018**, 559 (7712), 83–88.

(8) (a) Tsuritani, T.; Strotman, N. A.; Yamamoto, Y.; Kawasaki, M.; Yasuda, N.; Mase, T. N-Cyclopropylation of Indoles and Cyclic Amides with Copper(II) Reagent. Org. Lett. 2008, 10 (8), 1653-1655. (b) Bénard, S.; Neuville, L.; Zhu, J. Copper-Mediated N-Cyclopropylation of Azoles, Amides, and Sulfonamides by Cyclopropylboronic Acid. J. Org. Chem. 2008, 73 (16), 6441-6444. (c) Bénard, S.; Neuville, L.; Zhu, J. Copper-Promoted N-Cyclopropylation of Anilines and Amines by Cyclopropylboronic Acid. Chem. Commun. 2010, 46 (19), 3393-3395. (d) Tambe, Y. B.; Sharma, S.; Pathak, A.; Reddy, L. K. Microwave-Assisted N-Cyclopropylation of Pyridinols with Cyclopropyl Boronic Acid. Synth. Commun. 2012, 42 (9), 1341-1348. (e) Mudryk, B.; Zheng, B.; Chen, K.; Eastgate, M. D. Development of a Robust Process for the Preparation of High-Quality Dicyclopropylamine Hydrochloride. Org. Process Res. Dev. 2014, 18 (4), 520-527. (f) Shibue, T.; Fukuda, Y. Efficient Synthesis of Cyclopropylhydrazine Salts. Tetrahedron Lett. 2014, 55 (30), 4102-4104. (g) Derosa, J.; O'Duill, M. L.; Holcomb, M.; Boulous, M. N.; Patman, R. L.; Wang, F.; Tran-Dubé, M.; McAlpine, I.; Engle, K. M. Copper-Catalyzed Chan-Lam Cyclopropylation of Phenols and Azaheterocycles. J. Org. Chem. 2018, 83 (7), 3417-3425.

(9) (a) Mori-Quiroz, L. M.; Shimkin, K. W.; Rezazadeh, S.; Kozlowski, R. A.; Watson, D. A. Copper-Catalyzed Amidation of Primary and Secondary Alkyl Boronic Esters. *Chem. - Eur. J.* **2016**, *22* (44), 15654–15658. (b) Racine, E.; Monnier, F.; Vors, J.-P.; Taillefer, M. Direct N-Cyclopropylation of Secondary Acyclic Amides Promoted by Copper. *Chem. Commun.* **2013**, *49* (67), 7412–7414. (c) Sueki, S.; Kuninobu, Y. Copper-Catalyzed N - and O -Alkylation of Amines and Phenols Using Alkylborane Reagents. *Org. Lett.* **2013**, *15* (7), 1544–1547.

(10) Abdel-Magid, A. F.; Mehrman, S. J. A Review on the Use of Sodium Triacetoxyborohydride in the Reductive Amination of Ketones and Aldehydes. *Org. Process Res. Dev.* **2006**, *10* (5), 971–1031.

(11) Kong, D.; Moon, P. J.; Bsharat, O.; Lundgren, R. J. Direct Catalytic Decarboxylative Amination of Aryl Acetic Acids. *Angew. Chem.*, *Int. Ed.* **2020**, 59 (3), 1313–1319.

(12) An initial version of this work was deposited in ChemRxiv on September 7, 2020. Grayson, J. D.; Dennis, F. M.; Robertson, C. C.; Partridge, B. M. Chan-Lam Amination of Benzylic Secondary and Tertiary Boronic Esters. *ChemRxiv.* **2020**. DOI: 10.26434/chemrxiv.12925199.v2.

(13) Grayson, J. D.; Partridge, B. M. Mild Cu-Catalyzed Oxidation of Benzylic Boronic Esters to Ketones. *ACS Catal.* **2019**, *9* (5), 4296– 4301.

(14) For further details, see the Supporting Information.

(15) Lam, P. Y.; Vincent, G.; Clark, C. G.; Deudon, S.; Jadhav, P. K. Copper-Catalyzed General C-N and C-O Bond Cross-Coupling with Arylboronic Acid. *Tetrahedron Lett.* **2001**, *42* (20), 3415–3418.

(16) Vantourout, J. C.; Miras, H. N.; Isidro-Llobet, A.; Sproules, S.; Watson, A. J. B. Spectroscopic Studies of the Chan-Lam Amination: A Mechanism-Inspired Solution to Boronic Ester Reactivity. *J. Am. Chem. Soc.* **2017**, *139* (13), 4769–4779.

(17) Rossi, S. A.; Shimkin, K. W.; Xu, Q.; Mori-Quiroz, L. M.; Watson, D. A. Selective Formation of Secondary Amides via the Copper-Catalyzed Cross-Coupling of Alkylboronic Acids with Primary Amides. *Org. Lett.* **2013**, *15* (9), 2314–2317.

(18) (a) Liu, X.; Zhu, Q.; Chen, D.; Wang, L.; Jin, L.; Liu, C. Aminoazanium of DABCO: An Amination Reagent for Alkyl and Aryl Pinacol Boronates. *Angew. Chem., Int. Ed.* 2020, 59 (7), 2745–2749.
(b) Edelstein, E. K.; Grote, A. C.; Palkowitz, M. D.; Morken, J. P. A

Protocol for Direct Stereospecific Amination of Primary, Secondary, and Tertiary Alkylboronic Esters. Synlett 2018, 29 (13), 1749–1752. (c) Mlynarski, S. N.; Karns, A. S.; Morken, J. P. Direct Stereospecific Amination of Alkyl and Aryl Pinacol Boronates. J. Am. Chem. Soc. 2012, 134 (40), 16449–16451. (d) Bagutski, V.; Elford, T. G.; Aggarwal, V. K. Synthesis of Highly Enantioenriched C-Tertiary Amines From Boronic Esters: Application to the Synthesis of Igmesine. Angew. Chem., Int. Ed. 2011, 50 (5), 1080–1083. (e) Harris, M. R.; Li, Q.; Lian, Y.; Xiao, J.; Londregan, A. T. Construction of 1-Heteroaryl-3-Azabicyclo[3.1.0]Hexanes by Sp 3 -Sp 2 Suzuki-Miyaura and Chan-Evans-Lam Coupling Reactions of Tertiary Trifluoroborates. Org. Lett. 2017, 19 (9), 2450–2453.

(19) For selected examples, see: (a) Partridge, B. M.; Chausset-Boissarie, L.; Burns, M.; Pulis, A. P.; Aggarwal, V. K. Enantioselective Synthesis and Cross-Coupling of Tertiary Propargylic Boronic Esters Using Lithiation-Borylation of Propargylic Carbamates. *Angew. Chem., Int. Ed.* **2012**, *51* (47), 11795–11799. (b) de Meijere, A.; Khlebnikov, A. F.; Sünnemann, H. W.; Frank, D.; Rauch, K.; Yufit, D. S. Convenient Access to Various 1-Cyclopropylcyclopropane Derivatives. *Eur. J. Org. Chem.* **2010**, *2010* (17), 3295–3301.

(20) CCDC 2073100 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

(21) Kornfilt, D. J. P.; MacMillan, D. W. C. Copper-Catalyzed Trifluoromethylation of Alkyl Bromides. J. Am. Chem. Soc. 2019, 141 (17), 6853–6858.

(22) Skotnitzki, J.; Morozova, V.; Knochel, P. Diastereoselective Copper-Mediated Cross-Couplings between Stereodefined Secondary Alkylcoppers with Bromoalkynes. *Org. Lett.* 2018, 20 (8), 2365–2368.
(23) SADABS; Bruker: Madison, WI, 2016.

(24) Krause, L.; Herbst-Irmer, R.; Sheldrick, G. M.; Stalke, D. Comparison of Silver and Molybdenum Microfocus X-Ray Sources for Single-Crystal Structure Determination. *J. Appl. Crystallogr.* **2015**, *48* (1), 3–10.

(25) Sheldrick, G. M. {\it SHELXT} {-} Integrated Space-Group and Crystal-Structure Determination. *Acta Crystallogr., Sect. A: Found. Adv.* **2015**, 71 (1), 3–8.

(26) Sheldrick, G. M. Crystal Structure Refinement with SHELXL. Acta Crystallogr., Sect. C: Struct. Chem. **2015**, 71 (1), 3–8.

(27) Dolomanov, O. V.; Bourhis, L. J.; Gildea, R. J.; Howard, J. A. K.; Puschmann, H. OLEX2: A Complete Structure Solution, Refinement and Analysis Program. J. Appl. Crystallogr. **2009**, 42 (2), 339–341.

(28) Dennis, F. M.; Robertson, C. C.; Partridge, B. M. Nickel-Catalysed Allylboration of Aldehydes. *Synthesis* **2020**, *52* (13), 1903–1914.

(29) Bagutski, V.; Ros, A.; Aggarwal, V. K. Improved Method for the Conversion of Pinacolboronic Esters into Trifluoroborate Salts: Facile Synthesis of Chiral Secondary and Tertiary Trifluoroborates. *Tetrahedron* **2009**, *65* (48), 9956–9960.

(30) Noh, D.; Yoon, S. K.; Won, J.; Lee, J. Y.; Yun, J. An Efficient Copper(I)-Catalyst System for the Asymmetric Hydroboration of α -Substituted Vinylarenes with Pinacolborane. *Chem. - Asian J.* **2011**, 6 (8), 1967–1969.

(31) Noh, D.; Chea, H.; Ju, J.; Yun, J. Highly Regio- and Enantioselective Copper-Catalyzed Hydroboration of Styrenes. *Angew. Chem., Int. Ed.* **2009**, *48* (33), 6062–6064.

(32) Nandakumar, M.; Rubial, B.; Noble, A.; Myers, E. L.; Aggarwal, V. K. Ring-Opening Lithiation-Borylation of 2-Trifluoromethyl Oxirane: A Route to Versatile Tertiary Trifluoromethyl Boronic Esters. *Angew. Chem., Int. Ed.* **2020**, *59* (3), 1187–1191.

(33) Roesner, S.; Blair, D. J.; Aggarwal, V. K. Enantioselective Installation of Adjacent Tertiary Benzylic Stereocentres Using Lithiation-Borylation-Protodeboronation Methodology. Application to the Synthesis of Bifluranol and Fluorohexestrol. *Chem. Sci.* **2015**, 6 (7), 3718–3723.

(34) Aichhorn, S.; Bigler, R.; Myers, E. L.; Aggarwal, V. K. Enantiospecific Synthesis of Ortho-Substituted Benzylic Boronic Esters by a 1,2-Metalate Rearrangement/1,3-Borotropic Shift Sequence. J. Am. Chem. Soc. **2017**, 139 (28), 9519–9522.

(35) Lei, Q.; Wei, Y.; Talwar, D.; Wang, C.; Xue, D.; Xiao, J. Fast Reductive Amination by Transfer Hydrogenation "on Water. *Chem.* -*Eur. J.* **2013**, *19* (12), 4021–4029.

pubs.acs.org/joc

(36) Rucker, R. P.; Whittaker, A. M.; Dang, H.; Lalic, G. Synthesis of Hindered Anilines: Copper-Catalyzed Electrophilic Amination of Aryl Boronic Esters. *Angew. Chem., Int. Ed.* **2012**, *51* (16), 3953–3956.

(37) Johns, A. M.; Utsunomiya, M.; Incarvito, C. D.; Hartwig, J. F. A Highly Active Palladium Catalyst for Intermolecular Hydroamination. Factors That Control Reactivity and Additions of Functionalized Anilines to Dienes and Vinylarenes. J. Am. Chem. Soc. **2006**, 128 (6), 1828–1839.

(38) Su, Y.; Lu, M.; Dong, B.; Chen, H.; Shi, X. Silver-Catalyzed Alkyne Activation: The Surprising Ligand Effect. *Adv. Synth. Catal.* **2014**, 356 (4), 692–696.

(39) Lefranc, A.; Qu, Z.-W.; Grimme, S.; Oestreich, M. Hydrogenation and Transfer Hydrogenation Promoted by Tethered Ru-S Complexes: From Cooperative Dihydrogen Activation to Hydride Abstraction/Proton Release from Dihydrogen Surrogates. *Chem. - Eur. J.* **2016**, *22* (29), 10009–10016.

(40) Ge, X.; Qian, C.; Ye, X.; Chen, X. Asymmetric Reduction of Imines with Trichlorosilane Catalyzed by Valine-Derived Formamide Immobilized onto Magnetic Nano-Fe₃O₄. *RSC Adv.* **2015**, *5* (80), 65402–65407.

(41) Gatien, A. V.; Lavoie, C. M.; Bennett, R. N.; Ferguson, M. J.; McDonald, R.; Johnson, E. R.; Speed, A. W. H.; Stradiotto, M. Application of Diazaphospholidine/Diazaphospholene-Based Bisphosphines in Room-Temperature Nickel-Catalyzed C(Sp2)-N Cross-Couplings of Primary Alkylamines with (Hetero)Aryl Chlorides and Bromides. ACS Catal. 2018, 8 (6), 5328–5339.

(42) Emayavaramban, B.; Chakraborty, P.; Manoury, E.; Poli, R.; Sundararaju, B. Cp*Co(Iii)-Catalyzed N -Alkylation of Amines with Secondary Alcohols. *Org. Chem. Front.* **2019**, *6* (6), 852–857.

(43) Li, Q.; Hou, C.-J.; Liu, X.-N.; Huang, D.-Z.; Liu, Y.-J.; Yang, R.-F.; Hu, X.-P. Chiral Phosphine-Phosphoramidite Ligands for Highly Enantioselective Hydrogenation of N-Arylimines. *RSC Adv.* **2015**, *5* (18), 13702–13708.

(44) Sakai, N.; Takahashi, N.; Ogiwara, Y. Indium-Catalyzed Hydroamination/Hydrosilylation of Terminal Alkynes and Aromatic Amines through a One-Pot, Two-Step Protocol. *Eur. J. Org. Chem.* **2014**, 2014 (23), 5078–5082.

(45) Shahane, S.; Louafi, F.; Moreau, J.; Hurvois, J.-P.; Renaud, J.-L.; van de Weghe, P.; Roisnel, T. Synthesis of Alkaloids of Galipea Officinalis by Alkylation of an α -Amino Nitrile. *Eur. J. Org. Chem.* **2008**, 2008 (27), 4622–4631.

(46) Wu, Z.; Du, S.; Gao, G.; Yang, W.; Yang, X.; Huang, H.; Chang, M. Secondary Amines as Coupling Partners in Direct Catalytic Asymmetric Reductive Amination. *Chem. Sci.* **2019**, *10* (16), 4509–4514.

(47) Fleischer, S.; Zhou, S.; Junge, K.; Beller, M. An Easy and General Iron-Catalyzed Reductive Amination of Aldehydes and Ketones with Anilines. *Chem. - Asian J.* **2011**, *6* (9), 2240–2245.

(48) Saito, K.; Miyashita, H.; Akiyama, T. Asymmetric Transfer Hydrogenation of Ketimines by Indoline as Recyclable Hydrogen Donor. *Org. Lett.* **2014**, *16*, 5312–5315.

(49) Pan, H.-J.; Ng, T. W.; Zhao, Y. Iron-Catalyzed Transfer Hydrogenation of Imines Assisted by an Iron-Based Lewis Acid. *Org. Biomol. Chem.* **2016**, *14* (24), 5490–5493.

(50) Saito, K.; Miyashita, H.; Ito, Y.; Yamanaka, M.; Akiyama, T. Oxidative Kinetic Resolution of Acyclic Amines Based on Equilibrium Control. *Org. Lett.* **2020**, *22* (8), 3128–3134.

(51) Denmark, S. E.; Nakajima, N.; Stiff, C. M.; Nicaise, O. J. C.; Kranz, M. Studies on the Bisoxazoline- and (–)-Sparteine-Mediated Enantioselective Addition of Organolithium Reagents to Imines. *Adv. Synth. Catal.* **2008**, 350 (7–8), 1023–1045.

(52) Wei, Y.; Zhao, C.; Xuan, Q.; Song, Q. An Expedient and Novel Strategy for Reductive Amination by Employing H_2O as Both a Hydrogen Source and Solvent via $B_2(OH)_4/H_2O$ Systems. *Org. Chem. Front.* **2017**, *4* (12), 2291–2295.

(53) Grainger, R. S.; Betou, M.; Male, L.; Pitak, M. B.; Coles, S. J. Semipinacol Rearrangement of Cis -Fused β -Lactam Diols into Keto-

Bridged Bicyclic Lactams. Org. Lett. 2012, 14 (9), 2234–2237.

(54) Wang, M.; Xiao, F.; Bai, Y.; Hu, X. Reactions of Tertiary Allylic Amines and Dichlorocarbenes. *Synth. Commun.* **2015**, *45* (19), 2259–2265.

(55) Mishra, K.; Kim, S. H.; Lee, Y. R. Band-Gap Narrowing of Highly Stable Heterogeneous ZrO2-ZnO Nanocomposites for the Reductive Amination of Carbonyl Compounds with Formic Acid and Triethylamine. *ChemSusChem* **2018**, *12* (4), 881–889.

(56) Gandon, V.; Bertus, P.; Szymoniak, J. Zirconium-Catalyzed Ethylmagnesiation of Imines - Scope and Mechanism. *Eur. J. Org. Chem.* **2001**, 2001 (19), 3677–3681.

(57) Nguyen, Q. P. B.; Kim, T. H. S-Benzyl Isothiouronium Chloride as a Recoverable Organocatalyst for the Direct Reductive Amination of Ketones with Hantzsch Ester. *Synthesis* **2012**, *44* (13), 1977–1982.