# **Studies in Tertiary Amine Oxides**

Part V<sup> $\dagger$ </sup>—Carbon-13 Nuclear Magnetic Resonance Spectra of some N-Aryl Tertiary Amines, the Corresponding N-Oxides and the Meisenheimer Rearrangement Products

Abdul-Hussain Khuthier, Jasim M. A. Al-Rawi,\* Abdul K. Al-Kazzaz and Mohammed A. Al-Iraqi Department of Chemistry, College of Science, University of Mosul, Mosul, Iraq

The carbon-13 NMR spectra of some N-(2- or 4-nitrophenyl) tertiary amines and their corresponding N-oxides have been analysed. These N-oxides undergo thermal rearrangement to O-arylhydroxylamines, for which the <sup>13</sup>C NMR spectral assignments were also carried out. The N-oxidation effect on the aromatic and aliphatic fragments is calculated.

# **INTRODUCTION**

The chemistry of tertiary amine oxides has received considerable attention, partly because they play an important role in a number of biological systems<sup>1</sup> and partly because of their interesting thermal rearrangement to the substituted hydroxylamines. The isomerization of tertiary amine oxides in which a certain group migrates from N to O (e.g.  $R_1R_2R_3N \rightarrow O \rightarrow R_1R_2N \rightarrow O \rightarrow R_3$ ) is known as the Meisenheimer rearrangement.<sup>2</sup> The type of groups which can migrate to oxygen include allyl,<sup>3</sup> benzyl,<sup>4</sup> neopentyl,<sup>5</sup> homoadamantyl<sup>6</sup> and tetrachloropyridyl.<sup>7</sup> The benzene nucleus can undergo migration during the thermolysis of substituted aniline N-oxides<sup>8</sup> and, furthermore, a number of N-aryl cyclic amine Noxides have recently been shown to undergo isomerization to O-arylhydroxylamines<sup>9</sup> (Scheme 1).

There are few tertiary N-oxides for which  ${}^{13}C$  NMR spectral data are known. This paper describes the  ${}^{13}C$  NMR spectroscopy of the amines (1, 2) their N-oxides (3, 4) and their rearrangement products (5, 6), and the effect of the N-oxide on the chemical shifts of various carbons in the above compounds.

# 

# RESULTS

The assignment of the <sup>13</sup>C chemical shifts for the saturated heterocyclic ring

This was achieved for compounds 1-6 (Scheme 1) by using the reported values of the <sup>13</sup>C chemical shifts and the coupling pattern for analogous compounds.<sup>10,11</sup>

\* Author to whom correspondence should be addressed.

Scheme 1. The tertiary amines, the N-oxides and their rearrangement products.

# The assignment of the <sup>13</sup>C chemical shifts for the aromatic ring

#### N-Aryl-tert-amines 1 and 2.

(a) N-(4-Nitrophenyl)amines **1a-1f**: the broad band proton decoupled <sup>13</sup>C NMR spectra of the aromatic carbons in **1a-1f** (Table 1) were analysed using the

CCC-0030-4921/82/0018-0104\$02.50

<sup>†</sup> For Part IV, see Ref. 9.

Compound No.	C-1	C-2	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11	C-12
1a	154.4	110.4	126.1	137.3	126.1	110.4	40.2					_
Calc. <sup>b</sup>	156.5	113.6	123.2	136.7	123.2	113.6						
1b	155.1	112.4	126.0	137.8	126.0	112.4	48.4	25.4	24.2	25.4	48.4	
1c	154.8	112.3	126.1	137.5	126.1	112.3	49.8	30.5	18.4	25.5	41.8	14.5
1d	154.9	112.4	126.1	137.6	126.1	112.4	55.2	30.7	32.8	24.7	47.9	19.1
1e	154.9	112.5	126.1	137.6	126.1	112.5	47.8	33.5	30.7	33.5	47.8	21.6
1f	155.1	112.7	125.9	140.0	125.9	112.7	47.2	66.4		66.4	47.2	—
Calc.°	155.3	115.2	122.8	138.4	122.8	115.2						

Table 1. Carbon-13 chemical shift assignments relative to TMS for N-(4-nitrophenyl)amines (1)<sup>a</sup>

<sup>a</sup> The numbering of the carbon atoms corresponds with that in Scheme 1. <sup>b</sup> Calculated values were obtained using the NO<sub>2</sub> effect on the <sup>13</sup>C chemical shift of the aromatic carbons in *N*,*N*-dimethylaniline  $[\delta(CDCl_3), C-1 = 150.7; C-2,6 = 112.7; C-3,5 = 129.8; C-4 = 116.0 and C-7 = 40.3 ppm].$ 

° Calculated values were obtained using the NO<sub>2</sub> effect on the <sup>13</sup>C chemical shift of the aromatic carbons of *N*-phenylmorpholine  $[\delta(CDCl_3) C-1 = 149.5; C-2,6 = 114.3; C-3,5 = 127.6; C-4 = 118; C-7,11 = 48.7 and C-8 = 66.0 ppm].$ 

#### Table 2. Carbon-13 chemical shift assignments relative to TMS for N-(2-nitrophenyl)amines (2)<sup>a</sup>

Compound No.	C-1	C-2	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11	C-12
2a	146.2	139.5	126.5	118.3	133.2	117.9	42.3	_	_			
Calc. <sup>b</sup>	145.9	132.7	124.2	117.6	134.8	113.6						
2b	147.1	142.5	125.8	121.3	133.2	120.8	53.2	26.1	24.1	26.1	53.2	
2c	145.8	147.7	125.0	124.0	132.2	123.5	55.0	33.5	22.7	26.2	53.0	18.3
2d	146.9	142.7	126.0	121.0	133.4	120.6	59.7	31.1	32.6	25.5	52.6	19.11
2e	147.0	142.8	126.0	121.0	133.4	120.6	52.2	34.3	30.6	34.3	52.2	21.8
2f	146.2	144.8	125.8	122.5	133.5	121.3	52.3	66.9	_	66.9	52.3	_
Calc.°	144.7	134.3	122.8	119.3	133.4	115.2						

\* The numbering of the carbon atoms corresponds with that in Scheme 1. b,c See Table 1 footnotes.

Table 3. Carbon-13 chemic	al shift assignments	s relative to TMS :	for N-(4-nitro	phenyl)amine	N-oxides	(3)*
---------------------------	----------------------	---------------------	----------------	--------------	----------	------

Compound No.	C-1	C-2	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11	C-12
3a	158.2	122.0	125.0	148.2	125.0	122.0	62.6		_			
Calc. <sup>b</sup>	160.4	120.9	123.2	148.8	123.2	120.0						
Зb	161.2	122.5	124.5	147.9	124.5	122.5	69.2	21.3	21.9	21.3	69.2	_
3c	159.3	122.7	124.2	147.3	124.2	122.7	74.3	26.4	16.3	20.9	59.3	59.9
	(u) <sup>c</sup>	(u)	(u)	(u)	(u)	(u)	(71.3)	(28.7)	_	(21.0)		(15.2)
3d	160.8	122.4	124.5	147.9	124.5	122.4	75.2	26.9	30.5	21.2	68.7	18.6
3e	161.1	122.4	124.5	148.0	124.5	122.4	69.1	29.4	29.1	29.4	69.1	21.4
3f	159.6	122.3	124.9	148.1	124.9	122.3	68.0	62.2		62.2	68.0	—

\* The numbering of the carbon atoms corresponds with that in Scheme 1.

<sup>b</sup> Calculated values were obtained using the NO<sub>2</sub> effect on the <sup>13</sup>C chemical shift of the aromatic carbons in N,N-dimethylaniline N-oxide  $[\delta^{13}C(CDCl_3 50\% v/v) C-1 = 154.6; C-2,6 = 120.0; C-3,5 = 129.0; C-4 = 128.8 and C-7 = 63.5 ppm].$ 

<sup>c</sup> Values in parentheses are for the other rotamer of 3c.

u Indicates unresolved signal.

# Table 4. Carbon-13 chemical shift assignments relative to TMS for N-(2-nitrophenyl)amine N-oxides (4)\*

Compound No.	C-1	C-2	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11	C-12
4a	147.0	145.0	125.4	130.4	131.4	122.2	63.0		_			
Calc <sup>b</sup>	149.8	140.0	124.2	129.7	134.8	120.9						
4b	148.3	145.4	124.9	129.9	130.6	122.2	67.9	20.5	20.8	20.5	67.9	
4c	148.8°	147.9°	125.4	129.8	130.4	122.0	71.3	28.3	22.4	20.9	61.3	15.5
			(125.8) <sup>d</sup>	(u)	(u)	(122.7)	(71.7)	(26.3)	(u)	(20.6)	(u)	(16.1)
4d	147.7	145.7	125.2	129.8	130.6	121.1	74.4	26.4	30.4	20.7	68.1	18.3
			(125.5)	(u)	(130.3)	(u)	(73.9)	(28.4)	(29.3)	(19.8)	(69.9)	(19.3)
4e	147.0	145.8	125.3	130.2	131.0	121.4	68.4	28.9	28.8	28.9	68.4	21.8
4f	147.0	145.4	125.1	130.3	131.0	122.0	67.0	61.2	_	61.2	67.0	

 $^{\rm a}$  The numbering of the carbon atoms corresponds with that in Scheme 1.  $^{\rm b}$  See Table 3, footnote b.

<sup>c</sup> These assignments can be reversed.

<sup>d</sup> The values in parentheses are for the other rotamers of 4c and 4d.

u Indicates unresolved signal.

	C-2	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11	C-12
4.6 1	13.8	125.8	141.5	125.8	113.8	56.9	25.4	23.3	25.4	56.9	
5.9 1	14.1	125.8	141.8	125.8	114.1	62.6	34.2	23.8	25.7	56.7	19.6
4.6 1	13.8	125.8	141.7	125.8	113.8	67.2	31.8	31.8	24.9	64.0	19.3
4.6 1	13.9	125.8	141.6	125.8	113.8	56.5	33.9	30.0	33.9	56.5	21.2
3.9 1	13.8	126.0	135.5	126.0	113.8	56.3	66.0		66.0	56.3	
	1 4.6 1 5.9 1 4.6 1 4.6 1 3.9 1	1         C-2           4.6         113.8           5.9         114.1           4.6         113.8           4.6         113.9           3.9         113.8	1         C-2         C-3           4.6         113.8         125.8           5.9         114.1         125.8           4.6         113.8         125.8           4.6         113.9         125.8           3.9         113.8         126.0	1         C-2         C-3         C-4           4.6         113.8         125.8         141.5           5.9         114.1         125.8         141.8           4.6         113.8         125.8         141.7           4.6         113.9         125.8         141.6           3.9         113.8         126.0         135.5	1       C-2       C-3       C-4       C-5         4.6       113.8       125.8       141.5       125.8         5.9       114.1       125.8       141.8       125.8         4.6       113.8       125.8       141.7       125.8         4.6       113.9       125.8       141.6       125.8         3.9       113.8       126.0       135.5       126.0	1       C-2       C-3       C-4       C-5       C-6         4.6       113.8       125.8       141.5       125.8       113.8         5.9       114.1       125.8       141.8       125.8       114.1         4.6       113.8       125.8       141.7       125.8       113.8         4.6       113.9       125.8       141.6       125.8       113.8         3.9       113.8       126.0       135.5       126.0       113.8	-1       C-2       C-3       C-4       C-5       C-6       C-7         4.6       113.8       125.8       141.5       125.8       113.8       56.9         5.9       114.1       125.8       141.8       125.8       114.1       62.6         4.6       113.8       125.8       141.7       125.8       113.8       67.2         4.6       113.9       125.8       141.6       125.8       113.8       56.5         3.9       113.8       126.0       135.5       126.0       113.8       56.3	1       C-2       C-3       C-4       C-5       C-6       C-7       C-8         4.6       113.8       125.8       141.5       125.8       113.8       56.9       25.4         5.9       114.1       125.8       141.8       125.8       114.1       62.6       34.2         4.6       113.8       125.8       141.7       125.8       113.8       67.2       31.8         4.6       113.9       125.8       141.6       125.8       113.8       56.5       33.9         3.9       113.8       126.0       135.5       126.0       113.8       56.3       66.0	1       C-2       C-3       C-4       C-5       C-6       C-7       C-8       C-9         4.6       113.8       125.8       141.5       125.8       113.8       56.9       25.4       23.3         5.9       114.1       125.8       141.8       125.8       114.1       62.6       34.2       23.8         4.6       113.8       125.8       141.7       125.8       113.8       67.2       31.8       31.8         4.6       113.9       125.8       141.6       125.8       113.8       56.5       33.9       30.0         3.9       113.8       126.0       135.5       126.0       113.8       56.3       66.0       —	1       C-2       C-3       C-4       C-5       C-6       C-7       C-8       C-9       C-10         4.6       113.8       125.8       141.5       125.8       113.8       56.9       25.4       23.3       25.4         5.9       114.1       125.8       141.8       125.8       114.1       62.6       34.2       23.8       25.7         4.6       113.8       125.8       141.7       125.8       113.8       67.2       31.8       31.8       24.9         4.6       113.9       125.8       141.6       125.8       113.8       56.5       33.9       30.0       33.9         3.9       113.8       126.0       135.5       126.0       113.8       56.3       66.0        66.0	1       C-2       C-3       C-4       C-5       C-6       C-7       C-8       C-9       C-10       C-11         4.6       113.8       125.8       141.5       125.8       113.8       56.9       25.4       23.3       25.4       56.9         5.9       114.1       125.8       141.8       125.8       114.1       62.6       34.2       23.8       25.7       56.7         4.6       113.8       125.8       141.7       125.8       113.8       67.2       31.8       31.8       24.9       64.0         4.6       113.9       125.8       141.6       125.8       113.8       56.5       33.9       30.0       33.9       56.5         3.9       113.8       126.0       135.5       126.0       113.8       56.3       66.0        66.0       56.3

Table 5. Carbon-13 chemical shift assignments relative to TMS for O-(4-nitrophenyl)hydroxylamines (5)\*

<sup>a</sup> The numbering of the carbon atoms corresponds with that in Scheme 1.

#### Table 6. Carbon-13 chemical shift assignments relative to TMS for O-(2-nitrophenyl)hydroxylamine (6)\*

Compound No.	C-1	C-2	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11	C-12
6b	153.3	138.7	125.4	120.4	134.3	116.7	56.9	25.3	23.3	25.3	56.9	_
6c	154.4	137.7	125.4	120.0	134.2	116.8	62.6	34.2	23.8	25.9	56.9	19.8
6d	153.2	137.7	125.8	120.4	134.4	116.7	64.1	31.8	31.8	24.9	56.3	19.4
6e	153.3	137.8	125.4	120.4	134.4	116.4	56.5	34.0	30.0	34.0	56.5	21.2

\* The numbering of the carbon atoms corresponds with that in Scheme 1.

reported values for p-nitroaniline.<sup>12</sup> The good agreement between the calculated and the observed values for 1a and 1f (Table 1) supports the above analysis. The calculated values were obtained using the  $-NO_2$  effect<sup>13</sup> on the aromatic <sup>13</sup>C chemical shift of *N*,*N*dimethylaniline<sup>14</sup> and N-arylmorpholine, respectivelv.1

Inspection of the <sup>13</sup>C NMR spectra of **1a-1f**, all obtained using the same acquisition parameters, showed that the signals for the carbon bearing the nitro group are broader than that bearing the amine moiety, and this could be due to the Overhauser effect of the methyl (1a) or the methylene (1b-1f) groups connected to the nitrogen atom.

(b) N-(2-Nitrophenyl)amines **2a-2f**: the analysis of the <sup>13</sup>C chemical shifts for the aromatic ring carbons in **2a–2f** (Table 2) was carried out using the  ${}^{13}$ C chemical shift values for o-nitroaniline.<sup>16</sup> The agreement between the calculated and observed values for 2a and 2f supports the above assignment, except for C-2 where the discrepancy between the calculated and observed values could be attributed to steric interaction (induced shift).1'

# N-Aryl-tert-amine oxides 3 and 4.

(a) N-(4-Nitrophenyl)amine N-oxides 3a-3f: the <sup>13</sup>C NMR spectral assignments, carried out from the calculated shifts of 3a, are reported in Table 3.

(b) N-(2-nitrophenyl)amine N-oxides **4a-4f**: the broad band <sup>13</sup>C NMR spectra of **4a-4f** are reported in Table 4. The calculated <sup>13</sup>C signals for N,Ndimethylaniline N-oxide were used to confirm the spectral analysis for 4a. The difference between the calculated and observed <sup>13</sup>C chemical shift for C-2 in **4a** (~5 ppm) may be due to a steric effect.<sup>17</sup>

**O-Arvihvdroxylamines 5 and 6.** The proton decoupled <sup>13</sup>C spectra of **5** and **6** are reported in Tables 5 and 6, respectively. The assignments of the <sup>13</sup>C chemical carbon atoms in shifts for the arvl the 5b-5f and 2-nitro-4-nitrophenylhydroxylamines phenylhydroxylamines 6b-6f were made by using the reported <sup>13</sup>C chemical shift for p-nitro-<sup>18</sup> and o-nitrophenol, respectively.

#### DISCUSSION

#### The N-oxidation effect

#### The N-oxidation effect for the <sup>13</sup>C chemical shifts of N-(4nitrophenyl)amine N-oxides (3a-3f).

(a) N-oxidation effect on the aromatic ring: the substitution of one hydrogen in benzene by the N,Ndimethylamine group induces an upfield shift for the o- and p-carbons of -16 and -12 ppm, respectively. This has been attributed to the mesomeric effect, in which the paired electrons of the nitrogen are used up in the resonance.19



This resonance is interrupted by N-oxidation of the molecule, and so a deshielding is expected for the ortho- and para-carbons. Indeed, C-2 and C-4 are deshielded by +7.3 and +12.8 ppm, respectively, C-1 experiences a deshielding of +3.9 ppm and C-3,5 show no effect (Tables 1 and 3). Table 7 shows a similar pattern for the N-oxide effect on the p-nitrophenyl group in 3a-3f.

(b) The N-oxidation effect on the aliphatic carbon atoms: a downfield shift in the range of +17.5 to +24.5 ppm is observed for the ring carbons C-7 and C-11 ( $\beta$ -effect) due to N-oxidation (Table 7). Shielding is observed for the ring C-8 and C-10 carbons  $(-3.5 \text{ to } -4.6 \text{ ppm}; \gamma\text{-effect})$  and for C-9 (-1.6 to2.3 ppm;  $\delta$ -effect).

#### The N-oxidation effect for the N-(2-nitrophenyl)amines (4a-4f).

(a) N-oxidation effect on the aromatic ring: the N-oxidation effect for **4a-4f** is indicated in Table 8.

Compounds	ΔδC-1	ΔδC-2	<b>∆</b> δC-3	ΔδC-4	Δ8C-5	ΔδC-6	ΔδC-7	ΔδC-8	ΔδC-9	ΔδC-10	Δ8C-11	ΔδC-12
3a-1a	+3.8	+11.6	-1.1	+10.4	1.1	+11.6	+22.4			—		
3b–1b	+6.1	+10.1	-1.5	+10.1	-1.5	+10.1	+20.8	-4.1	2.3	-4.1	+20.8	
3c-1c	+4.5	+10.4	1.9	+9.8	-1 <i>.</i> 9	+10.4	+24.5	-4.1	-2.2	-4.6	+17.5	+1.4
3d1d	+5.9	+10.0	-1.6	+10.3	-1.6	+10.0	+20.0	-3.8	-2.3	-3.5	+20.8	-0.5
3e1e	+6.2	+9.9	-1.6	+10.4	-1.6	+9.9	+21.3	-4.1	-1.6	-4.1	+21.3	-0.2
3f-1f	+4.5	+9.6	-1.0	+8.1	-1.0	+9.6	+20.8	-4.2		-4.2	+20.8	

Table 7. N-oxidation effect on the <sup>13</sup>C chemical shift of N-(4-nitrophenyl)amines (1)\*

\*  $\Delta\delta C = \delta C$  of the *N*-oxide (3) –  $\delta C$  of the corresponding amine (1).

Table 8. N-oxidation effect on the <sup>13</sup>C chemical shift of N-(2-nitrophenyl)amines(2)<sup>a</sup>

Compounds	<b>Δ</b> 8C-1	<b>ΔδC-2</b>	<b>ΔδC-3</b>	∆8C-4	∆8C-5	ΔδC-6	ΔδC-7	ΔδC-8	∆8C-9	ΔδC-10	<u>∆</u> δC-11	Δ8C-12
4a2a	+0.8	+5.5	-1.1	+12.1	-1.8	+5.3	+20.7	_				—
4b2b	+1.2	+2.9	-0.9	+8.6	-2.6	+1.4	+14.7	-5.6	-3.3	-5.6	+14.3	
4c-2c	+3.0	+0.2	+0.4	+5.8	-1.8	-1.5	+16.3	-5.2	+0.2	5.6	8.3	2.8
4d-2d	+0.8	+3.0	-0.5	+8.8	-2.8	+0.6	+14.7	-4.7	-2.2	-4.8	+15.5	-0.8
4e2e	0	+3.1	0.7	+9.2	-2.4	+0.8	+16.2	-5.4	-1.8	-5.4	16.2	0
4f-2f	+0.8	+0.6	+0.7	+7.8	-2.5	+0.7	14.7	-5.7		-5.7	14.7	

 $\Delta\delta C = \delta C$  of the *N*-oxide (4) –  $\delta C$  of the corresponding amine (2).

Table 9. N-oxidat dimethyl	tion effect aniline	on (	the <sup>13</sup> C	chemical	shift	of 3-nit	ro-N,N-
	C-1	C-2	C-3	C-4	C-5	C-6	C-7
Amine <sup>a</sup>	150.8	106.0	149.4	110.5	129.5	117.6	40.1
N-oxide <sup>b</sup>	153.3	116.0	148.8	124.8	131.1	131.1	62.2
N-oxide effect	+2.5	+10	-0.8	+14.3	+1.6	13.5	12.1

<sup>a</sup> The assignment of 3-nitro-*N*,*N*-dimethylaniline was achieved using the reported value of *m*-nitroaniline.<sup>1</sup>

<sup>b</sup> The assignment of the N-oxide was achieved using the  $-NO_2$  effect on the <sup>13</sup>C chemical shifts of N,N-dimethylaniline N-oxide.

Compound **4a** showed a similar pattern on Noxidation as N,N-dimethylaniline. The effect of Noxidation on the aromatic ring in **4b-4f** showed a similar pattern for C-1, C-3 and C-5 to that of the corresponding carbons in **4a**. However, the effect on C-2 and C-6 (*ortho* effect) and C-4 (*para* effect) is much smaller than in **4a**, although in the same direction. This could be due to steric interaction between the neighbouring NO<sub>2</sub> group and the heterocyclic ring in the amine. This prevents the conjugated system from attaining coplanarity, thereby reducing the electronic interaction between the aromatic  $\pi$ -system and

the -N lone pair of electrons, and canonical forms

such as



are, thus, less important than in the p- and m-NO<sub>2</sub> aromatic compounds. Accordingly, **4b**-**4f** did not show the true *N*-oxidation effect seen with **4a** or with the *p*-nitro compounds (**3a**-**3f**) and 3-nitro-*N*,*N*-dimethylaniline (see Table 9). Support for the above argument is given by the <sup>13</sup>C chemical shift values of **2c** in which a methyl group is present in the heterocyclic ring at the 2-position, making steric crowding in this compound even more severe than in the other isomers. C-2, C-4 and C-6 are more deshielded in **2c** than in the other congeners (Table 2).

Dreiding models of 2c and 4c show clearly that the

o-nitrophenyl group is not coplanar with the heterocyclic ring, but is nearly perpendicular due to the steric hindrance to rotation. It is of interest to note the appearance of two <sup>13</sup>C signals for some of the aromatic and heterocyclic ring carbon atoms in **4c** (Table 4), thus suggesting the possibility of the existence of two rotational isomers. The same phenomenon is noted in **3c** and **4d**.

(b) The N-oxidation effect on the aliphatic carbon atoms: it is clear from Tables 7 and 8 that the Noxidation effect on the N-methyl carbon atoms in **4a** has a similar value as in the p-nitro isomer **3a**. However, in **4b**-**4f**, the N-oxidation effect on the N—CH<sub>2</sub> carbon atoms (C-7 and C-11) has a smaller value than for the N—CH<sub>3</sub> carbon in **4a**. The difference is due to the steric effect induced by the presence of the  $-NO_2$ group in the ortho-position, positioning the N—CH<sub>2</sub> carbon out of the electrical field effect of the aromatic ring. Indeed, the situation becomes more severe in compound **4c**, which has a methyl group at C-7.

# EXPERIMENTAL

# Synthesis

The tertiary amines were synthesized by nucleophilic substitution of the appropriate secondary amines with o-nitro- and p-nitro-fluorobenzene in dimethyl sulphoxide.<sup>9,20</sup> The N-oxides were prepared by oxidation

of the tertiary amines with performic acid (a mixture of 98% formic acid and 30% hydrogen peroxide).<sup>9,21</sup> Rearrangement of the tertiary amine N-oxides was performed by heating the free base in dioxane for 3 h.<sup>9</sup> The products were purified by column chromatography with neutral alumina. The preparative details of these compounds are reported in Ref. 9.

# **Carbon-13 NMR spectra**

Proton decoupled and coupled spectra were obtained

- R. Kuntzman, A. Phillips, I. Isai, A. Klutch and J. J. Burns, J. Pharmacol. Exp. Ther. **155**, 337 (1966); D. M. Ziegler and F. H. Pettit, Biochemistry **5**, 2932 (1966); J. R. Gillette, Adv. Pharmacol. **4**, 219 (1966).
- J. Meisenheimer, Ber. Dtsch. Chem. Ges. 42, 1667 (1919); Ber. Dtsch. Chem. Ges. 55, 513 (1922).
- R. F. Kleinschmidt and A. C. Cope. J. Am. Chem. Soc. 66, 1929 (1944).
- U. Schollkopf, M. Patsch and H. Schafe, Tetrahedron Lett. 2515 (1964).
- 5. I. I. Brauman and W. A. Sanderson, Tetrahedron 23, 37 (1967).
- B. L. Adams and P. Kovacic, J. Am. Chem. Soc. 96, 7014 (1974).
- 7. S. M. Roberts and H. Suschitzky, J. Chem. Soc. C 1537 (1968).
- 8. A. H. Khuthier, T. Y. Ahmed and L. I. Jallo, J. Chem. Soc. Chem. Commun. 1001 (1976).
- 9. A. H. Khuthier, A. K. S. Al-Kazzaz, J. M. A. Al-Rawi and M. A. Al-Iraqi, *J. Org. Chem.* accepted for publication.
- 10. M. A. Al-Iraqi, J. M. Al-Rawi and A. H. Khuthier, Org. Magn. Reson. 14, 161 (1980).
- 11. Jasim M. A. Al-Rawi and A. H. Khuthier, Org. Magn. Reson. accepted for publication.

on a Bruker WH 90 DS spectrometer equipped with an Aspect 2000, 32 K computer, operating at 22.63 MHz for <sup>13</sup>C NMR with internal deuterium lock. The spectral width was 6024 Hz; a flip angle of 45° was used and the free induction decay was accumulated in 8 K, and the frequency domain spectra obtained in 4 K, data points. Samples were dissolved in CDCl<sub>3</sub> 10–15% by v/v or w/v in a 10 mm o.d. Wilmad tube, and a few drops of TMS were added as internal reference.

#### REFERENCES

- 12. V. Formacek, L. Desnoyer, H. P. Kellerhals, T. Keller and J.
- T. Clerc, Carbon-13 Data Bank VI, p. 89. Bruker (1976).
  13. F. W. Wehrli and T. Wirthlin, Interpretation of Carbon-13 NMR Spectra, p. 47. Heyden, London (1978).
- V. Formacek, L. Desnoyer, H. P. Kellerhals, T. Keller and J. T. Clerc, Carbon-13 Data Bank, p. 193. Bruker (1976).
- V. Formacek, L. Desnoyer, H. P. Kellerhals, T. Keller and J. T. Clerc, Carbon-13 Data Bank, p. 560. Bruker (1976).
- V. Formacek, L. Desnoyer, H. P. Kellerhals, T. Keller and J. T. Clerc, Carbon-13 Data Bank, p. 87. Bruker (1976).
- 17. D. A. Netzel, Org. Magn. Reson. 11, 58 (1978).
- V. Formacek, L. Desnoyer, H. P. Kellerhals, T. Keller and J. T. Clerc, Carbon-13 Data Bank, p. 361. Bruker (1976).
- 19. J. B. Stothers, *Carbon-13 NMR Spectroscopy*, p. 428. Academic Press, New York (1972).
- 20. J. H. Suhr, Ann. Chem. 689, 109-117 (1965).
- S. M. Roberts and H. Suschitzky, J. Chem. Soc. C 1537 (1968).

Received 28 May 1981; accepted 23 September 1981

© Heyden & Son Ltd, 1982