

Preliminary communication

METALLATION OF ALIPHATIC CARBON ATOMS

II*, SYNTHESIS AND CHARACTERIZATION OF THE CYCLOPALLADATED COMPLEXES OF *N,N*-DIMETHYLNEOPENTYLAMINE

YOSHIO FUCHITA, KATSUMA HIRAKI* and YASUHIRO MATSUMOTO

Department of Industrial Chemistry, Faculty of Engineering, Nagasaki University, Bunkyo-machi, Nagasaki 852 (Japan)

(Received August 18th, 1984)

Summary

N,N-Dimethylneopentylamine reacts with $\text{Pd}(\text{MeCO}_2)_2$ to give a novel trinuclear cyclopalladated complex $[\text{Me}_2\text{NCH}_2\text{CMe}_2\text{CH}_2\text{Pd}(\mu\text{-MeCO}_2)_2\text{Pd}(\mu\text{-MeCO}_2)_2\text{PdCH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2] \cdot 0.5\text{C}_6\text{H}_6$ (I). The reaction of I with PPh_3 affords both *trans*- $[\text{Pd}(\text{MeCO}_2)_2(\text{PPh}_3)_2]$ (II) and $[\text{Pd}(\text{CH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2)(\text{MeCO}_2)(\text{PPh}_3)]$ (III). The reaction of III with LiCl yields a mononuclear cyclopalladated complex, $[\text{Pd}(\text{CH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2)\text{Cl}(\text{PPh}_3)]$ (IV).

Metallation of an aliphatic carbon atom has been one of the current topics in organometallic chemistry in association with the activation of C—H bonds by transition metals [1]. Recently, we [2] reported the syntheses of the six-membered cyclopalladated complexes of 2-neopentylpyridine through direct metallation of the aliphatic carbon atom by use of $\text{Pd}(\text{MeCO}_2)_2$. This communication deals with the cyclopalladation of *N,N*-dimethylneopentylamine by $\text{Pd}(\text{MeCO}_2)_2$, resulting in the formation of a novel trinuclear cyclopalladated complex.

N,N-Dimethylneopentylamine (5.5 mmol) reacted with $\text{Pd}(\text{MeCO}_2)_2$ (4.45 mmol) in benzene (20 ml) at 50–55°C for 1.5 h. After the mixture was filtered to remove the precipitated palladium black, the filtrate was concentrated under reduced pressure and diluted with hexane to give a novel trinuclear cyclopalladated complex, $[\text{Me}_2\text{NCH}_2\text{CMe}_2\text{CH}_2\text{Pd}(\mu\text{-MeCO}_2)_2\text{Pd}(\mu\text{-MeCO}_2)_2\text{PdCH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2] \cdot 0.5\text{C}_6\text{H}_6$ * (I) in 36% yield (Scheme 1).

*Satisfactory elemental analysis and ^1H NMR data were obtained.

Complex I was not isolated from the reaction mixture either in acetic acid at 70°C or in methanol at 60°C.


When *N,N*-dimethylnopentylamine was treated with $\text{Pd}(\text{MeCO}_2)_2$ in a molar ratio of 2/3 or 1/1 in CDCl_3 at 19°C, it was observed by ^1H NMR spectroscopy that I was formed smoothly. As for the reaction in the molar ratio of 2/1, I was produced in lower yield (about 10% after 4 h), and a large amount of palladium black was formed. In no case was any other cyclopalladated complex, such as a binuclear one, detected.

Complex I reacted with PPh_3 to produce *trans*- $[\text{Pd}(\text{MeCO}_2)_2(\text{PPh}_3)_2]$ (II) in 56% yield, as well as an oily product, $\text{Pd}(\text{CH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2)(\text{MeCO}_2)_2(\text{PPh}_3)^*$ (III). The IR spectrum of II coincided quite well with that of the authentic sample [3]. The oily product III was uncrystallizable and was converted into $\text{Pd}(\text{CH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2)\text{Cl}(\text{PPh}_3)$ (IV) in 55% yield (based on I) by reaction with LiCl.

The IR spectrum of I exhibited $\nu(\text{COO}^-)$ frequencies due to bridging acetato groups at 1610(s), 1570(s), 1405(s), 1380(s), and 1335(s) cm^{-1} , which were very similar to those of the trimeric palladium(II) acetate [3]. The ^1H NMR spectrum of IV in CDCl_3 exhibited two methylene proton signals at δ 1.34 (d, $^3J(\text{HP})$ 4.5 Hz; $\text{Pd}-\text{CH}_2$) and 2.50 ppm (s, $\text{N}-\text{CH}_2$) and two methyl proton signals at δ 1.02 (s, CMe_2) and 2.89 ppm (d, $^4J(\text{HP})$ 2.5 Hz; NMe_2). The resonance at δ 1.34 ppm was assigned to the protons of the palladium-bonded methylene group which was produced by the cyclopalladation of *N,N*-dimethylnopentylamine. The coupling of the protons of both $\text{Pd}-\text{CH}_2$ and NMe_2 with the ^{31}P nucleus confirmed strongly the presence of a $\text{CH}_2 \dots \text{NMe}_2$ chelate structure. The small coupling constant $^3J(\text{HP})$ for $\text{Pd}-\text{CH}_2$ suggests that PPh_3 is situated in a *cis* position to the palladated methylene carbon. These facts indicate unambiguously that I contains both the $\text{Pd}(\text{CH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2)(\text{MeCO}_2)_2$ and $\text{Pd}(\text{MeCO}_2)_2$ moieties. On the basis of these discussions, the elemental analysis, and the molecular weight (745.1 in benzene, calcd. 783.9), complex I was assigned to have trinuclear structure, which involves four acetato bridges and two $\text{Me}_2\text{NCH}_2\text{CMe}_2\text{CH}_2-\text{C}^1, \text{N}$ chelates. It is noteworthy that *N,N*-dimethylnopentylamine reacted with $\text{Pd}(\text{MeCO}_2)_2$ in hot benzene to afford directly the trinuclear cyclopalladated complex, I. Ukhin et al. [4] reported that $\text{Pd}(\text{MeCO}_2)_2$ reacted with 2,6-disubstituted pyrylium salts to afford trinuclear bis(1-3- η -1,3-diacylallyl)tripalladium(II) complexes having four μ -acetato ligands.

As for the ^1H NMR spectrum of I in CDCl_3 at 14°C, the acetato-methyl protons appeared as four singlets at δ 1.78 (2.4H), 1.80 (2.4H), 1.82 (3.6H), and 1.84 ppm (3.6H), indicating that I contains both an (*a*- C^1 , *b*-*N*)-(*k*- C^1 , *l*-*N*)-type (*ak*-type) isomer and an (*a*- C^1 , *b*-*N*)-(*k*-*N*, *l*- C^1)-type (*al*-type) isomer (Scheme 1) in a ratio of about 2/3. In addition, I exhibited two sets of signals corresponding to the two isomers. At 14°C, one set consisted of six signals at δ 1.33 (s, H_3CCCH_3), 1.36 (s, H_3CCCH_3), 2.40 (q, $\Delta\delta$ 0.29 ppm, $^2J(\text{HH})$ 8 Hz, $\text{Pd}-\text{CH}_2$), 2.45 (q, $\Delta\delta$ 0.29 ppm, $^2J(\text{HH})$ 9 Hz, $\text{N}-\text{CH}_2$), 2.88

*The ^1H NMR spectrum of III was very similar to that of IV, except for the acetato-methyl proton resonance at δ 1.54 ppm.



SCHEME 1. Reagents: i, PPh_3 ; ii, LiCl .

(s, H_3CNCH_3), and 3.01 ppm (s, H_3CNCH_3) and was temperature-dependent. At 43°C the two C-methyl proton signals united to give one singlet at δ 1.34 ppm, whereas the two N-methyl proton signals coalesced into one singlet at δ 2.98 ppm; then, the two quartets changed into broad signals. The other set involved four singlets at δ 1.27 [$\text{C}(\text{CH}_3)_2$], 2.73 ($\text{Pd}-\text{CH}_2$), 2.93 ($\text{N}-\text{CH}_2$), and 3.01 ppm [$\text{N}(\text{CH}_3)_2$] * and virtually did not change in the range -35 – 43°C . The temperature-dependency of the ^1H NMR spectra of I was associated with the inversion of the acetato bridges, as observed for the other binuclear acetato-bridged complexes [5,6]. The inversion corresponding to the former set was actually quenched below 20°C , whereas that corresponding to the latter set was not quenched even at -35°C .

The $^{13}\text{C}\{-^1\text{H}\}$ NMR data of I in CDCl_3 at 24°C are quite consistent with the presence of the two isomers, showing four carboxylato carbon signals at δ 181.6, 181.9, 183.4, and 183.5 ppm, two acetato-methyl carbon signals at δ 22.4 and 23.4 ppm, and two C-methyl carbon signals at δ 28.3 and 28.6 ppm. In addition, I exhibited four singlets at δ 35.9 ($\text{Pd}-\text{CH}_2$), 43.8 (CMe_2), 54.3 [$\text{N}(\text{CH}_3)_2$], and 80.3 (NCH_2).

The stability of I, III, and IV is associated with both the chelate structure of the $\text{PdCH}_2\text{CMe}_2\text{CH}_2\text{NMe}_2$ moiety and the lack of β -hydrogen in the 3-(dimethylamino)-2,2-dimethylpropyl group.

Acknowledgements. The present work was partially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education.

*The singlet at δ 3.01 ppm was ascribed to protons of one of two N-methyl groups of the former set and those of two N-methyl groups of the latter set.

References

- 1 D.E. Webster, *Adv. Organomet. Chem.*, 15 (1977) 147.
- 2 Y. Fuchita, K. Hiraki and T. Uchiyama, *J. Chem. Soc., Dalton Trans.*, (1983) 897.
- 3 T.A. Stephenson, S.M. Morehouse, A.R. Powell, J.P. Heffer and G. Wilkinson, *J. Chem. Soc.*, (1965) 3652.
- 4 L.Yu. Ukhin, N.A. Dolgoplova, L.G. Kuz'mina and Yu.T. Struchkov, *J. Organomet. Chem.*, 210 (1981) 263.
- 5 J. Powell, *J. Am. Chem. Soc.*, 91 (1969) 4311.
- 6 A.J. Deeming and I.P. Rothwell, *J. Organomet. Chem.*, 205 (1981) 117.

JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 280, NO. 2

AUTHOR INDEX

- | | | |
|-------------------------|----------------------|-------------------------|
| Arad-Yellin, R., 197 | Haaland, A., C43 | Oro, L.A., 261 |
| Assadourian, L., 153 | Häberle, K., 183 | |
| | Harris, R.K., 281 | Packer, K.J., 281 |
| Baccar, B., 165 | Hiraki, K., C51 | |
| Bellassoued, M., 165 | Horn, E., 289 | Pete, J.P., 269 |
| Bin Shawkataly, O., 289 | Hudson, A., 173 | Pinillos, M.T., 261 |
| Black, P., 159 | | |
| Bruce, M.I., 289 | Jackson, R.A., 173 | |
| Butler, I.R., C47 | | Raubenheimer, H.G., 241 |
| Bywater, S., 159 | Klein, H.-P., 203 | Reams, P., 281 |
| | Klein, S.I., 281 | Rhodes, C.J., 173 |
| Cullen, W.R., C47 | Klement, U., 215 | Riahi, A., 269 |
| | Kreiter, C.G., 225 | Ricci, A., 177 |
| Del Vecchio, A.L., 173 | Kroto, H.W., 281 | Riess, J.G., 215 |
| Döppert, K., 203 | Kruger, G.J., 241 | |
| Doyle, G., 253 | | Schilling, B.E.R., C43 |
| Dräger, M., 183 | Lachance, P., 159 | Shawkataly, O., 289 |
| Dyk, M. Van, 241 | Lappert, M.F., C43 | Snow, M.R., 289 |
| | Leyendecker, M., 225 | |
| El Borgi, A., 165 | Linford, L., 241 | |
| Engen, D. Van, 253 | Lotz, S., 241 | Tejel, C., 261 |
| | | Thewalt, U., 203 |
| Faure, R., 153 | Mann, B.E., C47 | Thorne, A.J., C43 |
| Fiorenza, M., 177 | Matsumoto, Y., C51 | |
| Fjeldberg, T., C43 | Meidine, M.F., 281 | Van Dyk, M., 241 |
| Fuchita, Y., C51 | Mordini, A., 177 | Van Engen, D., 253 |
| | Muzart, J., 269 | Volden, H.V., C43 |
| | | |
| Gau, G., 153 | Nixon, J.F., 281 | Wachter, J., 215 |
| Gaudemar, M., 165 | Nurse, C.R., C47 | Wudl, F., 197 |