## Diazoalkanes react with a bis(phosphino)borate copper(I) source to generate $[Ph_2BP^{tBu}_2]Cu(\eta^1-N_2CR_2)$ , $[Ph_2BP^{tBu}_2]Cu(CPh_2)$ , and $[Ph_2BP^{tBu}_2]Cu-N(CPh_2)(NCPh_2)^{\dagger}$

Neal P. Mankad and Jonas C. Peters\*

Received (in Berkeley, CA, USA) 6th September 2007, Accepted 14th December 2007 First published as an Advance Article on the web 17th January 2008 DOI: 10.1039/b713687k

 $[Ph_2BP^{tBu}_2]Cu-L \ complexes \ react \ with \ diazoalkanes \ to \ generate \ structurally \ unusual \ \eta^1-diazoalkane \ adducts, \ a \ terminal \ carbene, \ and \ an \ \eta^1-azine \ adduct.$ 

Metal-catalyzed carbene transfer from diazoalkanes to organic substrates has become a versatile technique in synthesis, and copper catalysts have been particularly well studied in this regard.<sup>1,2</sup> Careful study of copper carbene complexes (Cu=CR<sub>2</sub>), the presumed reactive intermediates in these reactions, serves to deepen understanding of these coppercatalyzed transformations. Identifying factors that affect copper carbene stability, characterizing specific decomposition pathways, and understanding the nature of copper–diazoalkane interactions prior to copper–carbene bond formation are therefore issues of fundamental interest.

Whereas  $\alpha$ -heteroatom stabilized copper carbenes are well known,<sup>3</sup> only two examples of non-heteroatom stabilized copper carbenes have been reported. Both of these derivatives are stabilized by bidentate, monoanionic N-chelates. The first such example was reported by Straub and Hofmann in 2001 and assigned using spectroscopic data.<sup>4</sup> Warren and co-workers later reported X-ray structural data for a  $\beta$ -diketiminato Cu=CPh<sub>2</sub> complex.<sup>5</sup> Both of these systems mediate catalytic cyclopropanation reactions and employ diazoalkane precursors to generate the carbenes of interest. In this context we note that the presumed diazoalkane adduct intermediates *en route* to carbene formation were not observed, and to our knowledge no such species have been previously characterized for copper systems.

Our group has had an interest in exploiting (phosphino)borate ligands to stabilize mid-to-late first-row metals featuring metal ligand multiple bonds.<sup>6</sup> In this paper, we describe the utility of the bulky bis(phosphino)borate ligand<sup>7</sup> [Ph<sub>2</sub>BP<sup>tBu</sup><sub>2</sub>] = Ph<sub>2</sub>B(CH<sub>2</sub>P<sup>t</sup>Bu<sub>2</sub>)<sub>2</sub> to generate unusual diazoalkane adducts of Cu<sup>I</sup>. In addition, we establish that N<sub>2</sub>CPh<sub>2</sub> reacts to generate the copper carbene complex [Ph<sub>2</sub>BP<sup>tBu</sup><sub>2</sub>]Cu=CPh<sub>2</sub> as verified by NMR spectroscopy and group transfer of the carbene unit. This latter species is unstable in the presence of additional N<sub>2</sub>CPh<sub>2</sub> in solution and as such generates the structurally novel azine adduct  $[Ph_2BP^{tBu}_2]$ -Cu–N(CPh<sub>2</sub>)(NCPh<sub>2</sub>) as a side product.

The synthesis of copper(1) species of the type  $[Ar_2BP_2]CuL$ (L = e.g. CH<sub>3</sub>CN, PR<sub>3</sub>, CO) was reported recently.<sup>8</sup> Preliminary group transfer studies with diazoalkanes suggested the necessity of a  $[Ar_2BP_2]CuL$  precursor with a sufficiently labile L ligand to expose reactions of interest. The aniline adduct  $[Ph_2BP^{1Bu}_2]Cu(NH_2Mes)$  (Mes = 2,4,6-trimethylphenyl) proved key in this regard. Its synthesis (Scheme 1) proceeded from the neutral pyridine adduct  $[Ph_2BP^{1Bu}_2]Cu-$ (pyridine) (1). Reaction of LiNHMes and excess 12-crown-4 with yellow 1 provided the anionic colorless amido complex  $\{[Ph_2BP^{tBu}_2]CuNHMes\}\{Li(12-crown-4)_2\}$  (2). Protonation of 2 with HBF<sub>4</sub> (toluene, -90 °C) produced the colorless aniline adduct  $[Ph_2BP^{tBu}_2]Cu(NH_2Mes)$  (3). Recrystallization of 3 from a THF-petroleum ether mixture provided the THF adduct  $[Ph_2BP^{tBu}_2]Cu(THF)$  (4).

The X-ray crystal structures of complexes **2** and **3**, which are related by a proton transfer, are shown in Fig. 1. The crystal structure of **3** features a longer Cu–N distance (2.0758(3) Å) and more compressed Cu–N–C angle (118.46(2)°) when compared to its conjugate base **2** (1.962(1) Å and 136.43(1)°, respectively).‡ Gunnoe and co-workers have recently reported that the structurally related but neutral complex (dtbpe)-Cu(NHPh) (dtbpe =  ${}^{t}Bu_{2}PCH_{2}CH_{2}P^{t}Bu_{2}$ ) has an appreciably shorter Cu–N distance of 1.890(6) Å.<sup>9</sup>

When either **3** or **4** were mixed with  $N_2C(SiMe_3)_2$  or  $N_2CMes_2$ , the stable diazoalkane adducts yellow  $[Ph_2BP^{tBu}_2]$ - $Cu(\eta^1-N_2C(SiMe_3)_2)$  (**5**) and red  $[Ph_2BP^{tBu}_2]Cu(\eta^1-N_2CMes_2)$  (**6**) formed rapidly and could be readily isolated (Scheme 1).



Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA. E-mail: jcpeters@mit.edu

<sup>&</sup>lt;sup>†</sup> Electronic supplementary information (ESI) available: Synthetic details, crystal structure diagrams of complexes 1–7 and spectroscopic data. See DOI: 10.1039/b713687k



Fig. 1 Solid-state structure of  $\{[PhBP^{tBu}_2]Cu(NH_2Mes)\}\{Li(12-crown-4)_2\}$  (2) (top) and  $[PhBP^{tBu}_2]Cu(NH_2Mes)$  (3) (bottom). Only the anion of 2 is shown. N–H hydrogens are shown in calculated positions.

We were surprised to find that these adducts are the only characterized examples of Cu<sup>1</sup>-diazoalkane complexes to be reported, though Hofmann and co-workers have described a formally Cu<sup>III</sup>-diazoalkane complex resulting from reduction of a chelating diazoalkane unit by a Cu<sup>I</sup> precursor.<sup>10</sup> The observed  $N_2CR_2$  <sup>13</sup>C{<sup>1</sup>H} chemical shifts ( $\delta$  29.5 for **5** and  $\delta$  99.3 for **6**) and  $CN_2$  vibrational frequencies (2108 cm<sup>-1</sup> for 5 and 2041  $cm^{-1}$  for 6) are very similar to the corresponding values for the free diazoalkanes,11 indicating that the diazoalkane moieties have not been significantly perturbed upon coordination to copper. The molecular structures of 5 and 6 (Fig. 2) feature N-N distances (1.200(8) Å for 5 and 1.1630(6) Å for 6), N-C distances (1.237(9) Å for 5 and 1.3188(7) for 6), and N-N-C angles  $(180.000(7)^{\circ} \text{ for } 5 \text{ and } 176.50(6)^{\circ} \text{ for } 6)$  that are notably similar to the corresponding parameters for the free diazoalkanes.<sup>11</sup> The Cu–N–N angle of 5 is linear (180.000(2)°) whereas for 6 the angle drops to 156.95(5)°.‡ For comparison, Hofmann's Cu<sup>III</sup>-diazoalkane complex has severely bent Cu-N-N and N-N-C angles (123.8(2) and 134.3(3)°, respectively) owing to the reduced diazoalkane ligand.<sup>10</sup>

Complex 5 is stable to 110 °C in toluene solutions even in the presence of Lewis acid catalysts. Complex 6 decomposes to a complicated mixture of products when heated in hydrocarbon solutions, and though the product profile simplifies when  $Sm(OTf)_3$  is added as a catalyst, no stable products were isolated from these thermolysis experiments, and no evidence for Cu=CR<sub>2</sub> bond formation was detected. The diazoalkane complex (dtbpe)Ni( $\eta^2$ -N<sub>2</sub>CPh<sub>2</sub>) has been isolated by Hillhouse and co-workers and was shown to extrude N<sub>2</sub> and form



Fig. 2 Solid-state structures of  $[Ph_2BP^{tBu}_2]Cu(N_2CMes_2)$  (6) (left) and  $[Ph_2BP^{tBu}_2]Cu-N(=CPh_2)NCPh_2$  (7) (right). Non-phosphorus atoms of the  $[Ph_2BP^{tBu}_2]$  ligands have been omitted for clarity.

(dtbpe)Ni=CPh<sub>2</sub> upon thermolysis in the presence of catalytic Sm(OTf)<sub>3</sub>.<sup>12</sup> Perhaps an important distinction to note between this Ni system and the Cu-diazoalkane adducts **5** and **6** is that N<sub>2</sub>CPh<sub>2</sub> binds in an  $\eta^2$ -NN mode to the L<sub>2</sub>Ni<sup>0</sup> fragment, whereas N<sub>2</sub>C(SiMe<sub>3</sub>)<sub>2</sub> and N<sub>2</sub>CMes<sub>2</sub> bind in an  $\eta^1$ -N fashion to the L<sub>2</sub>Cu<sup>I</sup> fragment described here.

When the less bulky diazoalkane reagent N2CPh2 was added to 3 or 4 at ambient temperature, N<sub>2</sub> release occurred spontaneously with concomitant formation of an inky blue solution and free NH<sub>2</sub>Mes (in the case of 3). A mixture of two Cucontaining species, ultimately assigned as [Ph<sub>2</sub>BP<sup>tBu</sup><sub>2</sub>]-Cu–N(=CPh<sub>2</sub>)NCPh<sub>2</sub> (7,  $\delta$  39.9) and [Ph<sub>2</sub>BP<sup>tBu</sup><sub>2</sub>]Cu=CPh<sub>2</sub> (8,  $\delta$  64.8), could be detected by <sup>31</sup>P{<sup>1</sup>H} NMR (Scheme 1). Quantitative conversion to 7 exclusively was achieved by the use of 2 equiv. or more of N<sub>2</sub>CPh<sub>2</sub>, and its assignment as a benzophenone azine adduct was verified by single-crystal Xray diffraction (Fig. 2).<sup>‡</sup> The N-N bond distance of 1.3962(3) A in 7 is consistent with the azine N–N single bond formulation. The Cu center in 7 is best described as trigonal planar, as the Cu– $N_{distal}$  distance (2.7649(3) Å) is much longer than the Cu-N<sub>proximal</sub> bond length of 2.0166(2) Å. Complex 7 presumably forms from the reaction between the intermediate carbene species 8 and unreacted N<sub>2</sub>CPh<sub>2</sub>; such C-N bond forming reactions between isolated terminal metal carbenes and diazoalkanes have been observed for metals other than copper.<sup>13</sup>

Under dilute conditions, as high as 70% conversion to carbene complex **8** has been achieved. This species features an intense band at  $\lambda_{max} = 583$  nm ( $\varepsilon = (est.)$  12 000 M<sup>-1</sup> cm<sup>-1</sup>) that likely arises from MLCT charge transfer into the Cu=*CPh*<sub>2</sub> unit. Warren has observed an optical transition of similar energy and intensity in the related  $\beta$ -diketiminato Cu=*CPh*<sub>2</sub> species.<sup>5</sup> To cement the assignment of **8**, solutions of [Ph<sub>2</sub>BP<sup>tBu</sup><sub>2</sub>]Cu=<sup>13</sup>CPh<sub>2</sub> (**8**-<sup>13</sup>C) were generated using N<sub>2</sub><sup>13</sup>CPh<sub>2</sub> in place of unlabelled diphenyldiazomethane. A diagnostic triplet at  $\delta$  331.5 was clearly visible in the <sup>13</sup>C{<sup>1</sup>H} NMR spectrum (<sup>2</sup>*J*<sub>PC</sub> = 41 Hz) (Fig. 3), and the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum (<sup>2</sup>*J*<sub>PC</sub> = 41 Hz). A slightly larger value for <sup>2</sup>*J*<sub>PC</sub> was observed for Hillhouse's (dtbpe)Ni=CPh<sub>2</sub> species ( $\delta$  222, <sup>2</sup>*J*<sub>PC</sub> = 51 Hz).<sup>12</sup> Interestingly, the Cu=*CP*h<sub>2</sub> <sup>13</sup>C{<sup>1</sup>H} NMR chemical shift in **8**-<sup>13</sup>C is considerably further downfield than for other reported copper carbene species,<sup>3-5</sup>



Fig. 3  $C{^{+}H}$  signal (left) and  ${^{-1}P}{^{+}H}$  signal (right) for  $[Ph_2BP^{tBu}_2]Cu={^{-1}3}CPh_2$  (8- ${^{13}C}$ ).

though it is certainly within the range typically observed for terminal metal carbene complexes.<sup>5,14</sup> As further evidence for the presence of the "CPh<sub>2</sub>" carbene functionality in **8**, we observed quantitative carbene transfer to CO to generate ketene Ph<sub>2</sub>C=CO and [Ph<sub>2</sub>BP<sup>tBu</sup><sub>2</sub>]Cu(CO) upon exposure of solutions of **8** to excess carbon monoxide (Scheme 1). Carbene transfer to CO is also characteristic of the Ni and Cu carbenes of Hillhouse and Warren, respectively.<sup>5,12</sup> Complex **8** did not readily transfer its carbene unit to olefins such as 1-hexene or styrene.

Whereas Warren's  $\beta$ -diketiminato copper carbene complex is stable at room temperature, complex **8** loses Ph<sub>2</sub>C==CPh<sub>2</sub> even at -30 °C. This fact has thus far precluded its crystallization from solution. The  $\beta$ -diketiminato Cu==CPh<sub>2</sub> species degrades similarly upon thermolysis.<sup>5</sup>

A computational study of 8 was undertaken to probe the Cu-C distance by DFT methods.<sup>15</sup> A diphenylcarbene unit was attached to the [Ph<sub>2</sub>BP<sup>tBu</sup><sub>2</sub>]Cu fragment taken from the atomic coordinates in the solid-state structure of 1. Geometry optimization calculations using various initial Cu-C distances all gave the same optimized geometry, featuring trigonal planar geometries at both the copper center and the carbene carbon atom, with a Cu-C distance of 1.933 A. Significantly shorter distances have been determined experimentally for the previously reported M=CR<sub>2</sub> (M = Cu, Ni) complexes (1.834-1.859 Å).<sup>5,12</sup> It is possible that the long predicted Cu-C distance in 8 is experimentally manifested by the relatively small value for  ${}^{2}J_{PC}$  and the significantly deshielded carbene carbon atom. It must however be noted that optimizing the geometry of 8 while fixing the Cu-C distance to be either 1.830 or 2.030 Å gave structures whose energies spanned a range of only 1.0 kcal  $mol^{-1}$ , indicating that the molecule's potential energy surface is quite flat with regard to the Cu-C distance.

This work was funded by BP ( $MC^2$  program) and the NSF (CHE-0132216). N. P. M. is grateful for an NSF Graduate Research Fellowship. J. Christopher Thomas is acknowledged for preliminary reaction screening. John Keith gave useful advice regarding DFT calculations, and Larry Henling provided crystallographic assistance. DFT calculations were done at the Materials and Process Simulation Center at the California Institute of Technology.

## Notes and references

‡ Details of the X-ray diffraction studies: 1,  $C_{18}H_{28}BCuNP_2$ : orthorhombic, *Fdd*2, a = 16.5848(13), b = 30.917(2), c = 13.3863(10) Å,

 $V = 6863.8(9) \text{ Å}^3$ , Z = 8,  $\mu$ (Mo-K $\alpha$ ) = 0.729 mm<sup>-1</sup>, 8664 total reflections, 2460 independent ( $R_{int} = 0.0409$ ) with  $I > 2\sigma(I)$ ; **2**, C<sub>60</sub>H<sub>106</sub>BCuLiNO<sub>8</sub>P<sub>2</sub>: monoclinic, P2<sub>1</sub>/c, a = 11.9146(9), b = 21.3858(16), c = 24.4619(18) Å,  $\beta = .92.5560(10)^\circ$ , V = .0406 $A^3, Z = 8, \mu(Mo-K\alpha) = 0.905 \text{ mm}^{-1}, 72625 \text{ total reflections}, 14800 \text{ independent} (R_{int} = 0.0841) \text{ with } I > 2\sigma(I);$ **3**,  $C_{39}H_{63}BCuNP_2$ : monoclinic,  $P2_1/c$ , a = 10.186(5), b = 17.079(8), 22.083(7) Å,  $\beta = 98.24(4)^{\circ}$ , V = 3802(3) Å<sup>3</sup>, Z = 4,  $\mu$ (Mo-K $\alpha$ ) =  $0.685 \text{ mm}^{-1}$ , 31590 total reflections, 7306 independent ( $R_{\text{int}} = 0.0990$ ) with  $I > 2\sigma(I)$ ; 4, C<sub>34</sub>H<sub>58</sub>BCuOP<sub>2</sub>: orthorhombic, *Fdd2*, *a* 17.548(3), b = 30.333(5), c = 12.818(3) Å, V = 6823(2) Å<sup>3</sup>, Z = 8,  $\mu$ (Mo-K $\alpha$ ) = 0.758 mm<sup>-1</sup>, 3639 total reflections, 2005 independent  $(R_{\text{int}} = 0.0385)$  with  $I > 2\sigma(I)$ ; **5**,  $C_{34}H_{59}BCuN_2P_2Si_2$ ; trigonal,  $R\bar{3}c$ , a = 21.4508(13), c = 47.633(3) Å, V = 18981(2) Å<sup>3</sup>, Z = 15,  $\mu$ (Mo-K $\alpha$ ) = 0.561 mm<sup>-1</sup>, 37667 total reflections, 3067 independent ( $R_{int}$  = 0.0834) with  $I > 2\sigma(I)$ ; 6, C<sub>49</sub>H<sub>72</sub>BCuN<sub>2</sub>P<sub>2</sub>: monoclinic,  $P2_1/c$ , a =11.6975(16), b = 17.955(2), c = 23.035(3) Å,  $\beta = 91.707(4)^\circ$ , V = 4836.1(12) Å<sup>3</sup>, Z = 4,  $\mu$ (Mo-K $\alpha$ ) = 0.550 mm<sup>-1</sup>, 22763 total reflections, 6798 independent ( $R_{\rm int}$  = 0.1309) with  $I > 2\sigma(I)$ . 7,  $C_{56}H_{70}BCuN_2P_2$ : monoclinic,  $P2_1/c$ , a = 12.0749(11), b $Z_{2,347(2)}$ , c = 19.2412(18) Å,  $β = 104.625(2)^{\circ}$ , V = 5023.8(8) Å<sup>3</sup>, Z = 4, μ(Mo-Kα) = 0.536 mm<sup>-1</sup>, 42567 total reflections, 11402 independent ( $R_{int} = 0.0617$ ) with  $I > 2\sigma(I)$ . CCDC 659790–659796. For crystallographic data in CIF or other electronic format see DOI: 10.1039/b713687k

- Selected reviews: W. Kirmse, Angew. Chem., Int. Ed., 2003, 42, 1088; H. M. L. Davies and S. A. Panaro, Tetrahedron, 2000, 56, 4871; M. P. Doyle and D. C. Forbes, Chem. Rev., 1998, 98, 911.
- 2 For recent advances see, for example: M. R. Fructos, T. R. Belderrain, M. C. Nicasio, S. P. Nolan, H. Kaur, M. M. Díaz-Requejo and P. J. Pérez, *J. Am. Chem. Soc.*, 2004, **126**, 10846, and references therein.
- 3 Selected examples: J. Barluenga, L. A. López, O. Löber, M. Tomás, S. García-Granda, C. Alvarez-Rúa and J. Borge, *Angew. Chem., Int. Ed.*, 2001, **126**, 10085; D. S. Laitar, P. Müller and J. P. Sadighi, *J. Am. Chem. Soc.*, 2005, **127**, 17196; H. Kaur, F. Kauer, E. D. Stevens and S. P. Nolan, *Organometallics*, 2004, **23**, 1157.
- 4 B. Straub and P. Hofmann, Angew. Chem., Int. Ed., 2001, 40, 1288.
- 5 Y. M. Badiei and T. H. Warren, J. Organomet. Chem., 2005, 690, 5989; X. Dai and T. H. Warren, J. Am. Chem. Soc., 2004, 126, 10085.
- Selected examples: D. M. Jenkins, T. A. Betley and J. C. Peters, J. Am. Chem. Soc., 2002, 124, 11238; S. D. Brown, T. A. Betley and J. C. Peters, J. Am. Chem. Soc., 2003, 125, 322; T. A. Betley and J. C. Peters, J. Am. Chem. Soc., 2004, 126, 6252; C. E. MacBeth, J. C. Thomas, T. A. Betley and J. C. Peters, Inorg. Chem., 2004, 43, 4645.
- 7 J. C. Thomas and J. C. Peters, J. Am. Chem. Soc., 2003, 125, 8870; J. C. Thomas and J. C. Peters, *Inorg. Chem.*, 2003, 42, 5055.
- 8 J. C. Thomas and J. C. Peters, Polyhedron, 2004, 23, 2901.
- 9 E. D. Blue, A. Davis, D. Conner, T. B. Gunnoe, P. D. Boyle and P. S. White, J. Am. Chem. Soc., 2003, 125, 9435.
- 10 B. F. Straub, F. Rominger and P. Hofmann, Organometallics, 2000, 19, 4305.
- E. Glozbach and J. Lorberth, J. Organomet. Chem., 1980, 191, 371;
  H. E. Zimmerman and D. H. Paskovich, J. Am. Chem. Soc., 1964, 86, 2149.
- 12 R. Waterman and G. L. Hillhouse, J. Am. Chem. Soc., 2003, 125, 13350; D. J. Mindiola and G. L. Hillhouse, J. Am. Chem. Soc., 2002, 124, 9976.
- 13 H. Zhu, J. Chai, A. Stasch, H. W. Roesky, T. Blunk, D. Vidovic, J. Magull, H.-G. Schmidt and M. Noltemeyer, *Eur. J. Inorg. Chem.*, 2004, 4046.
- 14 R. H. Crabtree, The Organometallic Chemistry of the Transition Metals, Wiley-Interscience, New York, 2001.
- 15 B3LYP functional; LACVP\*\* basis set for Cu atoms, 6-31G\*\* basis set for carbene C and P atoms; MIDI! basis set for all other atoms. See ESI† for additional details.