## Intramolecular alkene hydroaminations catalyzed by a bis(thiophosphinic amidate) Zr(IV) complex<sup>†</sup>

Hyunseok Kim,<sup>a</sup> Phil Ho Lee<sup>\*a</sup> and Tom Livinghouse<sup>\*b</sup>

Received (in Berkeley, CA, USA) 22nd April 2005, Accepted 15th July 2005 First published as an Advance Article on the web 20th September 2005 DOI: 10.1039/b505738h

A neutral Zr(TV) complex has been shown to be an effective precatalyst for intramolecular alkene hydroaminations that provide cyclic amines in good to excellent yields.

The intramolecular hydroamination of alkenes constitutes a powerful method for the synthesis of azacycles.<sup>1</sup> The seminal group 3 metallocenes developed by Marks and co-workers<sup>2</sup> have recently been joined by a variety of non-metallocene complexes of the group 3 metals as catalysts for this important reaction.<sup>3,4</sup> Although there are numerous accounts that describe both the intra- and intermolecular hydroamination of alkynes5a-d and allenes<sup>5e,f</sup> catalyzed by various complexes of the group 4 metals, there have been very few reports of intramolecular alkene hydroaminations mediated by complexes of metals belonging to this triad, and all but one<sup>6</sup> of these are restricted to cationic species.<sup>7</sup> We have previously disclosed that chelating bis(thiophosphinic amidate) "NPS" complexes of the group 3 metals (particularly Y and Nd) are potent catalysts for intramolecular alkene hydroamination.<sup>3d</sup> In this communication, we show that the neutral Zr(IV) bis(thiophosphinic amidate) complex (1) is a competent precatalyst for the cyclization of representative primary aminoalkenes (Scheme 1).



Scheme 1

The internal hydroamination of 2,2-dimethylpent-4-ene-1-amine (4a) was selected for initial examination as it was expected that

<sup>b</sup>Department of Chemistry, Montana State University, Bozeman, MT 59717, USA. E-mail: livinghouse@chemistry.montana.edu
† Electronic supplementary information (ESI) available: Experimental procedures and characterization data for all new compounds. See http:// dx.doi.org/10.1039/b505738h

cyclization of this substrate would be facilitated by a geminal Thorpe-Ingold effect.<sup>3f,6</sup> Successful generation of the desired NPS precatalyst 1 proved experimentally straightforward via amine elimination, involving commercially available Zr(NMe2)4 (2) and bis(thiophosphinic amide) (3)<sup>3d</sup> (1 equiv., C<sub>6</sub>D<sub>6</sub>, 25 °C, 10 min). The <sup>1</sup>H, <sup>31</sup>P and <sup>13</sup>C NMR spectra of 1 are consistent with a monomeric species possessing an octahedral structure with a mer configuration. The NMe2 resonance (500 MHz) appears as a sharp singlet at 3.11 ppm and the linker CH<sub>2</sub> as a doublet (2.69 ppm, J = 10 Hz). The signal for the CH adjacent to P appears as a welldefined septet centered at 2.00 ppm (J = 7 Hz), with the diastereomeric isopropyl methyls appearing as a set of doublets between 1.16 and 1.10 ppm (J = 7 Hz). The <sup>31</sup>P NMR spectrum of 1 reveals a singlet at 75.10 ppm.<sup>8</sup> The thermal stability of 1 was demonstrated by heating at 150 °C for 19 h, whereupon no alteration to the NMR spectra was observed. Addition of 4a to 1 (5 mol%), followed by heating at 100 °C, provided the pyrrolidine 5a in 97% yield<sup>9</sup> after 105 h. Alternatively, cyclization of 4a at 120 °C (C<sub>6</sub>D<sub>6</sub>) and 150 °C (toluene-d<sub>8</sub>) provided 5a in 94% (12 h) and 98% (2.5 h) yield, respectively. Closely-related reaction conditions were subsequently utilized for the cyclization of a series of representative primary amines 4b-f and, unsuccessfully, for the secondary amine 4g. A compilation of the reaction times and yields observed for the cyclization of aminoalkenes 4a-g in the presence of the Zr(IV)·NPS chelate (1) appears in Table 1. Significantly, cyclization of 4a, 4b and 4d on a preparative (3 mmol) scale followed by separation of the products from the catalyst by vacuum transfer and protonation (HCl-MeOH), furnished 5a, 5b and 5d as their hydrochloride salts in 90, 88 and 85% isolated yields, respectively.

Several of the trends that emerge from the foregoing examples are worthy of comment. The presence of geminal Thorpe-Ingold buttressing is helpful but not a prerequisite for successful cyclization, as 5-hexen-2-amine 4b and 4-penten-1-amine 4c partake in the reaction. Accordingly, 4c was smoothly converted to 5c in 91% yield in the presence of 1 (10 mol%) in 10 h at 150 °C. By way of comparison, cyclization of 4c using  $Zr(NMe_2)_4$  as the precatalyst (10 mol%, 150 °C, toluene-d<sub>8</sub>) gave 5c (91%) but required 28 h. In addition, the styrenyl substrate 4d, containing an internal alkene, underwent cyclization at 150 °C to provide 5d in 93% yield after 120 h. In this instance, the reaction time could be shortened to 39 h when 10 mol% of the precatalyst was employed. A similar result was observed in the case of the 1,1-disubstituted aminoalkene 4f. That the latter substrate undergoes cyclization to give the product derived from exocyclic addition to the alkene is consistent with the mechanism shown in Scheme 2. It is also of significance that the secondary

<sup>&</sup>lt;sup>a</sup>Department of Chemistry, Kangwon National University, Chunchon, 200-701, Republic of Korea

## **Table 1** Internal alkene hydroaminations catalyzed by $\mathbf{1}^a$

Entry	Substrate	Product	Temp./°C	Time/h	Yield $(\%)^b$
1	NH <sub>2</sub> 4a	NH 5a	120 150	12 2.5	94 98
2	NH <sub>2</sub> 4b	NH 5b	120 150	172 22	98 <sup>c</sup> 96 <sup>c</sup>
3	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	<b>NH</b> 5c	$\frac{120^d}{150^d}$	41 10	89 91
4	NH <sub>2</sub> Ph 4d	NH 5d	$150 \\ 150^d$	120 39	93 91
5	NH <sub>2</sub> 4e	NH 5e	120 150	9 1	99 99
6	NH <sub>2</sub> 4f	Sf	150 150 <sup>d</sup>	104 45	92 94
7	NH 4g	<br 5g	150	18	0

<sup>*a*</sup> Benzene-d<sub>6</sub> (120 °C) or toluene-d<sub>8</sub> (150 °C) were used as solvents, respectively. <sup>*b*</sup> NMR yields based on *p*-xylene as an internal standard. <sup>*c*</sup> trans : cis = 1.3 : 1.0. <sup>*d*</sup> 10 mol% Catalyst was used.



Seneme 4

aminoalkene **4g** is resistant to cyclization. This stands in sharp contrast to the results of Scott and Hultzsch who have reported that secondary, but not primary, aminoalkenes participate in internal hydroamination, catalyzed by cationic Zr(IV) complexes.<sup>7*a,b*</sup> Finally, the Zr(IV)·NPS complex **1** shows higher activity as a precatalyst than Zr(NMe<sub>2</sub>)<sub>4</sub> **2**.

The dynamics of the hydroamination involving precatalyst 1 can be conveniently monitored by <sup>31</sup>P NMR. Addition of 4a to a C<sub>6</sub>D<sub>6</sub> solution of 1 (5 mol%) results in the immediate disappearance of the phosphorus resonance at 75.10 ppm and the concomitant appearance of a new signal at 78.12 ppm. That this is accompanied by the production of 2 equiv. of Me<sub>2</sub>NH (<sup>1</sup>H NMR) is strongly indicative of a quantitative exchange of the amido ligands at zirconium, resulting in the incorporation of two substrate aminoalkene substituents. Significantly, cyclization of 4a at 120 °C over 12 h results in a 92% conversion to 5a with no change to the <sup>31</sup>P resonance at 78.12 ppm, thus providing evidence that the Zr catalyst is robust under the reaction conditions. In addition, at no time during this reaction did the 31P resonance associated with the free proligand 3 at 89.54 ppm appear. A probable mechanistic pathway for the intramolecular hydroamination of 4a, involving the putative Zr(IV) imido complex  $A^{10}$  and azazirconacyclobutane B based on these observations, is depicted in Scheme 2.

In conclusion, we have shown that the neutral Zr(IV)·NPS complex (1) is a competent precatalyst for intramolecular alkene hydroaminations involving primary amines. Although the catalytic activity of 1 is lower than that exhibited by the corresponding Y(III)·NPS chelate,<sup>3d</sup> the results presented here are among the first examples of internal alkene hydroamination catalyzed by a neutral complex of a group 4 metal.<sup>6</sup> Studies utilizing cationic Zr(IV)·NPS and related complexes for this process are currently under way.

Generous financial support for this research was provided by the National Institutes of Health and the National Science Foundation.

## Notes and references

- 1 (a) F. Pohlki and S. Doye, *Chem. Soc. Rev.*, 2003, **32**, 104; (b) T. E. Muller and M. Beller, *Chem. Rev.*, 1998, **98**, 675.
- 2 (a) S. Hong and T. J. Marks, Acc. Chem. Res., 2004, 39, 673 and references therein; (b) J. -S. Ryu, T. J. Marks and F. E. McDonald, J. Org. Chem., 2004, 69, 1038; (c) M. R. Gagné, C. L. Stern and T. J. Marks, J. Am. Chem. Soc., 1992, 114, 275; (d) S. Tian, V. M. Arredondo, C. L. Stern and T. J. Marks, Organometallics, 1999, 18, 4421; (e) J. -S. Ryu, T. J. Marks and F. E. McDonald, Org. Lett., 2001, 3, 3091; (f) V. M. Arredondo, S. Tian, F. E. McDonald and T. J. Marks, J. Am. Chem. Soc., 1999, 121, 3633 and references therein.
- 3 (a) J. -S. Ryu, G. Y. Li and T. J. Marks, J. Am. Chem. Soc., 2003, 125, 12584; (b) P. N. O'Shaughnessy, P. D. Knight, C. Morton, K. M. Gillespie and P. Scott, Chem. Commun., 2003, 1770; (c) D. V. Gribkov, K. C. Hultzsch, F. Hampel and T. Wagner, Organometallics, 2004, 23, 2601; (d) Y. K. Kim, T. Livinghouse and Y. Horino, J. Am. Chem. Soc., 2003, 125, 9560; (e) Y. K. Kim and T. Livinghouse, Angew. Chem., Int. Ed., 2002, 41, 3645; (f) Y. K. Kim, T. Livinghouse and J. E. Bercaw, Tetrahedron Lett., 2001, 42, 2933; (g) D. V. Gribkov, K. C. Hulzsch and F. Hampel, Chem. Eur. J., 2003, 9, 4796; (h) Z. Zhang and L. L. Schafer, Org. Lett., 2003, 5, 4733.
- 4 (a) Catalytic activity has also been observed for cationic scandium non-metallocenes: F. Lauterwasser, P. G. Hayes, S. Brase, W. Piers and L. L. Schafer, *Organometallics*, 2004, 23, 2234; as well as calcium derivatives: (b) M. R. Crimmin, I. J. Casely and M. S. Hill, *J. Am. Chem. Soc.*, 2005, 127, 2042.
- 5 (a) A. L. Odom, *Dalton Trans.*, 2005, 225 and references therein; (b) S. Doye, *Synlett*, 2004, 1653; (c) J. M. Hoover, J. R. Petersen, J. H. Pikul and A. R. Johnson, *Organometallics*, 2004, 23, 4614; (d) C. Li, R. K. Thomson, B. Gillon, B. O. Patrick and L. L. Schafer,

Chem. Commun., 2003, 2462; (e) L. Ackermann and R. G. Bergman, Org. Lett., 2002, 4, 1475; (f) Y. Shi, J. T. Ciszewski and A. L. Odom, Organometallics, 2001, 20, 3967.

- 6 J. A. Bexrud, J. D. Beard, D. C. Leitch and L. L. Schafer, Org. Lett., 2005, 7, 1959.
- 7 (a) P. D. Knight, I. Munslow, P. N. O'Shaughnessy and P. Scott, *Chem. Commun.*, 2004, 894; (b) D. V. Gribkov and K. C. Hultzsch, *Angew. Chem., Int. Ed.*, 2004, **43**, 5542.
- $8~^1H$  NMR,  $^{13}C$  NMR and  $^{31}P$  NMR of NPS and NPS-Zr(NMe\_2)\_2 are available as ESI.†.
- 9 J. Young NMR tubes, purchased from Aldrich or J. Young Ltd, were used under refluxing conditions (bath temperature, 120 °C or 150 °C) without any special precautions.
- (a) Y. Li, Y. Shi and A. L. Odom, J. Am. Chem. Soc., 2004, 126, 1794;
   (b) B. D. Ward, A. Maisse-Francois, P. Mountford and L. H. Gade, Chem. Commun., 2004, 704.