Palladium-Catalyzed Silylation of Electron-Deficient Aryl Iodides Using Triorganosilane in the Presence of Pyridine and LiCl

Muneaki Iizuka^[a] and Yoshinori Kondo^{*[a]}

Keywords: Silylation / Palladium / Aryl iodides / Pyridine / Lithium chloride

Palladium-catalysed silylation of aryl iodides with electronwithdrawing groups was efficiently achieved using pyridine and lithium chloride as additives and conducting the reaction at room temperature. Functionalized aryl[2-(hydroxymethyl)phenyl]dimethylsilanes were also prepared by palladiumcatalysed reaction using THP-protected [2-(hydroxymethyl)-

Introduction

The silicon-based cross coupling has become an important method for carbon-carbon bond formation, and the reaction is now widely used (Hiyama coupling).^[1] Various arylsilanes have been employed for the coupling partners and the transition-metal-catalysed silvlation of aryl halides should be one of the most versatile and chemoselective methods to prepare arylsilanes.^[2] Recently, Yamanoi reported the palladium-catalysed silvlation of aryl halides using *t*Bu₃P as a ligand (Figure 1).^[2h] Trialkylsilyl groups can be successfully transferred by the reaction; however, the reaction is not applicable to aryl halides with an electronwithdrawing group. Murata et al. also reported a palladium-catalysed silvlation of aryl halides with electron-donating groups using triorganosilanes.^[2e] Alami et al. reported a platinum oxide catalysed trialkylsilylation of various aryl halides,^[2f] but there has been no successful report of palladium-catalysed silvlations of aryl halides with electron-withdrawing groups.



Figure 1. Side reaction in the silylation of iodoarenes.^[2h]

Fax: +81-22-795-6804

WWW under http://www.eurjoc.org or from the author.

phenyl]dimethylsilane as a silylating agent followed by deprotection. Rhodium-catalysed 1,4-addition of the arylsilane was carried out using cyclohexenone as an enone in excellent yield.

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2008)

In connection with our recent study on the selective transformation of arylsilanes,^[3] we investigated the selective silylation of aryl halides with various functional groups catalysed by palladium.

Results and Discussion

We first investigated the palladium-catalysed silylation of 4-iodonitrobenzene (1), which was considered to be the most challenging substrate for the selective silylation as shown in Figure 1. In order to avoid the formation of reduced product 3, various reaction conditions were screened, and conditions free of phosphane ligands were suggested to be favorable. After the screening of various reaction conditions, we found that the use of pyridine and lithium chloride as additives is effective for selective silylation.

When 1-iodo-4-nitrobenzene (1) was treated with Et₃SiH in the presence of Pd(OAc)₂ (4 mol-%), pyridine (2.5 equiv.) and LiCl (4 equiv.) in DMI (N,N-dimethylimidazolin-2one), silvlated nitrobenzene 2 was obtained in 51% yield, and the ¹H NMR analysis of the crude material indicated a ratio of 2/3 = 8:2 (Table 1, Entry 1). The use of other solvents such as NMP or DMF gave poorer 2/3 ratios (Table 1, Entries 2, 3). When Et₃N was employed instead of pyridine, no formation of 2 was observed, and only 3 was produced (Table 1, Entry 4). The absence of base also resulted in no formation of 2 (Table 1, Entry 5). The use of NaOAc, LiBr, or LiI as an additive was not effective for the formation of 2, and only 3 was formed (Table 1, Entries 6-8). On the other hand, the use of Et_4NCl was effective like LiCl, and 2 was obtained in 56% yield (Table 1, Entry 9). When the amount of LiCl was reduced to 1 equiv., the ratio of 2/3 decreased (Table 1, Entry 10). Thus, pyridine was found to be essential for selective formation of 2, and also chloride anions play an important role for the selective transfer of a silvl group.

 [[]a] Graduate School of Pharmaceutical Sciences, Tohoku University, Aramaki Aza Aoba 6-3, Sendai 980-8578, Japan

E-mail: ykondo@mail.pharm.tohoku.ac.jp Supporting information for this article is available on the

SHORT COMMUNICATION

Table 1. Palladium-catalyzed silylation of 1-iodo-4-nitrobenzene.

0 ₂ N 1	base	iH, Pd(OAc) ₂ (4 e, additive, solve r.t., 24 h	ent O	2N 2	SIEt ₃ + O ₂ N 3
Entry	Base	Additive	Sovent	2/3 ^[a]	Yield (%) of 2 ^[b]
1	pyridine	LiCI (4 equiv.)	DMI	8:2	52
2	pyridine	LiCI (4 equiv.)	NMP	7:3	-
3	pyridine	LiCI (4 equiv.)	DMF	6:4	_
4	Et₃N	LiCI (4 equiv.)	DMI	0:10	_
5	-	LiCI (4 equiv.)	DMI	0:10	-
6	pyridine	NaOAc	DMI	0:10	-
7	pyridine	LiBr	DMI	0:10	_
8	pyridine	Lil	DMI	0:10	_
9	pyridine	Et ₄ NCI	DMI	8:2	56
10	pyridine	LiCI (1 equiv.)	DMI	6:4	-

[a] Estimated by ¹H NMR spectroscopy. [b] Isolated yields.

Further investigations were carried out for the silvlation of various aryl iodides using several silanes. On 1-iodo-4nitrobenzene (1), the use of PhMe₂SiH gave almost similar results with Et₃SiH, and A (2b) was isolated in 45% yield (Table 2, Entry 1). When Ph₃SiH was used, the yield of A (2c) was improved to 70% (Table 2, Entry 2). The reaction of 4-iodobenzonitrile (5a) with Et₃SiH gave A (6a) in 46% yield (Table 2, Entry 3). The use of PhMe₂SiH improved the yield of A (6b) up to 94%, and no formation of B was observed (Table 2, Entry 4). The reaction using Ph₃SiH gave A (6c) in 64% yield with no formation of B (Table 2, Entry 5). The reaction of 4-iodoacetophenone (5b) with Et₃SiH gave a 1:1 mixture of A/B, and the desired A (7a) was isolated in 38% yield (Table 2, Entry 6). When PhMe₂SiH was used, the ratio of A/B was improved to 8:2, and A (7b) was obtained in 73% yield (Table 2, Entry 7). The use of Ph₃SiH gave the best result, and A (7c) was obtained in 85% yield with no formation of B (Table 2, Entry 8). The reaction of methyl 4-iodobenzoate (5c) with Et₃SiH gave a 6:4 mixture of A/B, and A (8a) was isolated in 35% yiled (Table 2, Entry 9). The use of PhMe₂SiH gave a 9:1 mixture of A/B, and A (8b) was isolated in 82% yield (Table 2, Entry 10). The use of Ph₃SiH showed an excellent selectivity, and A (8c) was obtained in 91% yield with no formation of B (Table 2, Entry 11). 3-Iodopyridine (5d) was also treated with PhMe₂-SiH, and the corresponding silvlated pyiridine 9 was obtained in 87% yield (Table 2, Entry 12). The use of phenylsubstituted silanes showed better selectivities, and this tendency was also suggested by Murata's report.^[2e] At this moment, the silvlation of an electron-rich aryl halide such as 4iodoanisole was unsuccessful under our reaction conditions, and only a trace amount of silylated product was isolated.

With respect to the mechanism of the silylation reaction, two types of pathways have been suggested: (i) σ -bond metathesis, (ii) Pd^{IV} intermediate involvement.^[2a,2g] At this moment, it is not clear how pyridine and LiCl effectively form the silylated products; however, the combination is suggested to be favorable for the transition state of the metathesis or the formation of Pd^{IV} species. Table 2. Palladium-catalyzed silylation of iodoarenes.

R X	, 5	H, Pd /ridine r.t.,	(OAc) ₂ (4 s, LiCl, Di 24 h	4 mol-%) MI R	A Si	Et ₃ + R B
Entry	R	х	1 or 5	R₃SiH	A/B ^[a]	Yield (%) of $\mathbf{A}^{[b]}$
1	NO ₂	СН	1	PhMe ₂ SiH	9:1	45 (2b)
2		СН	1	Ph₃SiH	_	70 (2c) ^[c]
3	CN	СН	5a	Et ₃ SiH	_	46 (6a)
4	CN	СН	5a	PhMe ₂ SiH	10:0	94 (6b)
5	CN	СН	5a	Ph₃SiH	10:0	64 (6c)
6	COMe	СН	5b	Et₃SiH	5:5	38 (7a)
7	COMe	СН	5b	PhMe ₂ SiH	8:2	73 (7b)
8	COMe	СН	5b	Ph₃SiH	10:0	85 (7c)
9	COOMe	СН	5c	Et ₃ SiH	4:6	35 (8a)
10	COOMe	СН	5c	PhMe ₂ SiH	9:1	82 (8b)
11	COOMe	СН	5c	Ph₃SiH	10:0	91 (8c)
12	Н	Ν	5d	PhMe ₂ SiH	-	87 (9)

[a] Estimated by ¹H NMR spectroscopy. [b] Isolated yields. [c] DMF was used as a solvent.

Our next interest was focused on the formation of aryl[2-(hydroxymethyl)phenyl]dimethylsilanes, which recently were reported to be excellent aryl donors for various transition-metal-catalysed reactions.^[4] The THP-protected [2-(hydroxymethyl)phenyl]dimethylsilane was treated with 4-iodo-benzonitrile (**5a**) and 4-iodoacetophenone (**5b**) followed by the treatment with *p*-TsOH to remove THP protective group, and the silylated arenes **10** and **11** were obtained in 59% and 58% yields, respectively. The [2-(hydroxymethyl)phenyl]silylated acetophenone **11** was then treated with cyclohexenone in the presence of a rhodium catalyst which gave the 1,4-adduct **12** in 96% yield (Scheme 1).



Scheme 1. Selective transformation of arylsilanes.

Conclusions

Palladium-catalysed silylation of aryl iodides with electron-withdrawing groups was efficiently achieved using pyridine and lithium chloride as additives and conducting the reaction at room temperature. Aryl[2-(hydroxymethyl)phenyl]dimethylsilanes were also prepared by palladiumcatalysed reaction using protected [2-(hydroxymethyl)phenyl]dimethylsilane as a silylating agent followed by de-



protection. Rhodium-catalysed 1,4-addition of the arylsilane was carried out using cyclohexenone in excellent yield. Further investigations on the scope and limitations of the palladium-catalysed silylation and mechanistic studies are underway.

Supporting Information (see footnote on the first page of this article): Experimental procedure and spectroscopic data of synthesized compounds.

Acknowledgments

This work was partly supported by the Grant-in-Aid for Scientific Research on Priority Areas "Advanced Molecular Transformations of Carbon Resources" (No. 19020005) and "Synergistic Effects for Creation of Functional Molecules" (No. 19027008) and other Research Grants (No. 19390002 and No. 18659001) from the Ministry of Education, Science, Sports and Culture, Japan and a grant from Yamada Science Foundation. We thank Professor Miki Murata for his kind suggestions and discussions on this work.

219, p. 61; c) K. A. Horn, *Chem. Rev.* **1995**, *95*, 1317; d) Y. Hatanaka, T. Hiyama, *Synlett* **1991**, 845.

- [2] a) M. Murata, H. Yamasaki, K. Uogishi, S. Watanabe, Y. Masuda, Synthesis 2007, 2944; b) Y. Yamanoi, T. Taira, J. Sato, I. Nakamula, H. Hishihara, Org. Lett. 2007, 9, 4543; c) M. Murata, K. Oka, H. Yamasaki, S. Watanabe, Y. Masuda, Synlett 2007, 1387; d) M. Murata, H. Yamasaki, T. Ueta, M. Nagata, M. Ishikura, S. Watanabe, Y. Masuda, Tetrahedron 2007, 63, 4087; e) M. Murata, H. Ohara, R. Oiwa, S. Watanabe, Y. Masuda, Synthesis 2006, 1771; f) A. Hamze, O. Provot, M. Alami, J.-D. Brion, Org. Lett. 2006, 8, 931; g) D. Karchtedt, A. T. Bell, T. D. Tilley, Organometallics 2006, 25, 4471; h) Y. Yamanoi, J. Org. Chem. 2005, 70, 9607; i) S. E. Denmark, J. M. Kallemeyn, Org. Lett. 2003, 5, 3483; j) W. Gu, S. Liu, R. B. Silverman, Org. Lett. 2002, 4, 4171; k) A. S. Manoso, P. J. DeShong, J. Org. Chem. 2001, 66, 7455; l) M. Murata, K. Suzuki, S. Watanabe, Y. Masuda, Tetrahedron Lett. 1999, 40, 9255; m) M. Murata, K. Suzuki, S. Watanabe, Y. Masuda, J. Org. Chem. 1997, 62, 8569.
- [3] K. Suzawa, M. Ueno, A. E. H. Wheatley, Y. Kondo, *Chem. Commun.* **2006**, 4850.
- [4] a) Y. Nakao, H. Imanaka, A. H. Sahoo, A. Yada, T. Hiyama, J. Am. Chem. Soc. 2005, 127, 6952; b) Y. Nakao, J. Chen, H. Imanaka, T. Hiyama, Y. Ichikawa, W.-L. Duan, R. Shintani, T. Hayashi, J. Am. Chem. Soc. 2007, 129, 9137; c) Y. Nakao, J. Chen, M. Tanaka, T. Hiyama, J. Am. Chem. Soc. 2007, 129, 11694.

Received: December 2, 2007 Published Online: January 29, 2008

a) T. Hiyama, in *Metal-Catalyzed Cross-Coupling Reactions* (Eds.: F. Diederich, P. J. Stang), Wiley-VCH, Weinheim, **1998**, p. 421; b) T. Hiyama, E. Shirakawa, in *Topics in Current Chemistry* (Ed.: N. Miyaura), Springer-Verlag, Heidelberg, **2002**, vol.