

Ternary chlorides in the systems $ACl/YbCl_3$ ($A=Cs,Rb,K$)

Jörg Sebastian, Hans-Joachim Seifert*

FB19 Inorganic Chemistry, University Gh Kassel, Heinrich-Plett-Str. 40, D-34109 Kassel, Germany

Received 9 June 1997; accepted 15 October 1997

Abstract

The phase diagrams of the $ACl/YbCl_3$ ($A=Cs,Rb,K$) systems were investigated by DTA and XRD. Compounds A_3YbCl_6 , A_2YbCl_5 and AYb_2Cl_7 exist in all systems, enneachlorides $A_3Yb_2Cl_9$ in the systems with $A=Cs,K$. In the group AYb_2Cl_7 , the Yb ions have coordination number 7, in all other compounds the coordination is octahedral. Thermodynamic functions determined by solution calorimetry and emf vs T measurements in galvanic cells for solid electrolytes reveal that the ternary chlorides A_nYbCl_{3+n} are stable compared with mixtures ($nACl+YbCl_3$); the compounds Cs_2YbCl_5 and Rb_2YbCl_5 are stable only at temperatures >0 K. © 1998 Elsevier Science B.V.

Keywords: Phase diagrams; Ternary ytterbium chlorides; Thermodynamic properties

1. Introduction

In connection with our research work on ternary lanthanide chlorides, we have investigated the pseudo-binary systems of ytterbium(III) chloride with the alkali metal chlorides CsCl, RbCl and KCl. Hitherto, DTA measurements on the $KCl/YbCl_3$ [1] and $CsCl/YbCl_3$ systems [2] were performed and the unit cells of those compounds, which could be prepared directly from melts, were determined by Meyer [3] with X-ray powder patterns.

Our special interests were directed to the question of whether systematic rules found recently from investigations of the analogous systems with $HoCl_3$ [4], $ErCl_3$ [5] and $TmCl_3$ [6] are also valid for ytterbium compounds. In addition to DTA and XRD, we have measured solution enthalpies and (free) Gibbs' enthalpies with galvanic cells for solid electrolytes –

methods which have proved very successful for yielding predictions about the mutual stabilities of all ternary chlorides in the respective systems.

2. Experimental

2.1. Chemicals

The starting compound was $YbCl_3 \cdot 6H_2O$, prepared from a solution of Yb_2O_3 (99.9%, Heraeus/Hanau) in hot hydrochloric acid. Alkali metal chlorides were dried at $500^\circ C$.

2.2. Differential thermal analysis (DTA)

The DTA measurements were performed in a home-built device for samples (~ 0.5 g) in vacuum-sealed silica ampoules. Samples with <50 mol% $YbCl_3$ were homogenized by melting in a gas flame, shaking and

*Corresponding author. Fax: +49 561 804 4010.

annealing. Because YbCl_3 melts react with silica, YbCl_3 -rich mixtures were melted in the DTA furnace ≈ 20 K above the alleged liquidus temperatures. In general, heating curves were recorded (heating rate = 2 K min^{-1}).

2.2.1. X-ray powder patterns

Powder patterns at ambient temperature were taken with a Philips PW 1050/25 goniometer equipped with a proportional counter and a vacuum attachment. During exposure (CuK_α radiation) the samples were kept under He atmosphere. For high-temperature photographs, a Simon–Guinier camera (Enraf–Nonius) was applied. Corundum powder was used as the internal standard; at 20°C : $a=475.92 \text{ pm}$ and $c=1299.00 \text{ pm}$; at 500°C , $a=479.32$ and $c=1308.92 \text{ pm}$.

2.2.2. Solution calorimetry


The heats of solution were measured using an isoperibolic underwater calorimeter first described in 1978 [7]. In a silver vessel of 1.3 l volume, samples of 2 to 4 g in thin-walled glass ampoules are cracked under water at the beginning of the measurement. The change of temperature, T , against the surroundings, in a thermostat with a temperature constancy of $2\text{--}10^{-4} \text{ K}$, generated by the solution enthalpy is measured with a thermopile. The calibration is done by Joule heating.

At least two samples of each substance were measured. From the enthalpies of solution, $\Delta_{\text{sol}}H^0_{298}$, the enthalpies of formation of the ternary chlorides from ($n\text{AlCl} + \text{YbCl}_3$), $\Delta_f H^0_{298}$ were calculated according to the following:

$$\Delta_f H^0 = \{ \Delta_{\text{sol}} H^0(\text{YbCl}_3) + n \Delta_{\text{sol}} H^0(\text{AlCl}) \} - \Delta_{\text{sol}} H^0(A_n \text{YbCl}_{3+n})$$

2.2.3. Measurement of emf

A detailed description of the galvanic cell has been given earlier [8]. The measured potentials were generated by the solid state reactions:



in the $300\text{--}400^\circ\text{C}$ range. The solid electrolytes (compressed discs) were separated by A^+ -conducting diaphragms of sintered glass powder. It was not possible to prepare Cs-glasses, stable against YbCl_3 ; furthermore, YbCl_3 itself gave no stable potentials with each

kind of glass. Therefore, measurements could be performed only for Rb- and K-compounds, AYb_2Cl_7 excluded.

3. Results

3.1. Ytterbium(III) chloride

The starting compound $\text{YbCl}_3 \cdot 6\text{H}_2\text{O}$ with a correct content of crystal water was prepared by precipitation from a saturated solution at 0°C with HCl-gas. After filtration and washing with ether it was dried at 40°C . Surprisingly, its crystal structure was still unknown [9]. As for the other hexahydrates, it is isotypic to $\text{PrCl}_3 \cdot 6\text{H}_2\text{O}$ [10], space group $P2_1/n$, $Z=2$:

$$\begin{aligned} a &= 777.3(4) \text{ pm}; & b &= 643.0(5) \text{ pm}; \\ c &= 947.4(9) \text{ pm} \end{aligned}$$

In this structure type, the lanthanide ion is surrounded by two chloride ions and six water molecules, $[\text{YbCl}_2(\text{H}_2\text{O})_6]^+$. These cationic groups are held together by the third Cl^- -ions.

When heated in a drying oven at 100°C , a trihydrate, $\text{YbCl}_3 \cdot 3\text{H}_2\text{O}$, is formed. According to X-ray powder patterns, it is isotypic to $\text{ErCl}_3 \cdot 3\text{H}_2\text{O}$ and $\text{TmCl}_3 \cdot 3\text{H}_2\text{O}$; the structure is still unknown.

Further dehydration must occur in a stream of HCl to avoid hydrolysis. At 120°C , a monohydrate $\text{YbCl}_3 \cdot \text{H}_2\text{O}$ with unknown structure is formed. Above 180°C , the last water is removed. For the preparation of ca. 10 g YbCl_3 , the temperature must be raised slowly up to 350°C . (If the product looks grey because of impurities of carbon, it can be purified by heating for 1 h in a stream of Cl_2 at ca. 500°C .)

Laptev [11] has prepared YbCl_3 from the oxide and CCl_4 -gas at 700°C . His product crystallizes with the layer structure of AlCl_3 (space group $C2/m$, $Z=2$). If the compound is prepared at a lower temperature, e.g. by dehydration of the hydrate, the structure is more or less disturbed. We have already described this phenomenon previously for DyCl_3 [12].

3.2. Ternary ytterbium(III) chlorides from solutions

Some ternary chlorides could be prepared conveniently from aqueous solutions or from acetic acid, as far as they are congruently soluble in these solvents. In

the case of aqueous solutions, AlCl_3 and Yb_2O_3 were dissolved with the correct molar ratio in hydrochloric acid. These solutions were evaporated to dryness at temperatures of 100–120°C. The following compounds could be prepared.

Cs_4YbCl_7 – The crystal structure could be solved by single-crystal measurements, as previously described [13];

Cs_3YbCl_6 – In the orthorhombic modification, space group $Pbcm$ [14];

$\text{Cs}_2\text{YbCl}_5 \cdot \text{H}_2\text{O}$ – Erythro-siderite structure (space group $Pnma$, $Z=4$).

The orthorhombic cell parameters are: $a=1455.3(9)$ pm; $b=1046.6(9)$ pm; $c=751.9(2)$ pm – $\text{Cs}_2\text{YbCl}_5 \cdot \text{H}_2\text{O}$ can be dehydrated by heating in a stream of HCl gas at temperatures rising slowly from 200 to 300°C.

From acetic acid the compounds Cs_3YbCl_6 and Rb_3YbCl_6 , both with the Cs_3BiCl_6 structure (space group $C2/c$) could be precipitated with HCl gas, K_3YbCl_6 (space group $P2_1/c$) was formed with KCl . The solutions were prepared from $\text{YbCl}_3 \cdot 6\text{H}_2\text{O}$ and alkali metal carbonate in the correct molar ratio.

3.3. Phase diagrams and crystal structures

Fig. 1 illustrates the results of the DTA measurements on the systems $\text{AlCl}_3/\text{YbCl}_3$ ($A=\text{Cs}, \text{Rb}, \text{K}$). As already found for ErCl_3 , YbCl_3 also melts and reacts slowly with quartz. Therefore, the melting point was measured in a corundum crucible; the value is 1133 K (860°C). (In the literature: 864°C [11] and 854–880°C)

In the $\text{CsCl}/\text{YbCl}_3$ system, four compounds were found: one congruently melting (Cs_3YbCl_6) and the others, incongruently melting Cs_2YbCl_5 , $\text{Cs}_3\text{Yb}_2\text{Cl}_9$ and CsYb_2Cl_7 . This result is in good agreement with the findings of Blachnik and Selle [2].

In the $\text{RbCl}/\text{YbCl}_3$ system, one congruently melting compound, Rb_3YbCl_6 , and the incongruently melting compounds Rb_2YbCl_5 and RbYb_2Cl_7 exist.

In the KCl/YbCl_3 system, according to Novikov [1], one congruently melting compound, K_3YbCl_6 , exists with a phase transition at 385°C. We found this transition at 392°C and a second, which is not shown in Fig. 1, at 47°C. Another compound – KYb_2Cl_7 – for which the structure was determined by Meyer [17], melts incongruently at 453°C. Still unknown were two

other incongruently melting ternary chlorides: K_2YbCl_5 and $\text{K}_3\text{Yb}_2\text{Cl}_9$, which are stable only in the temperature range from 394° to 425°C.

A typical difficulty for the investigations of phase diagrams with incongruently melting compounds arose. Because of the large temperature difference of the melting point of K_3YbCl_6 (820°C) and the eutectic (409°C at 49.3 mol% YbCl_3) a large amount of K_3YbCl_6 crystallizes from the melt above the peritectic for K_2YbCl_5 at 455°C. The crystals separate from the melt and, therefore, the peritectic reaction at 455°C is not complete. As a consequence, all other effects at lower temperatures, namely the peritectic for $\text{K}_3\text{Yb}_2\text{Cl}_9$ at 425°C and the eutectic, will appear also in the range from K_3YbCl_6 to K_2YbCl_5 . Thus, the quenched samples have to be annealed to get equilibrium conditions. The proof for the existence of the two compounds was eventually found by a combination of DTA measurements for these annealed samples and X-ray patterns:

K_2YbCl_5 : A sample with 33.3 mol% YbCl_3 , annealed at ca. 430°C, only gave one effect at 455°C (peritectic temperature). The X-ray photo was different from that of a mixture between K_3YbCl_6 and KYb_2Cl_7 .

$\text{K}_3\text{Yb}_2\text{Cl}_9$: A 40% sample, annealed at ca. 380°C, gave two thermal effects in a heating curve at 394° and 425°C in addition to the peritectic effect of K_2YbCl_5 at 455°C. The X-ray pattern was a superposition of the peaks of K_2YbCl_5 and KYb_2Cl_7 . In a dynamic high temperature photo new lines appeared at ca. 400°C.

3.3.1. Thermodynamic measurements

Solution enthalpies could be measured for all compounds, $\text{K}_3\text{Yb}_2\text{Cl}_9$ excepted, since is not stable at ambient temperature. The results are given in Table 1, together with two values determined by Blachnik and Selle [15]. For the calculation of the $\Delta_f H^0$ values, the solution enthalpies of the alkali metal chlorides were taken from published papers: $\text{CsCl}=18.1$; $\text{RbCl}=17.6$; $\text{KCl}=17.9$ kJ mol⁻¹. The solution enthalpy of YbCl_3 was found to be $-212.8(7)$ kJ mol⁻¹ (Ref. [16]: 212.9–216.1 kJ mol⁻¹). In the last column of Table 1 ‘syn-reaction enthalpies’ are listed. These are enthalpies of formation from the two direct neighbours of a compound in the phase diagram at ambient temperature.



Fig. 1. Phase diagrams of ACl/YbCl₃ (A=Cs, Rb, K) systems.

Table 1

Solution enthalpies ($\Delta_{\text{sol}}H^0$), enthalpies of formation from the binary compounds (Δ_fH^0) and synreaction enthalpies ($\Delta_{\text{syn}}H^0_{289}$) (all in kJ mol^{-1}) (Range of error for the last digit within parentheses)

Compound	$\Delta_{\text{sol}}H^0_{289}$	Ref. [15]	$\Delta_fH^0_{298}$	$\Delta_{\text{syn}}H^0_{289}$
Cs_4YbCl_7	−54.(2)		−86.1	+8.8
Cs_3YbCl_6 (<i>Pbcm</i>)	−63.6(3)		−94.9	−14.3
Cs_3YbCl_6 (<i>C2/c</i>)	−64.6(1)	−65.3	−93.9	—
Cs_2YbCl_5	−101.4(1)		−75.2	+6.3
$1/2 \text{ Cs}_3\text{Yb}_2\text{Cl}_9$	−110.8(6)		−74.8	−14.8
$1/2 \text{ CsYb}_2\text{Cl}_7$	−174.2(3)		−29.5	−4.6
Rb_3YbCl_6	−74.6(1)		−85.4	−22.3
Rb_2YbCl_5	−114.5(1)		−63.1	+2.3
$1/2 \text{ RbYb}_2\text{Cl}_7$	−166.7(1)		−35.4	−21.5
K_3YbCl_6	−105.5(1)	−105.0	−53.8	−10.5
K_2YbCl_5	−133.7(1)		−43.3	−5.0
$1/2 \text{ KYb}_2\text{Cl}_7$	−188.8(4)		−15.1	−4.3

These reactions are:



Measurements on emf could be performed only for the compounds A_2YbCl_5 and A_3YbCl_6 ($\text{A}=\text{Rb}, \text{K}$) as pointed out in Section 2.2.3. The dependence of emf (ε) on T was linear. Thus, the measured regression lines $\varepsilon=a+b-T$ would be transformed to Gibbs–Helmholtz equation, $\Delta_rG^0=\Delta_rH^0-\Delta_rS^0T$, by means of the relation $G=-nF\varepsilon$.

3.3.2. Rubidium compounds

Reaction: $1.5 \text{ RbCl} + \text{Rb}_{0.5}\text{YbCl}_{3.5} = \text{Rb}_2\text{YbCl}_5$ ($T=620\text{--}690 \text{ K}$)

$$\text{emf/mV} = 211.3 + 0.1346 T/\text{K}$$

$$\Delta_rG^0/\text{kJ mol}^{-1} = -30.6 - 0.0195 T/\text{K}$$

Reaction: $\text{RbCl} + \text{Rb}_2\text{YbCl}_5 = \text{Rb}_3\text{YbCl}_6$ ($T=620\text{--}690 \text{ K}$)

$$\text{emf/mV} = 259.8$$

$$\Delta_rG^0/\text{kJ mol}^{-1} = -25.1$$

3.3.3. Potassium compounds

Reaction: $1.5 \text{ KCl} + \text{K}_{0.5}\text{YbCl}_{3.5} = \text{K}_2\text{YbCl}_5$
(Samples of 50 and 37.5 mol% YbCl_3 in the range of 580–635 K)

$$\text{emf/mV} = 179.4 + 0.1152 T/\text{K}$$

$$\Delta_rG^0/\text{kJ mol}^{-1} = -26.0 - 0.0167 T/\text{K}$$

Measurements above the formation temperature of $\text{K}_3\text{Yb}_2\text{Cl}_9$ yield two reactions:

50 mol%: $\text{KCl} + \text{K}_{0.5}\text{YbCl}_{3.5} = \text{K}_{1.5}\text{YbCl}_{4.5}$ ($T=630\text{--}670 \text{ K}$)


$$\text{emf/mV} = 252.3 + 0.0014 T/\text{K}$$

$$\Delta_rG^0/\text{kJ mol}^{-1} = -24.3$$

with $\text{emf}(\text{K}_2\text{YbCl}_5) = \text{emf}(\text{K}_{1.5}\text{YbCl}_{4.5})$, the formation temperature T_f for K_3YbCl_6 is 641 K (368°C)

37.5 mol%: $0.5\text{KCl} + \text{K}_{1.5}\text{YbCl}_{4.5} = \text{K}_2\text{YbCl}_5$ ($T=630\text{--}670 \text{ K}$)

$$\text{emf/mV} = 121.3 + 0.2045 T/\text{K}$$

Fig. 2. Plot of emf vs. T for K_3YbCl_6 .

$$\Delta_f G^0 / \text{kJ mol}^{-1} = -5.8 - 0.0099 T / \text{K}.$$

From $\text{emf}(\text{HT}) = \text{emf}(\text{LT})$, it follows that $T_f = 650 \text{ K}$ (377°C)

Reaction: $KCl + K_2YbCl_5 = H - K_3YbCl_6$ ($T = 650\text{--}680 \text{ K}$)

$$\text{emf} / \text{mV} = 9.4 + 0.3065 T / \text{K}$$

$$\Delta_f G^0 / \text{kJ mol}^{-1} = -0.9 - 0.0296 T / \text{K}$$

Reaction: $KCl + K_2YbCl_5 = M - K_3YbCl_6$ ($T = 590\text{ to } 650 \text{ K}$)

$$\text{emf} / \text{mV} = 90.4 + 0.1818 T / \text{K}$$

$$\Delta_f G^0 / \text{kJ mol}^{-1} = -8.7 - 0.0175 T / \text{K}$$

The results of the measurements are depicted in Fig. 2.

From $\text{emf}(H - K_3YbCl_6) = \text{emf}(M - K_3YbCl_6)$, it follows that $T = 650 \text{ K}$ (377°C). The transformation enthalpy is $\Delta_r H^0 = 7.8 \text{ kJ mol}^{-1}$, the entropy $\Delta_r S^0 = 12.1 \text{ J K}^{-1} \text{ mol}^{-1}$. It should be pointed out that T_f from the emf measurements is lower than the value from DTA heating curves (392°C) because of kinetic effects. In DTA cooling curves, the transformation could not be detected at all.

The $\Delta_f G^0$ values for the reactions $ACl + A_2YbCl_5 = A_3YbCl_6$ are identical with $\Delta_{\text{syn}} G^0$. For $T = 0$, both

($A = \text{Rb, K}$) are negative, and the compounds are stable at all temperatures.

The synreactions of the 2 : 1 compounds are $0.4A_{0.5}YbCl_{3.5} + 0.6A_3YbCl_6 = A_2YbCl_5$, the free enthalpies $\Delta_{\text{syn}} G^0 = \Delta_f G^0(A_2YbCl_5) - \{0.4\Delta_f G^0(A_{0.5}YbCl_{3.5}) + 0.6\Delta_f G^0(A_3YbCl_6)\}$. With $A_{0.5}YbCl_{3.5}$ as standard compound, one can formulate $\Delta_f G^0$ values:

$$\Delta_f G^0(A_{0.5}YbCl_{3.5}) = 0$$

$$\Delta_f G^0(A_2YbCl_5) = \Delta_r G^0$$

$$\Delta_f G^0(A_3YbCl_6) = \Delta_r G^0(A_2YbCl_5) + \Delta_f G^0(A_3YbCl_6).$$

Then: $\Delta_{\text{syn}} G^0(A_2YbCl_5) = 0.4\Delta_r G^0(A_2YbCl_5) - 0.6\Delta_r G^0(A_3YbCl_6)$.

$$\text{For } Rb_2YbCl_5: \Delta_{\text{syn}} G^0 / \text{kJ mol}^{-1} = +2.8 - 0.0078 T / \text{K}$$

$$\Delta_{\text{syn}} G^0 = 0 \text{ at } 359 \text{ K } (86^\circ\text{C})$$

$$\text{For } K_2YbCl_5: \Delta_{\text{syn}} G^0 / \text{kJ mol}^{-1} = -5.2 + 0.0038 T / \text{K}$$

3.3.4. Crystal structures

For several compounds, the unit cells have already been determined by Meyer et al. from powder patterns.

[17] $CsYb_2Cl_7$ and $RbDy_2Cl_7$ type; space group $RbYb_2Cl_7$: $Pnma$

[17] KYb_2Cl_7 : KDy_2Cl_7 type; space group $P2_1/c$

[18] Rb_3YbCl_6 : Cs_3BiCl_6 type; space group $C2/c$

- [19] Cs₂YbCl₅: Cs₂DyCl₅ type; space group *Pbnm* (id. *Pnma*)
 [20] Cs₃Yb₂Cl₉: Cs₃Tl₂Cl₉ type; space group *R* $\bar{3}$ _c

Our own results about hitherto unknown crystal structures are compiled in Table 3:

Cs₄YbCl₇: The structure for this compound was solved recently by single-crystal work [13].

Cs₃YbCl₆: The structure of the modification in *Pbcm* is discussed elsewhere [14], the structure of the modification in *C2/c* was determined by single-crystal work.

We succeeded in growing suitable crystals from supercritical acetic acid, as recently described for Cs₃CrCl₆

[21]. The results of the single-crystal work are shown in Tables 2 and 3.

H-Cs₃YbCl₆ and *H*-modifications of A₃YbCl₆ (A=Rb,K): Guinier patterns at 500°C could be indexed in the cubic space group *Fm* $\bar{3}$ *m*.

Rb₂YbCl₅: isotypic to Cs₂YbCl₅, space group *Pnma*.

L- and M-K₃YbCl₆: recently we found, by structure determination with single crystals of the analogous erbium compound, again obtained from supercritical acetic acid, that L-K₃ErCl₆ has the monoclinic K₃MoCl₆-structure [22] in space group *P2*₁/*c*. L-K₃YbCl₆ proved to be isotypic. From an X-ray photo at 100°C, we found that the monoclinic *a*- and *c*-axes had become equal, the angle β was 109.4°, which is

Table 2
Single crystal data for Cs₃YbCl₆ and structure refinement

Crystal data

Cs₃YbCl₆; M_r=784.47; monoclinic;
S.G. *C2/c*; Z=8

single-crystal data

a=26.748(7) Å
b=8.1010(5) Å
c=13.028(2) Å
 β =100.00(2)°
V=2780.1(9) Å³

Data collection

MoK α radiation
 Enraf Nonius CAD-4 diffr.
 μ =15.570 mm⁻¹
 ω scans

powder diffraction data

a=26.837(8) Å
b=8.174(4) Å
c=13.035(3) Å
 β =100.2(3)°

λ =0.71073 Å

θ =2–24°

T=293(2) K

h= –3 0 → 0; *k*= –9 → 0;

l= –14 → 14

Refinement

Full-matrix least squares on *F*²

R[*I*>2σ(*I*)]

R(all data)

*R*_{int}=0.0321

Goodness-of-fit on *F*²=1.077

*R*₁=0.0365, *wR*₂=0.1008

*R*₁=0.0459, *wR*₂=0.1073

Fractional atomic coordinates and equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> (eq)
Yb ₁	2500	2500	0	22(1)
Yb ₂	0	–2193(1)	–2500	21(1)
Cs ₃	1525(1)	3135(1)	–3545(1)	33(1)
Cs ₄	3367(1)	3095(1)	–3016(1)	35(1)
Cs ₅	–502(1)	2530(1)	–4330(1)	53(1)
Cl ₁	2502(1)	1154(3)	–1807(2)	32(1)
Cl ₂	561(1)	139(4)	–3076(2)	40(1)
Cl ₃	–568(1)	–2240(4)	–4306(2)	45(1)
Cl ₄	–557(1)	–4444(4)	–1847(2)	42(1)
Cl ₅	1825(2)	405(4)	367(2)	52(1)
Cl ₆	3260(1)	637(4)	786(2)	45(1)

Table 3

Hitherto unknown cell parameters of ternary ytterbium chlorides

Compound	Space group	Z	a/pm	b/pm	c/pm	$\beta/^\circ$
Cs ₄ YbCl ₇	$R\bar{3}m$	3	764.6(4)		2628.9(10)	120.00
Cs ₃ YbCl ₆ (aq.sol.)	<i>Pbcm</i>	8	808.9(3)	1307.7(3)	2636.6(7)	
L-Cs ₃ YbCl ₆	<i>C2/c</i>	8	2683.7(8)	817.4(4)	1303.5(3)	100.2(3)
H-Cs ₃ YbCl ₆	<i>Fm3m</i>	4	1152.2(3)			
H-Rb ₃ YbCl ₆	<i>Fm3m</i>	4	1114.5(2)			
Rb ₂ YbCl ₅	<i>Pnma</i>	4	952.7(3)	726.1(3)	1494.2(4)	
L-K ₃ YbCl ₆	<i>P2₁/c</i>	4	1299.8(5)	762.4(3)	1248.5(4)	109.85(2)
M-K ₃ YbCl ₆	<i>Cmmm</i>	8	1504.4(3)	2128.1(5)	752.7(2)	
H-K ₃ YbCl ₆	<i>Fm3m</i>	4	1086.4(2)			
K ₂ YbCl ₅	(<i>P2₁/c</i>)	4	1141	1008	824	104.1

the tetrahedral angle. Thus, the cell could be transformed into an orthorhombic C-centred cell, space group *Cmmm*. We tried to find the point positions by a Rietveld analysis, by constructing a suitable surveyor's cell.

K₂YbCl₅: The structure of this compound neither belongs to the K₂PrCl₅ nor to the Cs₂DyCl₅ type. Because of a certain similarity to the X-ray pattern of L-K₃YbCl₆, we could perform a monoclinic trial indexing with the extinctions of space group *P2₁/c*.

K₃Yb₂Cl₉: A dynamic Guinier photo in a small temperature range above ca. 380°C displayed lines in addition to those of the mixture (K₂YbCl₅+KYb₂Cl₇). The structure is probably that of the K₃W₂Cl₉ type, space group *P6₃/m*.

4. Discussion

Comparing the systems of ytterbium chloride with the foregoing systems AlCl/TmCl₃, only one difference exists; with ytterbium a high-temperature compound K₃Yb₂Cl₉ is formed. It does not have the structure of Cs₃Yb₂Cl₉, but probably belongs to the K₃W₂Cl₉ type. In the group ALn₂Cl₇, the lanthanide ion has coordination number 7, in other groups – A₃LnCl₆, A₂LnCl₅ and A₃Ln₂Cl₉ – the Ln³⁺ ions are octahedrally surrounded by chloride ions. That will be true also for the still unknown structures of K₂YbCl₅ and K₂TmCl₅. It is to be expected that there the coordination number of K⁺ will be smaller than 10/11 in the

Cs₂DyCl₅ type in which the compounds Cs₂YbCl₅ and Rb₂YbCl₅ crystallize.

The compounds Rb₂LnCl₅ have an increasing stability with decreasing radius of the Ln³⁺ ion. Rb₂HoCl₅ has only a range of existence between 687 and 785 K. Rb₂ErCl₅ is stable above 593 K, its synreaction enthalpy is +6.0 kJ mol⁻¹. The values for Rb₂TmCl₅ are 353 K and +2.4 kJ mol⁻¹, for the ytterbium compound – 359 K and +2.8 kJ mol⁻¹, respectively. The 10/11 interstices, generated by the [LnCl₆]-octahedra reach the optimal size for the Rb⁺ ions.

The synreaction enthalpies for the compounds Cs₂LnCl₅ are: –6.1 kJ mol⁻¹ for Ln=Ho, –1.7 for Er, –1.7 for Tm and +6.3 for Yb. The explanation must be, that the Cs⁺ ion increase for the 10/11 coordination.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie. Their help is gratefully acknowledged.

References

- [1] G.I. Novikov, O.G. Polyachenok, S.A. Frid, Zh. Neorg. Khim. 9 (1964) 472.
- [2] R. Blachnik, D. Selle, Z. Anorg. Allg. Chem. 454 (1979) 90.

- [3] G. Meyer, *Prog. Solid State Chem.* 14 (1982) 141.
- [4] M. Roffe, H.J. Seifert, *J. Alloys Comp.* 257 (1997) 128.
- [5] D. Büchel, J. Krok-Kowalski, H.J. Seifert, *Thermochim. Acta* 282 (1996) 297.
- [6] C. Zheng, H.J. Seifert, *J. Solid State Chem.*, (1997), in press..
- [7] H.J. Seifert, G. Thiel, *Thermochim. Acta* 22 (1978) 363.
- [8] H.J. Seifert, G. Thiel, *J. Chem. Thermodyn.* 14 (1982) 1159.
- [9] E.J. Graeber, G.H. Conrad, S.F. Duliere, *Acta. Crystallogr.* 21 (1966) 1012.
- [10] J. Chen, D. Xu, L.L.J. Wu, G. Xu, *Acta Crystallogr. Sect. C* 47 (1991) 1074.
- [11] D.M. Laptev, A.I. Poshevneva, I.S. Astakhova, N.M. Kulagin, *Zh. Neorg. Khim.* 21 (1976) 2317.
- [12] H.J. Seifert, R. Krämer, *Z. Anorg. Allg. Chem.* 620 (1994) 1543.
- [13] G. Reuter, J. Sebastian, G. Frenzen, *Acta Crystallogr. Sect. C* 52 (1996) 1859.
- [14] G. Reuter, G. Frenzen, *J. Solid State Chem.* 116 (1995) 329.
- [15] R. Blachnik, D. Selle, *Z. Anorg. Allg. Chem.* 454 (1979) 82.
- [16] R. Blachnik, D. Selle, *Thermochim. Acta* 33 (1979) 301.
- [17] G. Meyer, P. Ax, A. Cromm, H. Linzmaier, *J. Solid State Chem.* 98 (1984) 323.
- [18] H. Mattfeld, G. Meyer, *Z. Anorg. Allg. Chem.* 618 (1992) 13.
- [19] G. Meyer, J. Soose, A. Moritz, V. Vitt, T. Hollies, *Z. Anorg. Allg. Chem.* 521 (1985) 169.
- [20] G. Meyer, A. Schöнемund, *Mat. Res. Bull.* 15 (1980) 89.
- [21] M. Prien, H.J. Seifert, *J. Therm. Anal.* 45 (1995) 349.
- [22] Z. Amilius, B. van Laar, H.M. Rietveld, *Acta Crystallogr. Sect. B* 25 (1969) 400.