

Note

Selective Functionalization of Achmatowicz Rearrangement Products by Reactions with Potassium Organotrifluoroborates under Transition-Metal-Free Conditions

Silvia Roscales, Victor Ortega, and Aurelio Garcia Csaky

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.8b01643 • Publication Date (Web): 23 Jul 2018

Downloaded from <http://pubs.acs.org> on July 24, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036


Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Selective Functionalization of Achmatowicz Rearrangement Products by Reactions with Potassium Organotrifluoroborates under Transition-Metal-Free Conditions

Silvia Roscales, Víctor Ortega, and Aurelio G. Csáky*

Instituto Pluridisciplinar, Universidad Complutense, Campus de Excelencia Internacional Moncloa, Paseo de Juan XXIII, 1, 28040 Madrid, Spain


Supporting Information



ABSTRACT: The repertoire of synthetic transformations of the products of the Achmatowicz rearrangement has been expanded by exploring their reactivity with potassium organotrifluoroborates in the absence of transition-metals. Depending on the reaction conditions and the substitution pattern of the starting material, the reaction may lead to the stereoselective synthesis of dihydropyranones (2,6-*trans*), tetrahydropyranones (2,3-*cis*-2,6-*cis*) or functionalized 1,4-dicarbonyl compounds. The method has also been adapted for the one-pot synthesis of functionalized pyrroles.

The Achmatowicz rearrangement (AR) consists of an oxidative ring-expansion rearrangement of readily available 2-furfuryl alcohols **1** to 6-hydroxy-3-pyranones **2** (also known as pyranuloses) (Scheme 1).^{1,2} Due to their dense functionalization, the AR products **2** constitute useful starting materials for the preparation of an ample variety of molecules with dihydropyran or tetrahydropyran core, such as dihydropyranones, oxidopyryliums, δ -lactones, or pyranoses.³

Scheme 1. Previous Work: Synthesis and Transformations of AR Products.




Thus for example, AR products participate as glycosyl donors in Pd-catalyzed glycosylation reactions (**2** \rightarrow **3**, Feringa-O'Doherty *O*-glycosylation),⁴ in bicyclopentamerization reactions (**2** \rightarrow **4**),⁵ as 1,3-dipoles in [5+2]-cycloadditions with alkenes under basic conditions (**2** \rightarrow **5**),⁶ as 2π components in phosphine-catalyzed [3+2]-cycloaddition reactions with allenoates (**2** \rightarrow **6**),⁷ as 2π components in Pd-catalyzed oxa-[3+2]-cycloaddition reactions with 1,3-dicarbonyl compounds (**2** \rightarrow **7**),⁸ or in the Kishi reduction, which leads to *cis*-2,6-dihydropyranones **8**.⁹ Additionally, by means of a two-step sequence consisting of elimination (**2** \rightarrow **9**) followed by Heck-Matsuda arylation (Pd-catalyzed coupling with aryldiazonium salts), they may also give rise to *trans*-2,6-dihydropyranones **10**.¹⁰ Many of these transformations have been used in the synthesis of natural products, bioactive carbohydrates, and advanced pharmaceutical ingredients.

However, taking into consideration the simultaneous presence of the hemiacetal and α,β -unsaturated ketone moieties, the synthetic potential of these densely functionalized compounds as pluripotent molecular platforms for diversity oriented synthesis (DOS) is still poorly developed.¹¹ In this context, we were interested in finding new transformations of the AR products **2** using aryl, alkenyl and alkynyl potassium organotrifluoroborates **11** as reagents. The interest of these reagents in synthesis is due to the fact that they are bench-stable reagents that can be used without the need for protection from humidity,

and are compatible with unprotected functional groups, such as OH.¹² In particular, we have focused our attention on finding selective reaction conditions for the stereoselective replacement of the γ C-O bond in **2** by a γ C-C bond, and on stereoselective conjugate addition reactions to OH-unprotected **2**, both of them under transition-metal-free conditions (Scheme 2).^{13,14} Depending on the reactants and reaction conditions, these transformations will lead selectively to dihydropyranones **10**, tetrahydropyranones **12**,^{1,3} or 1,4-dicarbonyl compounds **13**.¹⁵ In addition, we will consider the application of these findings to the one-pot synthesis of highly functionalized pyrroles (**14**, **15**).¹⁶

Scheme 2. This Paper: Reactions of AR Products with Organotrifluoroborates under Transition-Metal-Free Conditions.




With regard to literature precedents for the replacement of the γ C-O bond in AR products **2** with a γ C-C(sp) bond, alkynylation reactions have only been carried out in connection with the synthesis of spirocyclic pyrans and macrocyclic lactones by the Lewis-acid catalyzed addition of alkynylsilanes.¹⁷ Concerning γ C-C(sp²) bonds, the direct arylation has been reported as a low-yielding process, and the synthesis of **10** has only been previously achieved by a sequence of elimination – Heck-Matsuda reaction.¹⁰ To the best of our knowledge, the direct vinylation reaction is unprecedented.

With respect to literature precedents for conjugate addition reactions to AR products **2**, the addition of alkyl groups using organometallic nucleophiles¹⁸ or radicals¹⁹ has only been carried out on *O*-protected derivatives of the AR products **2**, and is amply preceded. However, the addition of vinyl or aryl

groups is underdeveloped, and has only been carried out on *O*-protected AR products **2**.^{20,21} To the best of our knowledge, the conjugate addition of C(sp)-nucleophiles has not been previously reported.

We started our investigations by considering the selective replacement of the γ C-O bond in *O*-unprotected **2** with a γ C-C(sp²) bond (Scheme 3) toward the synthesis of **10**.²² The first challenge of this transformation was finding reaction conditions devoid of competence with conjugate addition to the α,β -unsaturated ketone moiety of the AR starting materials **2**, which would lead to **12**.²³ In addition, we were concerned about the stereochemical outcome of the transformation, which could lead to the dihydropyranones **10** either in 2,6-*cis* or 2,6-*trans* relative configuration. In particular, the diastereoselective obtention of 2,6-*trans*-dihydropyranones is challenging.¹⁰ After some experimentation using **2a** as starting material, we were pleased to find that the reaction with potassium (*E*)-phenylvinyl trifluoroborate **11a** gave rise to the selective formation of **10a** either when promoted by trifluoroacetic anhydride (TFAA, Scheme 3, conditions A) or by HBF₄ (Scheme 3, conditions B). Under both reaction conditions, **10a** was obtained as a single 2,6-*trans* diastereomer.²⁴

Scheme 3. Reactions of **2a** with **11a**




Complementarily, we looked for reaction conditions that could selectively afford the conjugate addition products **12** without formation of **10**. Again using **2a** as starting material, we found that the reaction with (*E*)-phenylvinyl potassium trifluoroborate **11a** promoted by Bu₄NHSO₄ (Scheme 3, conditions C)

produced selectively the conjugate addition product **12a**. However, since **12a** was unstable to purification by silica gel chromatography, the crude reaction product was acetylated *in situ* to **16a**. Under both reaction conditions, **16a** was obtained with high 2,3-*cis*-2,6-*cis* diastereoselectivity.^{24,25}


Under the optimum reaction conditions for the formation of **10a** (TFAA promotion, -15 °C to rt), we extended our observations to other AR products **2** for the synthesis of various 2,6-*trans*-dihydropyranones **10** (Scheme 4).^{24,26} We found that the additions of arylvinyltrifluoroborates and alkylvinyltrifluoroborates to **2a** took place smoothly to give the corresponding dihydropyranones **10b** – **10e** in good yields and with *trans* : *cis* dr > 98:02. Similar results were obtained for the reactions of **2a** with potassium aryltrifluoroborates (**10f** – **10h**). However, an erosion of the diastereoselectivity was observed in the addition of potassium alkynyltrifluoroborates (**10i**, **10j**). In these cases, we observed that lowering the reaction temperature to -40 °C afforded compounds **10i** and **10j** with increased *trans* : *cis* dr. Good yields and *trans* : *cis* dr > 98:02 were also obtained when starting from the AR products **2b** and **2c** (synthesis of **10k** – **10n**). We were also glad to observe that the vinylation and arylation reactions of compound **2d**, with a quaternary carbon center, also took place in good yield and full 2,6-*trans* diastereoselectivity (synthesis of **10o**, **10p**), although the alkynylation reaction (synthesis of **10q**), as well as the reactions of **2e** (synthesis of **10r**, **10s**), were less stereoselective. In these cases, we were not able to improve the diastereoselectivity by diminishing the reaction temperature.

Scheme 4. Diastereoselective synthesis of dihydropyranones 10b-s




In a similar fashion, under the optimum reaction conditions for the formation of **16a** (promotion with Bu_4NHSO_4 , rt), we extended our observations to other representative examples of potassium organotrifluoroborates **11** in their reactions with **2a** for the synthesis of various tetrahydropyranones **16** (Scheme 5). The new compounds **16b-e** have been also obtained with high 2,3-*cis*-2,6-*cis* diastereoselectivity.

Scheme 5. Diastereoselective synthesis of tetrahydropyranones 16b-e




However, when the γ -carbon of the AR product **2** was fully substituted (Scheme 6, **2f** – **2i**, $R^2 \neq H$), the outcome of the reaction with potassium organotrifluoroborates **11** changed drastically. Instead of the expected dihydropyranones **10** or tetrahydropyranones **12**, the reaction gave rise to the 1,4-dicarbonyl compounds **13**.¹⁵ The reaction was selective for α -functionalization adjacent to the hydroxymethylcarbonyl group. As observed for **13a**, this result was independent of the method of activation (**13a**: TFAA, 80%; HBF_4 , 62%; Bu_4NHOSO_4 , 75%). Under the best conditions found for **13a** (TFAA activation) the reaction was extended to several other examples (Scheme 6).

Scheme 6. Regioselective synthesis of 1,4-dicarbonyl compounds **13a-h**



The reactions of potassium organotrifluoroborates **11** with the AR products **2** can be understood (Scheme 7) by considering the direct conjugate addition to the enone moiety *versus* addition to the oxonium **I** that results from dehydroxylation.²²

Scheme 7. Proposed Reaction Course.




In the presence of TFAA or HBF₄, formation of the oxonium **I** is expected to be easy. Direct addition of **11** to the electrophilic γ -carbon in **I** accounts for the formation of dihydropyranones **10**.²⁷ The relative stereochemistry of the final products **10** is governed by *trans* addition to the most bulky group (L) on the starting material. This type of reaction would be hindered for R¹ \neq H. On the other hand, organotri-fluoroborates are known to be in equilibrium with organodifluoroboranes under a variety of circumstances.^{28,29} Thus, under reaction conditions that do not favor formation of **I** (Bu₄NHSO₄), coordination of a trivalent boron species to the OH group in **2** may trigger the conjugate addition reaction that leads to tetrahydropyranones **12**. This type of addition is presumed to take place from the β -face on a reactive conformation with the most bulky group (L) at a pseudoequatorial position.³⁰ The formation of the 1,4-dicarbonyl compounds **13** can be understood by an OH-triggered conjugate addition to the open chain intermediates **II**, which may be in equilibrium with **2** (R¹ \neq H).³¹

Finally, we have considered the application of this methodology to the one-pot synthesis of highly functionalized pyrroles (Scheme 8).¹⁶ Thus, reaction of **2a** or **2j** with compounds **11** in the presence of Bu₄NHSO₄ followed by the addition of benzylamine gave rise to pyrroles **14**, whereas reaction of **2f** or **2g** with compounds **11** in the presence of TFAA followed by the addition of benzylamine gave rise to pyrroles **15** in good yields. Both types of transformations were tolerant with the presence of a free OH

group. It is worth mentioning the complementary relative disposition of the β -substituent and the α -hydroxyalkyl chain in pyrroles **14** (2,4-relative disposition) and **15** (2,3-relative disposition).

Scheme 8. Synthesis of pyrroles **14** and **15**



In conclusion, the results presented in this paper put forward new valorizations of Achmatowicz rearrangement products **2** using potassium organotrifluoroborates **11** as reagents in the absence of transition-metals. The substitution pattern at position γ of the starting material **2** is crucial in determining the course of the reaction. When the γ -carbon is fully substituted, the reaction leads to 1,4-dicarbonyl compounds **13** independently of the reaction conditions. Compounds **13** show the particular feature of functionalization in α -position to the carbonyl group of a hydroxymethylcarbonyl group. When the γ -carbon of **2** is tertiary, the reaction can be controlled to obtain selectively the dihydropyranones **10** (TFAA-promotion, 2,6-*trans*) or the tetrahydropyranones **12** (Bu_4NHSO_4 -promotion, 2,3-*cis*-2,6-*cis*). Compounds **10**, **12** and **13** are useful intermediates for the synthesis of an ample diversity of products. As an

example, crude **12** and **13** have been transformed into functionalized pyrroles **14** and **15** bearing α -hydroxyalkyl chains in a one-pot operation.

EXPERIMENTAL SECTION

General information. Unless otherwise stated, all starting materials acids were commercially available research-grade chemicals and were used without further purification. Silica gel 60 F254 was used for TLC, and the spots were detected with UV light (254 and/or 366 nm) and/or vanillin solution. Flash column chromatography was carried out on silica gel 60. ^1H -NMR spectra were recorded at 300 or 500 MHz, ^{13}C -NMR spectra were recorded at 75 or 125 MHz, and ^{19}F -NMR spectra were recorded at 282 MHz, all of them in CDCl_3 , or acetone- d_6 solution.

Synthesis of 1a-g, 1i and 1j. 1-(Furan-2-yl)ethanol (**1a**),³² furan-2-yl(phenyl)methanol (**1b**),³³ 1-(furan-2-yl)-2-methylpropan-1-ol (**1c**),³⁴ 2-(furan-2-yl)-3-methylbutan-2-ol (**1d**),³⁵ 1-(furan-2-yl)-1-phenylethanol (**1e**),³⁶ (5-methylfuran-2-yl)methanol (**1f**),³⁷ (5-phenylfuran-2-yl)methanol (**1g**),³⁴ 1-(5-methylfuran-2-yl)ethanol (**1i**),³⁸ and furan-2-ylmethanol (**1j**)³⁹ were synthesized following previous literature procedures.

(4,5-Dimethylfuran-2-yl)methanol (1h): 4,5-Dimethylfurfural (300 mg, 2.2 mmol) was dissolved in anhydrous methanol (1.3 mL) and sodium borohydride (163.4 mg, 4.5 mmol) was added at 0°C. The mixture was stirred for 1 h at room temperature, water was then added, and the aqueous phase was extracted with dichloromethane. The combined organic layers were dried over MgSO_4 and filtered. The solvent was evaporated and the crude product was purified via flash column chromatography on silica gel eluting with hexane:AcOEt 8:2 (260.6 mg, 94% yield). Yellow oil. ^1H NMR (300 MHz, CDCl_3) δ : 6.05 (s, 1H), 4.50 (s, 2H), 2.19 (s, 3H), 1.91 (s, 3H), 1.76 (bs, 1H) ppm; ^{13}C NMR (75 MHz, CDCl_3) δ 151.1, 147.7, 114.8, 111.4, 57.7, 11.5, 9.91 ppm. Anal. calcd. for $\text{C}_7\text{H}_{10}\text{O}_2$: C, 66.65; H, 7.99. Found: C, 66.69; H, 7.98.

Synthesis of 2a-c, 2f, 2g, 2i, and 2j. 6-Hydroxy-2-methyl-2H-pyran-3(6H)-one (**2a**),⁴⁰ 6-hydroxy-2-phenyl-2H-pyran-3(6H)-one (**2b**),⁴¹ 6-hydroxy-2-isopropyl-2H-pyran-3(6H)-one (**2c**),⁴¹ 6-hydroxy-6-methyl-2H-pyran-3(6H)-one (**2f**),³¹ 6-hydroxy-6-phenyl-2H-pyran-3(6H)-one (**2g**),⁴¹ 6-

hydroxy-2,6-dimethyl-2H-pyran-3(6H)-one (**2i**),⁴² and 6-hydroxy-2H-pyran-3(6H)-one (**2j**)⁴³ were synthesized following previous literature procedures.

General procedure for the synthesis of 2d, 2e, and 2h. To a stirred solution of the furfuryl alcohol **1** (1 mmol) in DCM (8 mL) was added *m*-chloroperbenzoic acid (*m*CPBA, 1.5 mmol) at 0 °C. After completion of the addition, the reaction was allowed to stir at 0 °C for 1 h. Evaporation of the solvent gave the crude product which was purified via flash column chromatography on silica gel.

6-Hydroxy-2-isopropyl-2-methyl-2H-pyran-3(6H)-one (2d): Colorless oil. Major diastereomer. ¹H NMR (300 MHz, CDCl₃) δ: 6.83 (d, *J* = 10.3 Hz, 1H), 6.05 (d, *J* = 10.3 Hz, 1H), 5.70-5.69 (m, 1H), 4.55 (bs, 1H), 2.15-2.06 (m, 1H), 1.34 (s, 3H), 0.90-0.83 (m, 6H) ppm. ¹³C NMR (75 MHz, CDCl₃) δ: 200.0, 145.3, 126.9, 87.6, 84.3, 33.5, 21.9, 17.3, 16.3 ppm. Minor diastereomer. ¹H NMR (300 MHz, CDCl₃) δ: 6.84 (d, *J* = 10.3 Hz, 1H), 6.03 (d, *J* = 10.3 Hz, 1H), 5.68-5.67 (m, 1H), 4.55 (bs, 1H), 2.28-2.21 (m, 1H), 1.25 (s, 3H), 0.90-0.83 (m, 6H) ppm. ¹³C NMR (75 MHz, CDCl₃) δ: 200.1, 146.7, 127.4, 87.7, 83.8, 34.5, 18.9, 17.2, 16.1 ppm. Anal. calcd. for C₉H₁₄O₃: C, 63.51; H, 8.29. Found: C, 63.58; H, 8.30.

6-Hydroxy-2-methyl-2-phenyl-2H-pyran-3(6H)-one (2e): White solid (m.p: 77-79°C). Major diastereomer. ¹H NMR (300 MHz, CDCl₃) δ: 7.36-7.33 (m, 5H), 6.79 (d, *J* = 10.3 Hz, *J* = 1.3 Hz, 1H), 6.20 (d, *J* = 10.3 Hz, *J* = 1.7 Hz, 1H), 5.42 (s, 1H), 3.24 (bs, 1H), 1.67 (s, 3H) ppm. ¹³C NMR (750 MHz, CDCl₃) δ: 196.2, 147.4, 138.5, 128.9 (2C), 128.5, 128.4, 125.5 (2C), 88.6, 83.6, 27.8 ppm. Anal. calcd. for C₁₂H₁₂O₃: C, 70.57; H, 5.92. Found: C, 70.61; H, 5.92.

6-Hydroxy-5,6-dimethyl-2H-pyran-3(6H)-one (2h): Colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 5.90 (s, 1H), 4.48 (d, *J* = 16.9 Hz, 1H), 4.08 (d, *J* = 16.9 Hz, 1H), 3.51 (bs, 1H), 2.04-2.03 (m, 3H), 1.61 (s, 3H) ppm; ¹³C NMR (75 MHz, CDCl₃): δ 195.5, 161.0, 124.4, 95.4, 66.6, 25.9, 19.9 ppm. Anal. calcd. for C₇H₁₀O₃: C, 59.14; H, 7.09. Found: C, 59.08; H, 7.08.

General Procedure for the Synthesis of compounds 10 and 13, Conditions A: A stirred solution of compound **2** (0.23 mmol, 1.0 equiv) and the corresponding potassium trifluoroborate **11** (0.28 mmol, 1.2 equiv) in DCM (1.4 mL, 6 mL/mmol) was cooled to -15°C. After stirring for 5 min, TFAA (0.23-0.52 mmol, 1.0 - 2.2 equiv) was added and the reaction mixture was allowed to warm to rt. Then, the solvent was removed in vacuo and the crude product was purified by flash column chromatography on silica gel.

General Procedure for the Synthesis of compounds 10, Conditions B. A stirred solution of compound **2** (0.23 mmol, 1.0 equiv) and the corresponding potassium *trans*-styryltrifluoroborate **11** (0.35 mmol, 1.5 equiv) in MeCN (2.3 mL, 10 mL/mmol) was cooled to 0°C. After stirring for 5 min, HBF₄·OEt₂ (0.35 mmol, 1.5 equiv) was added, and the solution was stirred at 0 °C for 15 min. The reaction was quenched with water (15 mL) and extracted with ethyl acetate (3 x 10 mL). The organic layer was dried with MgSO₄ and concentrated. The crude product was purified by flash column chromatography on silica gel.

General Procedure for the Synthesis of compounds 16, Conditions C: A solution of compound **2** (0.23 mmol, 1.0 equiv), the corresponding potassium trifluoroborate **11** (0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogensulfate (0.12 mmol, 0.5 equiv) in DCM (4.6 mL, 20 mL/mmol) was allowed to stir at rt for 24-48h. After reaction completion as determined by TLC analysis, the crude mixture was cooled to 0°C, and Ac₂O (0.23 mmol, 1.0 equiv), DMAP (0.012 mmol, 5% mol) and Et₃N (0.47 mmol, 2.0 equiv) were added, and the reaction mixture was stirred at 0°C for 1 h. Then, the mixture was concentrated and purified via flash column chromatography on silica gel.

General Procedure for the one-pot Synthesis of pyrroles 14: A solution of compound **2** (0.23 mmol, 1.0 equiv), the corresponding potassium trifluoroborate **11** (0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogensulfate (0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL, 20 mL/mmol) was allowed to stir at rt for 24-48h. After reaction completion as determined by TLC analysis, the solvent was removed in vacuo and the residue was redissolved in toluene (1.1 mL, 5 mL/mmol). Benzylamine (0.23 mmol, 1 equiv) and *p*-toluenesulfonic acid (0.023 mmol, 0.1 equiv) were added and the mixture was

1 stirred at rt. Then, the mixture was concentrated and purified via flash column chromatography on silica
2 gel.
3

4
5 **General Procedure for the one-pot Synthesis of pyrroles 15:** A stirred solution of compound **2**
6 (0.23 mmol, 1.0 equiv) and the corresponding potassium trifluoroborate **11** (0.28 mmol, 1.2 equiv) in
7 DCM (1.4 mL, 6 mL/mmol) was cooled to -15°C. After stirring for 5 min, TFAA (0.23-0.52 mmol, 1.0 -
8 2.2 equiv) was added and the reaction mixture was allowed to warm to rt. Then, the solvent was re-
9 moved in vacuo and the residue was redissolved in toluene (1.1 mL, 5 mL/mmol). Benzylamine (0.23
10 mmol, 1 equiv) and *p*-toluenesulfonic acid (0.023 mmol, 0.1 equiv) were added and the mixture was
11 stirred at rt. Then, the mixture was concentrated and purified via flash column chromatography on silica
12 gel.
13
14
15
16
17
18
19
20
21
22

23 **(2*R*,6*R*)-2-Methyl-6-((*E*)-styryl)-2*H*-pyran-3(6*H*)-one (10a):** Following general procedure for
24 the synthesis of compounds **10**, Conditions A, starting from **2a** (30.1 mg, 0.23 mmol, 1.0 equiv), potas-
25 sium *trans*-styryltrifluoroborate **11a** (59.0 mg, 0.28 mmol, 1.2 equiv) and TFAA (33 µL, 0.23 mmol, 1.0
26 equiv) in DCM anh. (1.4 mL) for 1 h. The crude product was purified by flash column chromatography
27 on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (45.1 mg, 90% yield). Following
28 general procedure for the synthesis of compounds **10**, Conditions B, starting from **2a** (30.3 mg, 0.23
29 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (73.8 mg, 0.35 mmol, 1.5 equiv) and
30 HBF₄·OEt₂ (48 µL, 0.35 mmol, 1.5 equiv.) in MeCN anh. (2.3 mL). The crude product was purified by
31 flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (37.6
32 mg, 75% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.17-7.39 (m, 5H), 7.00 (dd, *J* = 10.4 Hz, *J* = 3.5 Hz,
33 1H), 7.60 (d, *J* = 16.1 Hz, 1H), 6.24 (dd, *J* = 16.1 Hz, *J* = 6.2 Hz, 1H), 6.10 (dd, *J* = 10.4 Hz, *J* = 1.7 Hz,
34 1H), 5.01-5.08 (m, 1H), 4.34 (c. *J* = 6.8 Hz, 1H), 1.34 (d, *J* = 6.8 Hz, 3H) ppm. ¹³C-NMR (75 MHz,
35 CDCl₃): δ 196.9, 148.9, 135.9, 134.6, 128.9, 128.6, 126.8, 126.3, 124.0, 72.8, 71.6, 15.6 ppm. Anal.
36 calcd. for C₁₄H₁₄O₂: C, 78.48; H, 6.59. Found: C, 78.55; H, 6.58.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 **(2*R*,6*R*)-6-((*E*)-4-Fluorostyryl)-2-methyl-2*H*-pyran-3(6*H*)-one (10b):** Following general pro-
57 cedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (29.7 mg, 0.23 mmol, 1.0
58
59
60

equiv), potassium 2-(4-fluorophenyl)vinyltrifluoroborate **11b** (64.1 mg, 0.28 mmol, 1.2 equiv) and TFAA (33 μ L, 0.23 mmol, 1.0 equiv) in DCM anh. (1.4 mL) for 2 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (51.6 mg, 95% yield). ^1H NMR (300 MHz, CDCl_3): δ 7.34-7.42 (m, 2H), 6.97-7.09 (m, 3H), 6.63 (d, J = 16.1 Hz, 1H), 6.23 (dd, J = 16.1 Hz, J = 6.0 Hz, 1H), 6.16 (dd, J = 10.4 Hz, J = 1.8 Hz, 1H), 5.06-5.13 (m, 1H), 4.40 (c. J = 6.8 Hz, 1H), 1.41 (d, J = 6.8 Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 196.8, 164.6, 161.3, 148.8, 133.4, 132.1, 132.0, 128.5, 128.4, 126.3, 123.8, 123.7, 116.0, 115.7, 72.9, 71.5, 15.6 ppm. Anal. calcd. for $\text{C}_{14}\text{H}_{13}\text{FO}_2$: C, 72.40; H, 5.64. Found: C, 72.49; H, 5.62.

(2R,6R)-6-((E)-4-Methoxystyryl)-2-methyl-2H-pyran-3(6H)-one (10c): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (29.7 mg, 0.23 mmol, 1.0 equiv), potassium 2-(4-methoxyphenyl)vinyltrifluoroborate **11c** (67.4 mg, 0.28 mmol, 1.2 equiv) and TFAA (33 μ L, 0.23 mmol, 1.0 equiv) in DCM anh. (1.4 mL) for 2 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (37.2 mg, 65% yield). ^1H NMR (500 MHz, CDCl_3): δ 7.34 (d, J = 8.7 Hz, 2H), 7.06 (dd, J = 10.8 Hz, J = 4.3 Hz, 1H), 6.87 (d, J = 8.7 Hz, 1H), 6.60 (d, J = 16.0 Hz, 1H), 6.14-6.20 (m, 2H), 5.06-5.11 (m, 1H), 4.40 (c. J = 6.8 Hz, 1H), 3.82 (s, 3H), 1.41 (d, J = 6.8 Hz, 3H) ppm. ^{13}C -NMR (125 MHz, CDCl_3): δ 197.0, 160.0, 149.2, 134.3, 128.7, 128.1, 126.2, 121.6, 114.3, 72.7, 71.8, 55.5, 15.6 ppm. Anal. calcd. for $\text{C}_{15}\text{H}_{16}\text{O}_3$: C, 73.75; H, 6.60. Found: C, 73.64; H, 6.58.

(2R,6S)-6-((E)-3-Chloroprop-1-en-1-yl)-2-methyl-2H-pyran-3(6H)-one (10d): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (30.1 mg, 0.23 mmol, 1.0 equiv), potassium *trans*-2-chloromethylvinyltrifluoroborate **11d** (51.2 mg, 0.28 mmol, 1.2 equiv) and TFAA (39 μ L, 0.28 mmol, 1.2 equiv) in DCM anh. (1.4 mL) for 2 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (40.2 mg, 92% yield). ^1H NMR (300 MHz, CDCl_3): δ 6.96 (dd, J = 10.4 Hz, J = 3.4 Hz, 1H), 6.12 (dd, J = 10.4 Hz, J = 1.7 Hz, 1H), 5.86-6.01 (m, 2H), 4.93-5.00 (m, 1H), 4.32 (c. J = 6.8 Hz, 1H), 4.00-

4.16 (m, 2H), 1.38 (d, $J = 6.8$ Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 196.5, 148.1, 130.7, 129.4, 126.3, 73.0, 70.4, 43.8, 15.4 ppm. Anal. calcd. for $\text{C}_9\text{H}_{11}\text{ClO}_2$: C, 57.92; H, 5.94. Found: C, 57.99; H, 5.92.

(2*R*,6*R*)-2-Methyl-6-((*E*)-3-phenylprop-1-en-1-yl)-2*H*-pyran-3(6*H*)-one (10e): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (30.0 mg, 0.23 mmol, 1.0 equiv), potassium *trans*-3-phenyl-1-propen-1-yltrifluoroborate **11e** (63.0 mg, 0.28 mmol, 1.2 equiv) and TFAA (39 μL , 0.28 mmol, 1.2 equiv) in DCM anh. (1.4 mL) for 2 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (49.7 mg, 93% yield). ^1H NMR (300 MHz, CDCl_3): δ 7.05-7.31 (m, 5H), 6.89 (dd, $J = 10.2$ Hz, $J = 3.4$ Hz, 1H), 6.00 (dd, $J = 10.2$ Hz, $J = 1.3$ Hz, 1H), 5.87 (dt, $J = 15.5$ Hz, $J = 16.7$ Hz, 1H), 5.59 (dd, $J = 15.5$ Hz, $J = 6.7$ Hz, 1H), 4.82-4.90 (m, 1H), 4.26 (c. $J = 6.8$ Hz, 1H), 3.37 (d, $J = 6.7$ Hz, 2H), 1.31 (d, $J = 6.8$ Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 197.0, 149.4, 139.3, 135.3, 128.7, 128.6, 126.5, 126.2, 125.9, 72.7, 71.3, 38.9, 15.5 ppm. Anal. calcd. for $\text{C}_{15}\text{H}_{16}\text{O}_2$: C, 78.92; H, 7.06. Found: C, 78.90; H, 7.07.

(2*R*,6*S*)-2-Methyl-6-phenyl-2*H*-pyran-3(6*H*)-one (10f): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (29.5 mg, 0.23 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (51.7 mg, 0.28 mmol, 1.2 equiv) and TFAA (39 μL , 0.28 mmol, 1.2 equiv) in DCM anh. (1.4 mL) for 2 h. Then, a second equivalent of TFAA (33 μL , 0.23 mmol, 1.0 equiv) was added and the reaction mixture was allowed to warm to room temperature for 4 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (37.4 mg, 85% yield). ^1H NMR (300 MHz, CDCl_3): δ 7.37-7.43 (m, 5H), 7.16 (dd, $J = 10.4$ Hz, $J = 3.1$ Hz, 1H), 6.23 (dd, $J = 10.4$ Hz, $J = 1.9$ Hz, 1H), 5.48-5.54 (m, 1H), 4.26 (c. $J = 6.9$ Hz, 1H), 1.40 (d, $J = 6.9$ Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 197.0, 149.3, 136.9, 129.0, 128.9, 128.2, 126.2, 73.3, 72.8, 15.4 ppm. Anal. calcd. for $\text{C}_{12}\text{H}_{12}\text{O}_2$: C, 76.57; H, 6.43. Found: C, 76.62; H, 6.42.

(2*R*,6*S*)-6-(4-Fluorophenyl)-2-methyl-2*H*-pyran-3(6*H*)-one (10g): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (29.9 mg, 0.23 mmol, 1.0 equiv), potassium 4-fluorophenyltrifluoroborate **11g** (56.8 mg, 0.28 mmol, 1.2 equiv) and TFAA (39 μ L, 0.28 mmol, 1.2 equiv) in DCM anh. (1.4 mL) for 2 h. Then, the mixture was cooled at 0°C and a second equivalent of TFAA (33 μ L, 0.23 mmol, 1.0 equiv) was added and the reaction mixture was allowed to warm to room temperature for 4 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (44.9 mg, 93% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.34-7.44 (m, 2H), 7.04-7.17 (m, 3H), 6.23 (dd, J = 10.4 Hz, J = 1.9 Hz, 1H), 5.46-5.50 (m, 1H), 4.23 (c. J = 6.8 Hz, 1H), 1.40 (d, J = 6.8 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.8, 164.7, 161.4, 148.9, 132.9, 132.8, 130.0, 129.9, 126.3, 116.0, 115.8, 73.3, 72.0, 15.4 ppm. Anal. calcd. for C₁₂H₁₁FO₂: C, 69.89; H, 5.38. Found: C, 69.85; H, 5.37.

(2*R*,6*S*)-6-(2-Methoxyphenyl)-2-methyl-2*H*-pyran-3(6*H*)-one (10h): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (30.1 mg, 0.23 mmol, 1.0 equiv), potassium 2-methoxyphenyltrifluoroborate **11h** (60.1 mg, 0.28 mmol, 1.2 equiv) and TFAA (33 μ L, 0.23 mmol, 1.0 equiv) in DCM anh. (1.4 mL) for 3 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (28.1 mg, 55% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.30-7.40 (m, 2H), 7.08 (dd, J = 10.3 Hz, J = 2.8 Hz, 1H), 6.92-7.01 (m, 2H), 6.17 (dd, J = 10.4 Hz, J = 2.2 Hz, 1H), 5.88-5.94 (m, 1H), 4.38 (c. J = 7.0 Hz, 1H), 3.89 (s, 3H), 1.43 (d, J = 7.0 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 197.5, 157.1, 150.2, 130.0, 128.3, 125.5, 125.2, 120.7, 111.1, 74.1, 66.8, 55.8, 15.4 ppm. Anal. calcd. for C₁₃H₁₄O₃: C, 71.54; H, 6.47. Found: C, 71.61; H, 6.45.

(2*R*,6*S*)-2-Methyl-6-(phenylethynyl)-2*H*-pyran-3(6*H*)-one (10i): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (30.1 mg, 0.23 mmol, 1.0 equiv), potassium phenylethynyltrifluoroborate **11i** (58.5 mg, 0.28 mmol, 1.2 equiv) and TFAA (33 μ L, 0.23 mmol, 1.0 equiv) in DCM anh. (1.4 mL) at -40°C for 3 h. The crude product was purified by flash col-

umn chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (40.2 mg, 81% yield, *dr* 91:09). ¹H NMR (300 MHz, CDCl₃): δ 7.43-7.50 (m, 2H), 7.32-7.38 (m, 3H), 7.03 (dd, *J* = 10.2 Hz, *J* = 4.2 Hz, 1H), 6.11 (dd, *J* = 10.2 Hz, *J* = 1.5 Hz, 1H), 5.37 (dd, *J* = 4.2 Hz, *J* = 1.5 Hz, 1H), 4.67 (c. *J* = 6.7 Hz, 1H), 1.44 (d, *J* = 6.7 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.4, 146.3, 132.1, 129.2, 128.5, 125.8, 121.8, 87.9, 81.8, 73.3, 63.4, 15.4 ppm. Anal. calcd. for C₁₄H₁₂O₂: C, 79.22; H, 5.70. Found: C, 79.29; H, 5.69.

(2*R*,6*S*)-6-(Cyclohexylethynyl)-2-methyl-2*H*-pyran-3(6*H*)-one (10j): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2a** (30.4 mg, 0.23 mmol, 1.0 equiv), potassium cyclohexylethynyltrifluoroborate **11j** (60.1 mg, 0.28 mmol, 1.2 equiv) and TFAA (33 μL, 0.23 mmol, 1.0 equiv) in DCM anh. (1.4 mL) at -40°C for 3 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (37.3 mg, 72% yield). ¹H NMR (300 MHz, CDCl₃): δ 6.92 (dd, *J* = 10.1 Hz, *J* = 4.2 Hz, 1H), 6.00 (dd, *J* = 10.1 Hz, *J* = 1.4 Hz, 1H), 5.09-5.17 (m, 1H), 4.56 (c. *J* = 6.7 Hz, 1H), 2.36-2.47 (m, 1H), 1.40 (d, *J* = 6.7 Hz, 3H), 1.18-1.84 (m, 10H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.7, 147.4, 125.1, 93.2, 72.9, 63.1, 32.5, 32.4, 29.2, 25.9, 24.9, 15.3 ppm. Anal. calcd. for C₁₄H₁₈O₂: C, 77.03; H, 8.31. Found: C, 77.00; H, 8.25.

(2*S*,6*S*)-2-Phenyl-6-((*E*)-styryl)-2*H*-pyran-3(6*H*)-one (10k): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2b** (30.0 mg, 0.16 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (39.9 mg, 0.19 mmol, 1.2 equiv) and TFAA (23 μL, 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (28.7 mg, 65% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.43-7.31 (m, 10H), 7.07 (dd, *J* = 10.4 Hz, *J* = 2.8 Hz, 1H), 6.73 (d, *J* = 16.0 Hz, 1H), 6.28-6.36 (m, 2H), 5.34 (s, 1H), 5.04-5.06 (m, 1H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 194.4, 149.5, 136.0, 134.8, 134.4, 128.9 (2C), 128.7 (3C), 128.6, 127.7 (2C), 127.0, 126.9 (2C), 124.6, 79.4, 71.1 ppm. Anal. calcd. for C₁₉H₁₆O₂: C, 82.58; H, 5.84. Found: C, 82.51; H, 5.86.

(2*S*,6*R*)-2,6-Diphenyl-2*H*-pyran-3(6*H*)-one (10l): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2b** (30.0 mg, 0.16 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (35.0 mg, 0.19 mmol, 1.2 equiv) and TFAA (23 μ L, 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 24 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (22.8 mg, 57% yield). ^1H NMR (300 MHz, CDCl_3): δ 7.41-7.39 (m, 10H), 7.10 (dd, $J = 10.4$ Hz, $J = 2.4$ Hz, 1H), 6.37 (dd, $J = 10.4$ Hz, $J = 2.4$ Hz, 1H), 5.35 (t, $J = 2.4$ Hz, 1H), 5.31 (s, 1H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 194.5, 150.3, 137.8, 134.4, 129.1 (2C), 129.0, 128.8 (2C), 128.7, 127.9 (2C), 127.6 (2C), 126.8, 79.9, 72.3 ppm. Anal. calcd. for $\text{C}_{17}\text{H}_{14}\text{O}_2$: C, 81.58; H, 5.64. Found: C, 81.55; H, 5.71.

(2*S*,6*S*)-2-Isopropyl-6-((*E*)-styryl)-2*H*-pyran-3(6*H*)-one (10m): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2c** (29.4 mg, 0.19 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (47.5 mg, 0.23 mmol, 1.2 equiv) and TFAA (26 μ L, 0.19 mmol, 1.0 equiv) in DCM anh. (1.1 mL) for 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (31.7 mg, 69% yield). ^1H NMR (300 MHz, CDCl_3): δ 7.41-7.28 (m, 5H), 7.03 (dd, $J = 10.4$ Hz, $J = 3.4$ Hz, 1H), 6.66 (d, $J = 16.2$ Hz, 1H), 6.26 (dd, $J = 16.2$ Hz, $J = 6.0$ Hz, 1H), 6.14 (dd, $J = 10.4$ Hz, $J = 1.9$ Hz, 1H), 5.10-5.12 (m, 1H), 4.00 (d, $J = 5.3$ Hz, 1H), 2.41-2.30 (m, 1H), 1.01 (d, $J = 6.9$ Hz, 3H), 1.00 (d, $J = 6.7$ Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 196.4, 148.6, 136.0, 134.2, 128.8 (2C), 128.5, 126.9, 126.8 (2C), 124.5, 81.4, 71.5, 28.4, 19.0, 17.7 ppm. Anal. calcd. for $\text{C}_{16}\text{H}_{18}\text{O}_2$: C, 79.31; H, 7.49. Found: C, 79.20; H, 7.37.

(2*S*,6*R*)-2-Isopropyl-6-phenyl-2*H*-pyran-3(6*H*)-one (10n): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2c** (29.1 mg, 0.19 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (41.2 mg, 0.22 mmol, 1.2 equiv) and TFAA (26 μ L, 0.19 mmol, 1.0 equiv) in DCM anh. (1.1 mL) for 24 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (25.5 mg, 62% yield). ^1H NMR (300 MHz,

CDCl₃): δ 7.40-7.39 (m, 5H), 7.13 (dd, J = 10.4 Hz, J = 3.1 Hz, 1H), 6.20 (dd, J = 10.4 Hz, J = 2.1 Hz, 1H), 5.51-5.49 (m, 1H), 3.80 (d, J = 6.4 Hz, 1H), 2.38-2.27 (m, 1H), 1.03 (d, J = 6.7 Hz, 3H), 0.94 (d, J = 6.9 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.4, 149.0, 137.6, 128.9 (2C), 128.8, 128.0 (2C), 126.7, 82.1, 72.6, 28.1, 18.8, 18.2 ppm. Anal. calcd. for C₁₄H₁₆O₂: C, 77.75; H, 7.46. Found: C, 77.73; H, 7.58.

(2*S*,6*S*)-2-Isopropyl-2-methyl-6-((*E*)-styryl)-2*H*-pyran-3(6*H*)-one (10o): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2d** (29.9 mg, 0.18 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (44.3 mg, 0.21 mmol, 1.2 equiv) and TFAA (25 μ L, 0.18 mmol, 1.0 equiv) in DCM anh. (1.1 mL) for 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (39.2 mg, 85% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.42-7.28 (m, 5H), 6.84 (dd, J = 10.3 Hz, J = 1.5 Hz, 1H), 6.72 (d, J = 15.9 Hz, 1H), 6.20 (dd, J = 15.9 Hz, J = 7.0 Hz, 1H), 6.00 (dd, J = 10.3 Hz, J = 2.5 Hz, 1H), 5.03-5.06 (m, 1H), 2.32-2.23 (m, 1H), 1.26 (s, 3H), 1.07 (d, J = 6.9 Hz, 3H), 0.84 (d, J = 6.7 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 198.7, 148.2, 136.2, 133.1, 128.7 (2C), 128.3, 126.8 (3C), 124.7, 83.6, 70.1, 29.4, 17.3, 17.0, 16.3 ppm. Anal. calcd. for C₁₇H₂₀O₂: C, 79.65; H, 7.86. Found: C, 79.74; H, 7.76.

(2*S*,6*R*)-2-Isopropyl-2-methyl-6-phenyl-2*H*-pyran-3(6*H*)-one (10p): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2d** (29.5 mg, 0.17 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (38.3 mg, 0.21 mmol, 1.2 equiv) and TFAA (24 μ L, 0.17 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 24 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (22.7 mg, 58% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.40-7.36 (m, 5H), 6.87 (dd, J = 10.3 Hz, J = 1.7 Hz, 1H), 6.04 (dd, J = 10.3 Hz, J = 2.6 Hz, 1H), 5.40-5.42 (m, 1H), 2.42-2.32 (m, 1H), 1.25 (s, 3H), 1.09 (d, J = 6.8 Hz, 3H), 0.86 (d, J = 6.8 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 198.7, 149.2, 139.5, 129.0 (2C), 128.7, 127.5 (2C), 124.3, 84.0, 71.6, 29.2, 17.4, 17.0, 16.0 ppm. Anal. calcd. for C₁₅H₁₈O₂: C, 78.23; H, 7.88. Found: C, 78.15; H, 7.93.

(2*S*,6*R*)-2-Isopropyl-2-methyl-6-(phenylethynyl)-2*H*-pyran-3(6*H*)-one (10q): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2d** (30.1 mg, 0.18 mmol, 1.0 equiv), potassium phenylethynyltrifluoroborate **11i** (44.2 mg, 0.28 mmol, 1.2 equiv) and TFAA (25 μ L, 0.18 mmol, 1.0 equiv) in DCM anh. (1.1 mL) at -40°C for 3 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (37.5 mg, 82% yield, *dr* = 85:15). ¹H NMR (300 MHz, CDCl₃): δ 7.48-7.46 (m, 2H), 7.33-7.31 (m, 3H), 6.93 (dd, *J* = 10.2 Hz, *J* = 2.4 Hz, 1H), 6.05 (dd, *J* = 10.2 Hz, *J* = 2.5 Hz, 1H), 5.45 (t, *J* = 2.5 Hz, 1H), 2.22 (heptet, *J* = 6.8 Hz, 1H), 1.37 (s, 3H), 1.03 (d, *J* = 6.8 Hz, 3H), 0.88 (d, *J* = 6.8 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 198.1, 145.7, 132.0 (2C), 129.0, 128.4 (2C), 125.1, 122.1, 86.4, 85.0, 84.4, 60.6, 30.6, 17.9, 17.2, 16.8 ppm. Anal. calcd. for C₁₇H₁₈O₂: C, 80.28; H, 7.13. Found: C, 80.30; H, 7.24.

(2*S*,6*S*)-2-Methyl-2-phenyl-6-((*E*)-styryl)-2*H*-pyran-3(6*H*)-one (10r): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2e** (30.0 mg, 0.15 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (37.1 mg, 0.18 mmol, 1.2 equiv) and TFAA (21 μ L, 0.15 mmol, 1.0 equiv) in DCM anh. (0.9 mL) for 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford the desired compound as a separable mixture of diastereomers (30.9 mg, 72% yield, *dr* = 78:22).

(2*S*,6*S*)-2-Methyl-2-phenyl-6-((*E*)-styryl)-2*H*-pyran-3(6*H*)-one. Major diastereomer. Colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 7.43-7.30 (m, 10H), 6.80-6.71 (m, 2H), 6.28-6.17 (m, 2H), 4.80-4.77 (m, 1H), 1.66 (s, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.4, 149.4, 138.6, 136.2, 133.3, 128.8 (2C), 128.7 (2C), 128.4, 128.3, 126.8 (2C), 126.2, 126.0, 125.8 (2C), 83.1, 71.0, 27.5 ppm. Anal. calcd. for C₂₀H₁₈O₂: C, 82.73; H, 6.25. Found: C, 82.86; H, 6.18.

(2*R*,6*S*)-2-Methyl-2-phenyl-6-((*E*)-styryl)-2*H*-pyran-3(6*H*)-one. Minor diastereomer. Colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 7.57 (d, *J* = 8.0 Hz, 2H), 7.38-7.29 (m, 8H), 7.00 (dd, *J* = 10.3 Hz, *J* = 1.9 Hz, 1H), 6.73 (d, *J* = 16.0 Hz, 1H), 6.21-6.10 (m, 2H), 5.25 (d, *J* = 7.1 Hz, 1H), 1.80 (s, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.8, 148.9, 141.7, 136.2, 133.3, 128.7 (2C), 128.3, 128.1 (2C), 127.9,

126.8 (2C), 126.6 (2C), 126.4, 125.6, 81.9, 71.2, 22.5 ppm. Anal. calcd. for C₂₀H₁₈O₂: C, 82.73; H, 6.25. Found: C, 82.80; H, 6.17.

(2*S*,6*R*)-2-Methyl-2,6-diphenyl-2*H*-pyran-3(6*H*)-one (10*s*): Following general procedure for the synthesis of compounds **10**, Conditions A, starting from **2e** (30.0 mg, 0.15 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (33.1 mg, 0.18 mmol, 1.2 equiv) and TFAA (21 μ L, 0.15 mmol, 1.0 equiv) in DCM anh. (0.9 mL) for 24 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford the desired compound as a separable mixture of diastereomers (22.2 mg, 56% yield, *dr* = 75:25).

(2*S*,6*R*)-2-Methyl-2,6-diphenyl-2*H*-pyran-3(6*H*)-one. Colorless oil Major diastereomer. ¹H NMR (300 MHz, CDCl₃): δ 7.40-7.38 (m, 10H), 6.82 (dd, *J* = 10.3 Hz, *J* = 1.6 Hz, 1H), 6.22 (dd, *J* = 10.4 Hz, *J* = 2.7 Hz, 1H), 5.14-5.13 (m, 1H), 1.67 (s, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.4, 150.3, 139.1, 138.6, 129.0 (2C), 128.8 (2C), 128.7, 128.3, 127.5 (2C), 125.9 (2C), 125.8, 83.4, 72.9, 27.6 ppm. Anal. calcd. for C₁₈H₁₆O₂: C, 81.79; H, 6.10. Found: C, 81.64; H, 6.02.

(2*R*,6*R*)-2-Methyl-2,6-diphenyl-2*H*-pyran-3(6*H*)-one. Colorless oil. Minor diastereomer. ¹H NMR (300 MHz, CDCl₃): δ 7.60 (d, *J* = 7.6 Hz, 2H), 7.49-7.29 (m, 8H), 7.03 (dd, *J* = 10.3 Hz, *J* = 1.6 Hz, 1H), 6.20 (dd, *J* = 10.3 Hz, *J* = 2.4 Hz, 1H), 5.65 (s, 1H), 1.87 (s, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 196.8, 149.9, 141.3, 139.3, 129.1 (2C), 128.7, 128.1 (2C), 127.8, 127.5 (2C), 126.7 (2C), 125.4, 82.2, 72.2, 21.5 ppm. Anal. calcd. for C₁₈H₁₆O₂: C, 81.79; H, 6.10. Found: C, 81.85; H, 6.00.

(*E*)-1-Hydroxy-3-styrylhexane-2,5-dione (13*a*): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2f** (30.0 mg, 0.23 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (59.1 mg, 0.28 mmol, 1.2 equiv) and TFAA (32 μ L, 0.23 mmol, 1.0 equiv) in DCM anh. (1.4 mL) for 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (42.7 mg, 80% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.33-7.27 (m, 5H), 6.56 (d, *J* = 15.8 Hz, 1H), 6.00 (dd, *J* = 15.8 Hz, *J* = 9.3 Hz, 1H), 4.48 (s, 2H), 3.75 (td, *J* = 9.5 Hz, *J* = 3.8 Hz, 1H), 3.29 (dd, *J* = 18.3 Hz, *J* = 9.8 Hz, 1H), 2.92 (bs, 1H), 2.74 (dd, *J* = 18.3 Hz, *J* = 3.8 Hz, 1H), 2.17 (s, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 209.5, 206.3, 136.1, 134.7, 128.8 (2C), 128.4,

126.5 (2C), 124.7, 67.6, 47.4, 45.5, 29.8 ppm. Anal. calcd. for C₁₄H₁₆O₃: C, 72.39; H, 6.94. Found: C, 72.33; H, 6.80.

1-Hydroxy-3-phenylhexane-2,5-dione (13b): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2f** (30.0 mg, 0.23 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (51.5 mg, 0.28 mmol, 1.2 equiv) and TFAA (32 μ L, 0.23 mmol, 1.0 equiv) in DCM anh. (1.4 mL) for 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (21.3 mg, 45% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.33-7.28 (m, 3H), 7.21-7.18 (m 2H), 4.42 (d, J = 18.9 Hz, 1H), 4.23 (d, J = 18.9 Hz, 1H), 4.17 (dd, J = 10.7 Hz, J = 3.6 Hz, 1H), 3.55 (dd, J = 18.3 Hz, J = 10.7 Hz, 1H), 2.86 (bs, 1H), 2.74 (dd, J = 18.3 Hz, J = 3.6 Hz, 1H), 2.17 (s, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 208.9, 206.6, 136.7, 129.4 (2C), 128.2 (2C), 128.1, 67.4, 49.3, 46.9, 29.8 ppm. Anal. calcd. for C₁₂H₁₄O₃: C, 69.88; H, 6.84. Found: C, 69.99; H, 6.95.

(E)-5-Hydroxy-1-phenyl-3-styrylpentane-1,4-dione (13c): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2g** (29.7 mg, 0.16 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (39.4 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μ L, 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (38.6 mg, 82% yield). ¹H NMR (500 MHz, CDCl₃): δ 7.96 (d, J = 8.6 Hz, 2H), 7.59 (t, J = 7.4 Hz, 1H), 7.47 (t, J = 7.9 Hz, 2H), 7.36-7.28 (m, 5H), 6.63 (d, J = 15.8 Hz, 1H), 6.12 (dd, J = 15.8 Hz, J = 9.0 Hz, 1H), 4.59 (q, J = 19.0 Hz, 2H), 3.95 (td, J = 9.5 Hz, J = 3.4 Hz, 1H), 3.86 (dd, J = 17.8 Hz, J = 9.5 Hz, 1H), 3.31 (dd, J = 17.8 Hz, J = 3.4 Hz, 1H), 3.00 (bs, 1H), ppm. ¹³C-NMR (125 MHz, CDCl₃): δ 209.6, 197.7, 136.2, 136.1 134.7, 133.7, 128.9 (4C), 128.4, 128.3 (2C); 126.6 (2C); 125.1, 67.7, 47.5, 41.3 ppm. Anal. calcd. for C₁₉H₁₈O₃: C, 77.53; H, 6.16. Found: C, 77.67; H, 6.02.

5-Hydroxy-1,3-diphenylpentane-1,4-dione (13d): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2g** (29.9 mg, 0.16 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (34.8 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μ L, 0.16 mmol, 1.0 equiv) in DCM

anh. (1.0 mL) 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (34.7 mg, 81% yield). ^1H NMR (300 MHz, CDCl_3): δ 7.95 (d, $J = 8.1$ Hz, 2H), 7.58 (t, $J = 7.5$ Hz, 1H), 7.46 (t, $J = 7.5$ Hz, 2H), 7.36-7.27 (m, 5H), 4.58 (d, $J = 18.8$ Hz, 1H), 4.40-4.30 (m, 2H), 4.11 (dd, $J = 18.0$ Hz, $J = 10.6$ Hz, 1H), 3.32 (dd, $J = 18.0$ Hz, $J = 3.4$ Hz, 1H), 3.17 (bs, 1H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 209.0, 197.9, 136.9, 136.1, 133.7, 129.5 (2C), 128.8 (2C), 128.3 (2C), 128.2 (2C); 128.1, 67.5, 49.4, 42.8 ppm. Anal. calcd. for $\text{C}_{17}\text{H}_{16}\text{O}_3$: C, 76.10; H, 6.01. Found: C, 75.93; H, 5.86.

(E)-6-Hydroxy-4-styrylheptane-2,5-dione (13e): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2i** (29.8 mg, 0.21 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (52.9 mg, 0.25 mmol, 1.2 equiv) and TFAA (29 μL , 0.21 mmol, 1.0 equiv) in DCM anh. (1.3 mL) for 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford the desired compound as a mixture of diastereomers (66:33 mixture, 34.1 mg, 66% yield). Mayor diastereomer. Colorless oil ^1H NMR (300 MHz, CDCl_3): δ 7.32-7.28 (m, 5H), 6.56 (d, $J = 15.9$ Hz, 1H), 5.98 (dd, $J = 15.9$ Hz, $J = 9.4$ Hz, 1H), 4.50 (q, $J = 7.1$, 1H), 3.75 (td, $J = 9.7$ Hz, $J = 4.0$ Hz, 1H), 3.40 (bs, 1H), 3.30 (dd, $J = 18.3$ Hz, $J = 10.0$ Hz, 1H), 2.66 (dd, $J = 18.3$ Hz, $J = 3.8$ Hz, 1H), 2.18 (s, 3H), 1.50 (d, $J = 7.1$ Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 212.3, 206.6, 136.1, 134.7, 128.8 (2C), 128.3, 126.5 (2C), 125.2, 71.8, 45.9, 45.3, 29.9, 20.3 ppm. Anal. calcd. for $\text{C}_{15}\text{H}_{18}\text{O}_3$: C, 73.15; H, 7.37. Found: C, 73.22; H, 7.39.

(E)-1-Hydroxy-4-methyl-3-styrylhexane-2,5-dione (13f): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2h** (30.0 mg, 0.21 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (53.3 mg, 0.25 mmol, 1.2 equiv) and TFAA (29 μL , 0.21 mmol, 1.0 equiv) in DCM anh. (1.3 mL) 1 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford the desired compound as a mixture of diastereomers (90:10 mixture, 41.3 mg, 80% yield). Mayor diastereomer. Colorless oil. ^1H NMR (300 MHz, CDCl_3): δ 7.34-7.29 (m, 5H), 6.60 (d, $J = 15.7$ Hz, 1H), 6.00 (dd, $J = 15.7$ Hz, $J = 10.0$ Hz, 1H), 4.43 (s, 2H), 3.52 (t, $J = 10.0$ Hz, 1H), 3.28-3.17 (m, 1H), 2.86 (bs, 1H), 2.22 (s, 3H), 1.16 (d, $J = 7.5$ Hz, 3H) ppm.

¹³C-NMR (75 MHz, CDCl₃): δ 211.4, 210.2, 136.2, 136.1, 128.9 (2C), 128.4, 126.6 (2C), 124.4, 67.5, 54.9, 48.5, 28.7, 14.9 ppm. Anal. calcd. for C₁₅H₁₈O₃: C, 73.15; H, 7.37. Found: C, 73.19; H, 7.21.

5-Hydroxy-3-(4-methoxyphenyl)-1-phenylpentane-1,4-dione (13g): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2g** (30.1 mg, 0.16 mmol, 1.0 equiv), potassium 4-methoxyphenyltrifluoroborate **11m** (40.7 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μL, 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 3 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (36.3 mg, 76% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.96-7.93 (m, 2H), 7.57 (t, *J* = 7.4 Hz, 1H), 7.46 (t, *J* = 7.8 Hz, 2H), 7.20 (d, *J* = 8.7 Hz, 2H), 6.88 (d, *J* = 8.7 Hz, 2H), 4.55 (d, *J* = 18.7 Hz, 1H), 4.36-4.29 (m, 2H), 4.07 (dd, *J* = 18.2 Hz, *J* = 10.5 Hz, 1H), 3.80 (s, 3H), 3.28 (dd, *J* = 18.2 Hz, *J* = 3.5 Hz, 1H), 2.86 (bs, 1H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 209.2, 198.0, 159.5, 136.2, 133.6, 129.4 (2C), 128.8 (3C), 128.2 (2C), 114.8, 67.3, 55.5, 48.5, 42.8 ppm. Anal. calcd. for C₁₈H₁₈O₄: C, 72.47; H, 6.08. Found: C, 72.61; H, 6.20.

(E)-5-Hydroxy-1-phenyl-3-(3-phenylprop-1-en-1-yl)pentane-1,4-dione (13h): Following general procedure for the synthesis of compounds **13**, Conditions A, starting from **2g** (29.8 mg, 0.16 mmol, 1.0 equiv), potassium *trans*-3-Phenyl-1-propen-1-yltrifluoroborate **11e** (42.6 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μL, 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 3 h. The crude product was purified by flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (38.0 mg, 77% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.93 (d, *J* = 7.6 Hz, 2H), 7.58 (t, *J* = 7.4 Hz, 1H), 7.46 (t, *J* = 7.6 Hz, 2H), 7.32-7.21 (m, 3H), 7.14 (d, *J* = 7.4 Hz, 2H), 5.94-5.84 (m, 1H), 5.46 (dd, *J* = 15.3 Hz, *J* = 7.8 Hz, 1H), 4.59 (s, 2H), 3.81-3.70 (m, 2H), 3.37 (d, *J* = 6.6 Hz, 2H), 3.25-3.16 (m, 1H), 2.98 (bs, 1H), ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 209.8, 197.8, 139.4, 136.2, 134.9, 133.7, 128.8 (2C), 128.7 (2C), 128.6 (2C), 128.2 (2C), 127.0, 126.5, 67.5, 47.2, 41.1, 39.0 ppm. Anal. calcd. for C₂₀H₂₀O₃: C, 77.90; H, 6.54. Found: C, 78.03; H, 6.49.

(E)-(1-Benzyl-4-styryl-1H-pyrrol-2-yl)methanol (14a): Following general procedure for the synthesis of pyrroles **14**, starting from **2j** (35.0 mg, 0.31 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (96.8 mg, 0.46 mmol, 1.5 equiv), tetrabutylammonium hydrogensulfate (52.6

mg, 0.16 mmol, 0.5 equiv) in DCM anh. (6.2 mL) at rt for 48h. Then, benzylamine (34 μ L, 0.31 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (5.9 mg, 0.03 mmol, 0.1 equiv) in toluene (1.6 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (67.3 mg, 75% yield). ^1H NMR (500 MHz, acetone- d_6): δ 7.44 (d, J = 8.2 Hz, 2H), 7.33 (t, J = 7.6 Hz, 2H), 7.29 (t, J = 8.1 Hz, 3H), 7.18 (d, J = 7.8 Hz, 2H), 7.14 (t, J = 7.5 Hz, 1H), 7.05 (d, J = 16.3 Hz, 1H), 6.92 (d, J = 1.8 Hz, 1H), 6.76 (d, J = 16.3 Hz, 1H), 6.36 (d, J = 1.8 Hz, 1H), 5.23 (s, 2H), 4.47 (s, 2H), 2.81 (bs, 1H) ppm. ^{13}C -NMR (125 MHz, acetone- d_6): δ 139.8, 139.6, 135.2, 129.4 (2C), 129.3 (2C), 128.1, 127.8 (2C), 127.0, 126.4 (2C), 124.1, 123.7, 123.4, 122.6, 106.5, 56.7, 50.9 ppm. Anal. calcd. for $\text{C}_{20}\text{H}_{19}\text{NO}$: C, 83.01; H, 6.62. Found: C, 82.90; H, 6.74.

(*E*)-1-(1-Benzyl-4-styryl-1*H*-pyrrol-2-yl)ethanol (14b): Following general procedure for the synthesis of pyrroles **14**, starting from **2a** (29.7 mg, 0.23 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (73.8 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogenosulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) at rt for 48h. Then, benzyl amine (26 μ L, 0.23 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (4.4 mg, 0.02 mmol, 0.1 equiv) in toluene (1.2 mL) at rt for 1 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (57.5 mg, 81% yield). ^1H NMR (300 MHz, acetone- d_6): δ 7.42-7.47 (m, 2H), 7.22-7.37 (m, 6H), 7.12-7.18 (m, 2H), 7.06 (d, J = 16.2 Hz, 1H), 6.88 (d, J = 1.7 Hz, 1H), 6.77 (d, J = 16.2 Hz, 1H), 6.40 (d, J = 1.7 Hz, 1H), 5.36 (d, J = 16.0 Hz, 1H), 5.20 (d, J = 16.0 Hz, 1H), 4.70 (q, J = 6.3 Hz, 1H), 3.91 (d, J = 6.3 Hz, 1H), 1.48 (d, J = 6.3 Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 140.0, 139.6, 139.0, 129.4, 129.3, 128.0, 127.7, 126.9, 126.3, 124.0, 123.8, 123.2, 122.3, 103.6, 62.2, 50.9, 22.9 ppm. Anal. calcd. for $\text{C}_{21}\text{H}_{21}\text{NO}$: C, 83.13; H, 6.98. Found: C, 83.09; H, 6.99.

(*E*)-1-(1-Benzyl-4-(3-phenylprop-1-en-1-yl)-1*H*-pyrrol-2-yl)ethanol (14c): Following general procedure for the synthesis of pyrroles **14**, starting from **2a** (30.3 mg, 0.23 mmol, 1.0 equiv), *trans*-3-phenyl-1-propen-1-yltrifluoroborate **11e** (78.5 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogenosulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) rt for 48h. Then, benzyl amine (26 μ L, 0.23 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (4.4 mg, 0.02 mmol, 0.1 equiv) in toluene (1.2

mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (57.9 mg, 78% yield). ¹H NMR (300 MHz, acetone-d₆): δ 7.06-7.38 (m, 10H), 6.69 (d, *J* = 1.9 Hz, 1H), 6.30 (d, *J* = 15.5 Hz, 1H), 6.21 (d, *J* = 1.9 Hz, 1H), 5.94 (dt, *J* = 15.5 Hz, *J* = 7.1 Hz, 1H), 5.32 (d, *J* = 16.1 Hz, 1H), 5.14 (d, *J* = 16.1 Hz, 1H), 4.68 (q, *J* = 6.5 Hz, 1H), 3.83 (d, *J* = 6.8 Hz, 1H), 3.44 (d, *J* = 6.8 Hz, 2H), 1.43 (d, *J* = 6.5 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 142.3, 140.2, 138.5, 129.4, 129.2, 128.0, 127.7, 126.7, 125.6, 124.3, 122.2, 121.8, 103.6, 62.3, 50.8, 40.0, 23.0 ppm. Anal. calcd. for C₂₂H₂₃NO: C, 83.24; H, 7.30. Found: C, 83.31; H, 7.28.

(1-Benzyl-4-(phenylethynyl)-1*H*-pyrrol-2-yl)methanol (14d): Following general procedure for the synthesis of pyrroles **14**, starting from **2j** (36.0 mg, 0.31 mmol, 1.0 equiv), potassium phenylethynyltrifluoroborate **11i** (95.7 mg, 0.46 mmol, 1.5 equiv) and tetrabutylammonium hydrogenosulfate (52.6 mg, 0.16 mmol, 0.5 equiv) in DCM anh. (6.2 mL) at rt for 48h. Then, benzylamine (34 μL, 0.31 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (5.9 mg, 0.03 mmol, 0.1 equiv) in toluene (1.6 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (57.0 mg, 64% yield). ¹H NMR (300 MHz, acetone-d₆): δ 7.45-7.29 (m, 8H), 7.20 (d, *J* = 7.5 Hz, 2H), 7.09 (d, *J* = 1.8 Hz, 1H), 6.22 (d, *J* = 1.8 Hz, 1H), 5.28 (s, 2H), 4.48-4.47 (m, 2H), 4.08 (bs, 1H) ppm. ¹³C-NMR (75 MHz, acetone-d₆): δ 139.2, 134.4, 131.7 (2C), 129.4 (2C), 129.2 (2C), 128.3, 128.2, 127.8 (2C), 127.5, 125.4, 112.2, 103.6, 87.9, 86.5, 56.3, 51.0 ppm. Anal. calcd. for C₂₀H₁₇NO: C, 83.59; H, 5.96. Found: C, 83.76; H, 6.08.

1-(1-Benzyl-4-(phenylethynyl)-1*H*-pyrrol-2-yl)ethanol (14e): Following general procedure for the synthesis of pyrroles **14**, starting from **2a** (30.3 mg, 0.23 mmol, 1.0 equiv), potassium phenylethynyltrifluoroborate **11i** (73.1 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogenosulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) at rt for 48h. Then, benzyl amine (26 μL, 0.23 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (4.4 mg, 0.02 mmol, 0.1 equiv) in toluene (1.2 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (57.2 mg, 81% yield). ¹H NMR (300 MHz, acetone-d₆): δ

7.25-7.46 (m, 8H), 7.17 (d, $J = 7.4$ Hz, 2H), 7.05 (d, $J = 1.7$ Hz, 1H), 6.23 (d, $J = 1.0$ Hz, 1H), 5.41 (d, $J = 15.8$ Hz, 1H), 5.25 (d, $J = 15.8$ Hz, 1H), 4.73 (q, $J = 6.5$ Hz, 1H), 4.04 (d, $J = 6.5$ Hz, 1H), 1.43 (d, $J = 6.5$ Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 139.4, 138.2, 131.7, 129.4, 129.3, 128.2, 128.1, 127.7, 127.2, 125.4, 109.6, 103.4, 87.9, 86.6, 62.0, 51.1, 22.8 ppm. Anal. calcd. for $\text{C}_{21}\text{H}_{19}\text{NO}$: C, 83.69; H, 6.35. Found: C, 83.60; H, 6.34.

(*E*)-(1-Benzyl-5-phenyl-3-styryl-1*H*-pyrrol-2-yl)methanol (15a): Following general procedure for the synthesis of pyrroles **15**, starting from **2g** (29.7 mg, 0.16 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (39.4 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μL , 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 1 h. Then, benzylamine (17 μL , 0.16 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (3.0 mg, 0.02 mmol, 0.1 equiv) in toluene (0.8 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (42.1 mg, 72% yield). ^1H NMR (300 MHz, acetone- d_6): δ 7.55-7.53 (m, 2H), 7.35-7.26 (m, 12H), 6.98-6.96 (m, 3H), 6.61 (s, 1H), 5.41 (s, 2H), 4.62 (s, 2H), 4.07 (bs, 1H) ppm. ^{13}C -NMR (75 MHz, acetone- d_6): δ 140.3, 139.6, 137.1, 134.0, 133.7, 129.6 (2C), 129.5 (2C), 129.4 (2C), 129.3 (2C), 128.2, 127.8, 127.2, 126.6 (2C), 126.4 (2C), 125.4, 122.2, 122.0, 106.1, 53.8, 48.2 ppm. Anal. calcd. for $\text{C}_{26}\text{H}_{23}\text{NO}$: C, 85.45; H, 6.34. Found: C, 85.35; H, 6.48.

(*E*)-(1-Benzyl-5-phenyl-3-(3-phenylprop-1-en-1-yl)-1*H*-pyrrol-2-yl)methanol (15b): Following general procedure for the synthesis of pyrroles **15**, starting from **2g** (29.8 mg, 0.16 mmol, 1.0 equiv), potassium *trans*-3-phenyl-1-propen-1-yltrifluoroborate **11e** (42.6 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μL , 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 3 h. Then, benzylamine (17 μL , 0.16 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (3.0 mg, 0.02 mmol, 0.1 equiv) in toluene (0.8 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (46.7 mg, 77% yield). ^1H NMR (500 MHz, acetone- d_6): δ 7.32-7.26 (m, 11H), 7.22-7.19 (m, 2H), 6.92 (d, $J = 7.3$ Hz, 2H), 6.56 (d, $J = 15.6$ Hz, 1H), 6.40 (s, 1H), 6.11 (dt, $J = 15.6$ Hz, $J = 7.1$ Hz, 1H), 5.36 (s, 2H), 4.49 (d, $J = 5.3$ Hz, 2H), 3.93 (t, $J = 5.3$ Hz, 1H), 3.52 (d, $J = 7.1$ Hz, 2H) ppm.

¹³C-NMR (125 MHz, acetone-d₆): δ 142.1, 140.5, 136.6, 134.2, 132.1, 129.6 (2C), 129.5 (2C), 129.4 (2C), 129.3 (2C), 129.2 (2C), 128.0, 127.7, 126.7, 126.4 (2C), 125.9, 124.0, 122.1, 106.3, 53.9, 48.1, 40.2 ppm. Anal. calcd. for C₂₇H₂₅NO: C, 85.45; H, 6.64. Found: C, 85.31; H, 6.75.

(E)-(1-Benzyl-5-methyl-3-styryl-1H-pyrrol-2-yl)methanol (15c): Following general procedure for the synthesis of pyrroles **15**, starting from **2f** (30.5 mg, 0.24 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (65.0 mg, 0.29 mmol, 1.2 equiv) and TFAA (33 μL, 0.24 mmol, 1.0 equiv) in DCM anh. (1.4 mL) for 1 h. Then, benzylamine (26 μL, 0.24 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (4.6 mg, 0.02 mmol, 0.1 equiv) in toluene (1.2 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (36.4 mg, 50% yield). ¹H NMR (300 MHz, acetone-d₆): δ 7.50 (d, *J* = 7.6 Hz, 2H), 7.34-7.13 (m, 8H), 7.01 (d, *J* = 7.5 Hz, 2H), 6.78 (d, *J* = 16.1 Hz, 1H), 6.23 (s, 1H), 5.28 (s, 2H), 4.62-4.61 (m, 2H), 3.92 (t, *J* = 5.4 Hz, 1H), 2.11 (s, 3H) ppm. ¹³C-NMR (75 MHz, acetone-d₆): δ 139.9, 139.8, 132.2, 131.2, 129.4 (2C), 129.3 (2C), 127.8, 126.9, 126.7 (2C), 126.4 (2C), 124.2, 122.5, 120.9, 104.0, 53.5, 47.4, 12.3 ppm. Anal. calcd. for C₂₁H₂₁NO: C, 83.13; H, 6.98. Found: C, 83.29; H, 6.90.

(1-Benzyl-3,5-diphenyl-1H-pyrrol-2-yl)methanol (15d): Following general procedure for the synthesis of pyrroles **15**, starting from **2g** (30.3 mg, 0.16 mmol, 1.0 equiv), potassium phenyltrifluoroborate **11f** (34.8 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μL, 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 1 h. Then, benzylamine (17 μL, 0.16 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (3.0 mg, 0.02 mmol, 0.1 equiv) in toluene (0.8 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (38.0 mg, 70% yield). ¹H NMR (300 MHz, acetone-d₆): δ 7.62-7.58 (m, 2H), 7.42-7.22 (m, 11H), 6.98 (d, *J* = 7.1 Hz, 2H), 6.44 (s, 1H), 5.47 (s, 2H), 4.55-4.53 (m, 2H), 4.07 (t, *J* = 4.8 Hz, 1H) ppm. ¹³C-NMR (75 MHz, acetone-d₆): δ 140.4, 137.5, 136.1, 134.2, 130.8, 129.6 (2C), 129.5 (2C), 129.3 (2C), 129.2 (2C), 129.0 (2C), 128.1, 127.8, 126.5, 126.4 (2C), 125.8, 109.4, 54.7, 48.3 ppm. Anal. calcd. for C₂₄H₂₁NO: C, 84.92; H, 6.24. Found: C, 85.11; H, 6.15.

(1-Benzyl-3-(4-methoxyphenyl)-5-phenyl-1*H*-pyrrol-2-yl)methanol (15e): Following general procedure for the synthesis of pyrroles **15**, starting from **2g** (30.4 mg, 0.16 mmol, 1.0 equiv), potassium 4-methoxyphenyltrifluoroborate **11m** (40.7 mg, 0.19 mmol, 1.2 equiv) and TFAA (22 μ L, 0.16 mmol, 1.0 equiv) in DCM anh. (1.0 mL) for 3 h. Then, benzylamine (17 μ L, 0.16 mmol, 1.0 equiv) and *p*-toluenesulfonic acid (3.0 mg, 0.02 mmol, 0.1 equiv) in toluene (0.8 mL) at rt for 12 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 (47.8 mg, 81% yield). ^1H NMR (300 MHz, acetone- d_6): δ 7.52 (d, J = 8.9 Hz, 2H), 7.37-7.22 (m, 8H), 6.99-6.96 (m, 4H), 6.39 (s, 1H), 5.44 (s, 2H), 4.51 (s, 2H), 4.06 (bs, 1H), 3.81 (s, 3H) ppm. ^{13}C -NMR (75 MHz, acetone- d_6): 159.1, 140.5, 139.5, 134.3, 130.4, 130.0 (2C), 129.9, 129.6 (2C), 129.4 (2C), 129.3 (2C), 128.0, 127.8, 126.4 (2C), 125.6, 114.7 (2C), 109.3, 55.5, 54.9, 48.3 δ ppm. Anal. calcd. for $\text{C}_{25}\text{H}_{23}\text{NO}_2$: C, 81.27; H, 6.27. Found: C, 81.22; H, 6.38.

(2*S*,3*R*,6*R*)-6-Methyl-5-oxo-3-((*E*)-styryl)tetrahydro-2*H*-pyran-2-yl acetate (16a): Following general procedure for the synthesis of compounds **16**, Conditions C, starting from **2a** (29.4 mg, 0.23 mmol, 1.0 equiv), potassium *trans*-styryltrifluoroborate **11a** (73.8 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogensulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) at rt for 48h. Then, Ac_2O (22 μ L, 0.23 mmol, 1.0 equiv), DMAP (1.4 mg, 0.012 mmol, 5% mmol) and Et_3N (65 μ L, 0.47 mmol, 2.0 equiv) at 0°C for 1 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (52.6 mg, 82% yield, *dr* 90:10). ^1H NMR (300 MHz, CDCl_3): δ 7.14-7.34 (m, 5H), 6.42 (d, J = 15.8 Hz, 1H), 6.09 (d, J = 5.4 Hz, 1H), 6.03 (dd, J = 15.8 Hz, J = 7.7 Hz, 1H), 4.31 (q, J = 6.8 Hz, 1H), 2.93-3.06 (m, 1H), 2.51-2.69 (m, 2H), 2.08 (s, 3H), 1.27 (d, J = 6.8 Hz, 3H) ppm. ^{13}C -NMR (75 MHz, CDCl_3): δ 208.9, 169.9, 136.4, 133.0, 128.8, 128.2, 126.6, 126.5, 94.4, 73.2, 42.2, 39.5, 21.3, 15.1 ppm. Anal. calcd. for $\text{C}_{16}\text{H}_{18}\text{O}_4$: C, 70.06; H, 6.61. Found: C, 70.14; H, 6.59.

(2*S*,3*R*,6*R*)-6-Methyl-3-(2-methylprop-1-en-1-yl)-5-oxotetrahydro-2*H*-pyran-2-yl acetate (16b): Following general procedure for the synthesis of compounds **16**, Conditions C, starting from **2a**

(29.9 mg, 0.23 mmol, 1.0 equiv), potassium 2-methyl-1-propenyltrifluoroborate **11k** (56.9 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogensulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) at rt for 48h. Then, Ac₂O (22 μ L, 0.23 mmol, 1.0 equiv), DMAP (1.4 mg, 0.012 mmol, 5% mmol) and Et₃N (65 μ L, 0.47 mmol, 2.0 equiv) at 0°C for 1 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (33.9 mg, 64% yield, *dr* 95:5). ¹H NMR (300 MHz, CDCl₃): δ 5.97 (d, *J* = 5.3 Hz, 1H), 5.02 (dt, *J* = 9.4 Hz, *J* = 1.3 Hz, 1H), 4.33 (c. *J* = 6.8 Hz, 1H), 3.02-3.13 (m, 1H), 2.36-2.57 (m, 2H), 2.12 (s, 3H), 1.71 (s, 3H), 1.64 (s, 3H), 1.31 (d, *J* = 6.8 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 209.7, 170.0, 136.4, 121.8, 95.2, 73.0, 40.1, 37.9, 25.9, 21.4, 18.3, 15.1 ppm. Anal. calcd. for C₁₂H₁₈O₄: C, 63.70; H, 8.02. Found: C, 63.75; H, 8.03.

(2*S*,3*R*,6*R*)-3-(Cyclohex-1-en-1-yl)-6-methyl-5-oxotetrahydro-2*H*-pyran-2-yl acetate (16c):

Following general procedure for the synthesis of compounds **16**, Conditions C, starting from **2a** (29.5 mg, 0.23 mmol, 1.0 equiv), potassium cyclohexenyltrifluoroborate **11l** (66.0 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogensulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) at rt for 18h. Then, Ac₂O (22 μ L, 0.23 mmol, 1.0 equiv), DMAP (1.4 mg, 0.012 mmol, 5% mmol) and Et₃N (65 μ L, 0.47 mmol, 2.0 equiv) at 0°C for 1 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (31.3 mg, 53% yield, *dr* 89:11). ¹H NMR (300 MHz, CDCl₃): δ 6.12 (d, *J* = 5.4 Hz, 1H), 5.55 (s, 1H), 4.28 (c. *J* = 6.6 Hz, 1H), 2.41-2.70 (m, 3H), 2.11 (s, 3H), 1.90-2.05 (m, 4H), 1.51-1.65 (m, 4H), 1.31 (d, *J* = 6.6 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 210.9, 170.1, 134.4, 125.5, 94.0, 72.6, 45.6, 38.5, 26.4, 25.3, 22.8, 22.2, 21.4, 15.3 ppm. Anal. calcd. for C₁₄H₂₀O₄: 66.65; H, 7.99. Found: 66.59; H, 8.01.

(2*S*,3*R*,6*R*)-3-(Benzofuran-2-yl)-6-methyl-5-oxotetrahydro-2*H*-pyran-2-yl acetate (16d): Fol-

lowing general procedure for the synthesis of compounds **16**, Conditions C, starting from **2a** (29.4 mg, 0.23 mmol, 1.0 equiv), potassium benzofuran-2-trifluoroborate **11m** (78.7 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogensulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) at rt

for 24h. Then, Ac₂O (22 μL, 0.23 mmol, 1.0 equiv), DMAP (1.4 mg, 0.012 mmol, 5% mmol) and Et₃N (65 μL, 0.47 mmol, 2.0 equiv) at 0°C for 1 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a yellow oil (53.3 mg, 79% yield, *dr* 92:8). ¹H NMR (300 MHz, CDCl₃): δ 7.49-7.55 (m, 1H), 7.41-7.47 (m, 1H), 7.18-7.31 (m, 2H), 6.47-6.58 (m, 2H), 4.43 (c. *J* = 6.6 Hz, 1H), 3.64-3.78 (m, 1H), 2.92-3.03 (m, 2H), 2.17 (s, 3H), 1.35 (d, *J* = 6.6 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 207.6, 169.6, 155.0, 154.7, 128.1, 124.5, 123.1, 121.1, 111.3, 104.2, 92.5, 73.3, 39.3, 38.0, 21.2, 15.2 ppm. Anal. calcd. for C₁₆H₁₆O₅: C, 66.66; H, 5.59. Found: C, 66.58; H, 5.60.

(2*S*,3*R*,6*R*)-6-Methyl-5-oxo-3-(phenylethynyl)tetrahydro-2*H*-pyran-2-yl acetate (16e): Following general procedure for the synthesis of compounds **16**, Conditions C, starting from **2a** (29.8 mg, 0.23 mmol, 1.0 equiv), potassium phenylethynyltrifluoroborate **11i** (73.1 mg, 0.35 mmol, 1.5 equiv) and tetrabutylammonium hydrogensulfate (39.7 mg, 0.12 mmol, 0.5 equiv) in DCM anh. (4.6 mL) at rt for 48h. Then, Ac₂O (22 μL, 0.23 mmol, 1.0 equiv), DMAP (1.4 mg, 0.012 mmol, 5% mmol) and Et₃N (65 μL, 0.47 mmol, 2.0 equiv) at 0°C for 1 h. The crude product was purified via flash column chromatography on silica gel eluting with Hexane:AcOEt 8:2 to afford a colorless oil (47.4 mg, 75% yield, *dr* 98:2). ¹H NMR (300 MHz, CDCl₃): δ 7.23-7.46 (m, 5H), 6.36 (d, *J* = 4.6 Hz, 1H), 4.36 (q, *J* = 6.9 Hz, 1H), 3.37-3.45 (m, 1H), 2.89 (dd, *J* = 16.0 Hz, *J* = 5.4 Hz, 1H), 2.78 (dd, *J* = 16.0 Hz, *J* = 8.0 Hz, 1H), 2.18 (s, 3H), 1.38 (d, *J* = 6.5 Hz, 3H) ppm. ¹³C-NMR (75 MHz, CDCl₃): δ 206.9, 169.6, 131.8, 128.7, 128.4, 122.4, 93.0, 85.5, 85.1, 73.3, 39.5, 32.8, 21.3, 15.3 ppm. Anal. calcd. for C₁₆H₁₆O₄: C, 70.57; H, 5.92. Found: C, 70.50; H, 5.93.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI:

Copies of ¹H NMR and ¹³C NMR spectra, and NOE determinations.

AUTHOR INFORMATION

Corresponding Author

*csaky@ucm.es

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

This work was supported by projects CTQ2014-52213-R from the Spanish government (MICINN).

REFERENCES

(1) See for example: (a) Cavill, G. W. K.; Laing, D. G.; Williams, P. J. Synthesis of the Juvenile Hormone Methyl *cis*-10,11-Epoxy-7-ethyl-3,11-dimethyltrideca-*trans,trans*-2,6-dienoate. *Aust. J. Chem.* **1969**, 22, 2145–2160. (b) Achmatowicz, O.; Bukowski, P.; Szechner, B.; Zwierzchowska, Z.; Zamojski, A. Synthesis of Methyl 2,3-dideoxy-*DL*-alk-2-enopyranosides from Furan Compounds. General Approach to the Total Synthesis of Monosaccharides. *Tetrahedron* **1971**, 27, 1973-1996. (c) Ghosh, A. K.; Brindisi, M. Achmatowicz Reaction and its Application in the Syntheses of Bioactive Molecules. *RSC Adv.* **2016**, 6, 111564-111598. (d) Deska, J.; Thiel, D.; Gianolio, E. The Achmatowicz Rearrangement - Oxidative Ring Expansion of Furfuryl Alcohols. *Synthesis* **2015**, 47, 3435-3450. (e) Plutschack, M. B.; Seeberger, P. H.; Gilmore, K. Visible-light-mediated Achmatowicz Rearrangement. *Org. Lett.* **2017**, 19, 30-33, and references cited therein.

(2) Although all material in this paper was racemic, the AR products can be obtained optically pure by dynamic kinetic resolution. See for example: (a) Wang, H. -Y.; Yang, K.; Yin, D.; Liu, C.; Glazier, D. A.; Tang, W. Chiral Catalyst-Directed Dynamic Kinetic Diastereoselective Acylation of Lactols for De Novo Synthesis of Carbohydrate. *Org. Lett.* **2015**, 17, 5272-5275. (b) Zhao, C.; Li, F.; Wang, J. N-Heterocyclic Carbene Catalyzed Dynamic Kinetic Resolution of Pyranones. *Angew. Chem. Int. Ed.* **2016**, 55, 1820-1824.

(3) See ref 1, and in addition: (a) Aljahdali, A. Z.; Shi, P.; Zhong, Y.; O'Doherty, G. A. De Novo Asymmetric Synthesis of the Pyranoses: From Monosaccharides to Oligosaccharides. *Adv. Carb. Chem. Biochem.* **2013**, *69*, 55-123. (b) Palfram, M. J.; Pattenden, G. The Versatility of Furfuryl Alcohols and Furanoxonium Ions in Synthesis. *Chem. Commun.* **2014**, *50*, 7223-7242. (c) Song, W.; Wang, S.; Tang, W. De Novo Synthesis of Mono- and Oligosaccharides via Dihydropyran Intermediates. *Chem. Asian J.* **2017**, *12*, 1027-1042. (d) Vetica, F.; Chauhan, P.; Dochain, S.; Enders, D. Asymmetric Organocatalytic Methods for the Synthesis of Tetrahydropyrans and their Application in Total Synthesis. *Chem. Soc. Rev.* **2017**, *46*, 1661-1674.

(4) (a) Comely, A. C.; Eelkema, R.; Minnaard, A. J.; Feringa, B. L. De Novo Asymmetric Bio- and Chemo-Catalytic Synthesis of Saccharides - Stereoselective Formal O-Glycoside Bond Formation Using Palladium Catalysis. *J. Am. Chem. Soc.* **2003**, *125*, 8714-8715. (b) Babu, R. S.; O'Doherty, G. A. A Palladium-Catalyzed Glycosylation Reaction: The de Novo Synthesis of Natural and Unnatural Glycosides. *J. Am. Chem. Soc.* **2003**, *125*, 12406-12407. (c) Babu, R. S.; Zhou, M.; O'Doherty, G. A. De Novo Synthesis of Oligosaccharides Using a Palladium-Catalyzed Glycosylation Reaction. *J. Am. Chem. Soc.* **2004**, *126*, 3428-3429.

(5) (a) Taniguchi, T.; Ohnishi, H.; Ogasawara, H. An Expedient Preparation of Chiral Building Blocks having Levoglucosenone Chromophore: A New Enantiocontrolled Route to (-)- β -Multistriatin And (+)-exo-Brevicomine. *Chem. Commun.* **1996**, 1477-1478. (b) Takeuchi, M.; Taniguchi, T.; Ogasawara, K. Back to the sugars. A New Enantio- and Diastereocontrolled Route to Hexoses from Furfural. *Synthesis* **1999**, 341-354. (c) Ren, J.; Liu, Y.; Song, L.; Tong, R. Scalable Asymmetric Total Syntheses of (+)-Psoracorylifol B and (+)-ent-Psoracorylifol C. *Org. Lett.* **2014**, *16*, 2986-2989. (d) Ren, J.; Tong, R. Asymmetric Total Synthesis of (+)-Didemniserinolipid B via Achmatowicz Rearrangement / Bicyclopentaketone ketalization. *J. Org. Chem.* **2014**, *79*, 6987-6995.

(6) (a) Burns, N. Z.; Witten, M. R.; Jacobsen, E. N. Dual Catalysis in Enantioselective Oxidopyrylium-Based [5 + 2] Cycloadditions. *J. Am. Chem. Soc.* **2011**, *133*, 14578-14581. (b) Pellissier, H. Recent Developments in the [5+2] Cycloaddition. *Adv. Synth. Catal.* **2011**, *353*, 189-218. (c) Ylijoki,

1 K. E. O.; Stryker, J. M. [5 + 2] Cycloaddition Reactions in Organic and Natural Product Synthesis.
2 *Chem. Rev.* **2013**, *113*, 2244–2266. (d) Witten, M. R.; Jacobsen, E. N. Catalytic Asymmetric Synthesis
3 of 8-Oxabicyclooctanes by Intermolecular [5+2] Pyrylium Cycloadditions. *Angew. Chem. Int. Ed.* **2014**,
4 *53*, 5912–5916.
5
6
7

8
9 (7) Jones, R. A.; Krische, M. J. Asymmetric Total Synthesis of the Iridoid β -Glucoside (+)-Geniposide
10 via Phosphine Organo-Catalysis. *Org. Lett.* **2009**, *11*, 1849-1851.
11
12

13
14 (8) Yu, J.; Ma, H.; Yao, H.; Cheng, H. ; Tong, R. Diastereoselective and Regiodivergent Oxa-[3 +
15 2]Cycloaddition of Achmatowicz Products and Cyclic 1,3-Dicarbonyl Compounds. *Org. Chem. Front.*
16 **2016**, *3*, 714-719.
17
18
19

20
21 (9) (a) Lewis, M. D.; Cha, J. K.; Kishi, Y. Highly Stereoselective Approaches to α - and β -C-
22 Glycopyranosides. *J. Am. Chem. Soc.* **1982**, *104*, 4976–4978. (b) Zhu, L.; Song, L.; Tong, R. Diastere-
23 oselective Reductive Ring Expansion of Spiroketal Dihydropyranones to cis-Fused Bicyclic Ethers.
24 *Org. Lett.* **2012**, *14*, 5892-5895.
25
26
27

28
29 (10) (a) Li, Z.; Ip, F. C. F.; Ip, N. Y.; Tong, R. Highly trans-Selective Arylation of Achmatowicz Rear-
30 rangement Products by Reductive γ -Deoxygenation and Heck-Matsuda Reaction: Asymmetric Total
31 Synthesis of (-)-Musellarins A-C and Their Analogues. *Chem. Eur. J.* **2015**, *21*, 11152-11157. (b) Li, Z.;
32 Tong, R. Asymmetric Total Syntheses of (-)-Hedycoropyrans A and B. *J. Org. Chem.* **2017**, *82*, 1127-
33 1135.
34
35
36

37
38 (11) O'Connor, C. J.; Beckmann, H. S. G.; Spring, D. R. Diversity-oriented Synthesis: Producing
39 Chemical Tools for Dissecting Biology. *Chem. Soc. Rev.* **2012**, *41*, 4444-4456.
40
41
42

43
44 (12) Molander, G. A. Organotrifluoroborates: Another Branch of the Mighty Oak. *J. Org. Chem.* **2015**,
45 *80*, 7837-7848.
46
47
48

49
50 (13) For a review on C-C bond forming reactions of boronic acid derivatives under transition-metal-
51 free conditions, see: Roscales, S.; Csáky, A. G. Transition-metal-free C-C Bond Forming Reactions of
52 Aryl, Alkenyl and Alkynylboronic Acids and their Derivatives. *Chem. Soc. Rev.* **2014**, *43*, 8215-8225.
53
54
55
56
57
58
59
60

(14) For reviews on conjugate addition reactions of boronic acid derivatives under transition-metal-free conditions, see: (a) Sanchez-Sancho, F.; Csáký, A. G. C-C Bond Formation with Boronic Acids and Derivatives by Transition-Metal-Free Conjugate Addition Reactions. *Synthesis* **2016**, *48*, 2165-2177. (b) Nguyen, T. N.; May, J. A. Enantioselective Organocatalytic Conjugate Addition of Organoboron Nucleophiles. *Tetrahedron Lett.* **2017**, *58*, 1535-1544.

(15) Due to their interest in the preparation of cyclopentenones, 5-membered heterocycles and 6-membered heterocycles, the synthesis of 1,4-diketones has received considerable attention in recent literature. For some recent reports, see: (a) Setzer, P.; Beauseigneur, A.; Pearson-Long, M. S. M.; Bertus, P. Titanium-Mediated Synthesis of 1,4-Diketones from Grignard Reagents and Acyl Cyanohydrins. *Angew. Chem. Int. Ed.* **2010**, *49*, 8691-8694. (b) Parida, B. B.; Das, P. P.; Niocel, M.; Cha, J. K. C-Acylation of Cyclopropanols: Preparation of Functionalized 1,4-Diketones. *Org. Lett.* **2013**, *15*, 1780-1783. (c) Zhang, F.; Du, P.; Chen, J.; Wang, H.; Luo, Q.; Wan, X. Co-Catalyzed Synthesis of 1,4-Dicarbonyl Compounds Using TBHP Oxidant. *Org. Lett.* **2014**, *16*, 1932-1935. (d) Luo, J.; Jiang, Q.; Chen, H.; Tang, Q. Catalyst-Free Formation of 1,4-Diketones by Addition of Silyl Enolates to Oxyallyl Zwitterions In Situ Generated From α -Haloketones. *RSC Adv.* **2015**, *5*, 67901-67908. (e) Stepherson, J. R.; Fronczek, F. R.; Kartika, R. An Expedient Synthesis of Functionalized 1,4-Diketone-Derived Compounds via Silyloxyallyl Cation Intermediates. *Chem. Commun.* **2016**, *52*, 2300-2303. (f) Yin, H.; Nielsen, D. U.; Johansen, M. K.; Lindhardt, A. T.; Skrydstrup, T. Development of a Palladium-Catalyzed Carbonylative Coupling Strategy to 1,4-Diketones. *ACS Catal.* **2016**, *6*, 2982-2987.

(16) (a) Khaghaninejad, S.; Heravi, M. M. Paal-Knorr Reaction in the Synthesis of Heterocyclic Compounds. *Adv. Het. Chem.* **2014**, *III*, 95-146. (b) Vessally, E. A New Avenue to The Synthesis of Highly Substituted Pyrroles: Synthesis from N-Propargylamines. *RSC Adv.* **2016**, *6*, 18619-18631.

(17) (a) Hobson, S. J.; Marquez, R. Stereocontrolled Synthesis of Highly Functionalized Spirocyclic Pyrans. *Org. Biomol. Chem.* **2006**, *4*, 3808-3814. (b) Srihari, P.; Sridhar, Y. Total Synthesis of Both Enantiomers of Macrocyclic Lactone Aspergillide C. *Eur. J. Org. Chem.* **2011**, *2011*, 6690-6697.

(18) For representative examples, see: (a) Yunker, M. B.; Plaumann, D. E.; Bert Fraser-Reid, B. The Stereochemistry of Conjugate Addition of Lithium Dialkyl Cuprate Reagents to some Carbohydrate α -Enones. *Can. J. Chem.* **1977**, *55*, 4002-4009. (b) Paulsen, H.; Koebernick, W. Branched-chain sugars. XVIII. Stereoselective Synthesis of Branched-Chain Sugars by 1,4-Addition of Carbanions to Pyranoside Enones. *Chem. Ber.* **1977**, *110*, 2127-2145. (c) Martin, S. F.; Dodge, J. A.; Burgess, L. E.; Hartmann, M. A Formal Total Synthesis of (+)-Macbecin I. *J. Org. Chem.* **1992**, *57*, 1070-1072. (d) Lipshutz, B. H.; Wood, M. R.; Tirado, R. Michael Additions of Functionalized Organozinc Reagents Mediated by Catalytic Quantities of Copper(I). *J. Am. Chem. Soc.* **1995**, *117*, 6126-6127. (e) Achmatowicz, O.; Szechner, B.; Maurin, J. K. Addition of Organolithium Reagents to some Carbohydrate Enones. *Tetrahedron* **1997**, *53*, 6035-6044. (f) van der Deen, H.; Kellogg, R. M.; Feringa, B. L. Remarkable O(2)-Effect in 1,4-Additions of Diethylzinc to 6-Acyloxy-2H-pyran-3(6H)-ones and 6-Alkoxy-2H-pyran-3(6H)-ones. *Org. Lett.* **2000**, *2*, 1593-1595. (g) Baba, T.; Huang, G.; Isobe, M. Synthesis of the JKLM-Ring Fragment of Ciguatoxin. *Tetrahedron* **2003**, *59*, 6851-6872. (h) Sugawara, K.; Imanishi, Y.; Hashiyama, T. A Total Synthesis of (17S)-Hexahydro-TMC-69. *Heterocycles* **2007**, *71*, 597-607.

(19) For representative examples, see: (a) Fraser-Reid, B.; Holder, N. L.; Yunker, M. B. Ground and Excited State 1,4-Addition Reactions of Carbohydrate Enones. *J. Chem. Soc. Chem. Commun.* **1972**, 1286-1287. (b) Fraser-Reid, B.; Holder, N. L.; Hicks, D. R.; Walker, D. L. Synthetic Applications of the Photochemically Induced Addition of Oxycarbonyl Species to α -Enones. Part I. The Addition of Simple Alcohols. *Can. J. Chem.* **1977**, *55*, 3978-3985. (c) Sibi, M. P.; Yu, A. Diastereoselective Conjugate Radical Additions to β -Pyranones. *Synlett* **2007**, 3193-3197.

(20) Cu-catalyzed addition of vinylmagnesium bromide: Lipshutz, B. H.; Sengupta, S. Organocopper Reagents: Substitution, Conjugate Addition, Carbo/Metallocupration, and other Reactions. *Org. React.* **1992**, *41*, 135-631.

(21) Cu-catalyzed addition of phenylmagnesium bromide: Sugawara, K.; Hashiyama, T. Stereoselective Nucleophilic Substitution Of 6-(tert-Butyldimethylsilyloxy)-3,6-dihydro-2H-pyran-3-yl Acetate: Application to the Synthesis of a NK1 Receptor Antagonist. *Tetrahedron Lett.* **2007**, *48*, 3723-3726.

(22) For the reaction of boronic acids and potassium organotrifluoroborates with oxonium cations derived from acetals under transition-metal-free conditions, see ref 13 and in addition: (a) Baxter, M.; Bolshan, Y. General Access to Propargylic Ethers through Bronsted Acid Catalyzed Alkynylation of Acetals and Ketals with Trifluoroborates. *Chem. Eur. J.* **2015**, *21*, 13535-13538. (b) Luan, Y.; Barbato, K. S.; Moquist, P. N.; Kodama, T.; Schaus, S. E. Enantioselective Synthesis of 1,2-Dihydronaphthalene-1-carbaldehydes by Addition of Boronates to Isochromene Acetals Catalyzed by Tartaric Acid. *J. Am. Chem. Soc.* **2015**, *137*, 3233-3236. (c) Qian, H.; Zhao, W.; Wang, Z.; Sun, J. Organocatalytic Enantio- and Diastereoselective Synthesis of 1,2-Dihydronaphthalenes from Isobenzopyrylium Ions. *J. Am. Chem. Soc.* **2015**, *137*, 560-563. (d) Fisher, K. M.; Bolshan, Y. Bronsted Acid-Catalyzed Direct Substitution of 2-Ethoxytetrahydrofuran with Trifluoroborate Salts. *Catalysts* **2016**, *6*, 94-98.

(23) For conjugate additions of boronic acids to γ -oxygen-substituted enones, see: Roscales, S.; Csáký, A. G. Trifluoroacetic Anhydride Promoted Tandem Conjugate Addition of Boronic Acids/Acetal Ring Opening. *Org. Lett.* **2012**, *14*, 1187-1189.

(24) The stereoselectivity was determined by inspection of the ^1H -NMR (CHCl_3 , 300 MHz) spectra of the crude reaction products. The relative stereochemistry was determined by NOE measurements. See Supporting Information.

(25) The stereoselectivity was evaluated on the crude reaction product leading to **12a**, prior to acetylation to **16a**.

(26) Unless otherwise stated (see text and Scheme 4), only one diastereomer (2,6-*trans*) was observed on the ^1H -NMR (CHCl_3 , 300 MHz) spectra of the crude reaction products.

(27) Organotrifluoroborates are nucleophilic enough to transfer their organic backbone to electrophilic centers directly without the need of coordination. On the other hand, boronic acids require previous coordination to be transformed into reactive boronate species. See ref. 12-14 and in addition: Lee, S.; MacMillan, D. W. C. Organocatalytic Vinyl and Friedel-Crafts Alkylations with Trifluoroborate Salts. *J. Am. Chem. Soc.* **2007**, *129*, 15438-15439.

(28) See for example: (a) Batey, R. A.; Thadani, A. N.; Smil, D. V.; Lough, A. J. Diastereoselective Allylation and Crotylation Reactions of Aldehydes with Potassium Allyl- and Crotyltrifluoroborates under Lewis Acid Catalysis. *Synthesis* **2000**, 990-998. (b) Mitchell, T. A.; Bode, J. W. Synthesis of Dialkyl Ethers from Organotrifluoroborates and Acetals. *J. Am. Chem. Soc.* **2009**, *131*, 18057-18059. (c) Vo, C.-V. T.; Mitchell, T. A.; Bode, J. W. Expanded Substrate Scope and Improved Reactivity of Ether-Forming Cross-Coupling Reactions of Organotrifluoroborates and Acetals. *J. Am. Chem. Soc.* **2011**, *133*, 14082-14089. (d) Baxter, M.; Bolshan, Y. A General Access to Propargylic Ethers through Bronsted Acid Catalyzed Alkynylation of Acetals and Ketals with Trifluoroborates. *Chem. Eur. J.* **2015**, *21*, 13535. (e) Shih, J. -L.; Nguyen, T. S.; May, J. A. *Angew. Chem. Int. Ed.* **2015**, *54*, 9931-9935.

(29) See in addition: (a) Roscales, S.; Ortega, V.; Csáký, A. G. Transition-Metal-Free Reactions of Boronic Acids: 1,3-Stereochemical Induction in the Substrate-Controlled Conjugate Addition. *J. Org. Chem.* **2013**, *78*, 12825-12830. (b) Roscales, S.; Csáký, A. G. Metal-Free Ring-Opening of Epoxides With Potassium Trifluoroborates. *Chem. Commun.* **2014**, *50*, 454-456. (c) Ortega, V.; Csáký, A. G. Ring-Opening of Donor-Acceptor Cyclopropanes by Boronic Acids and Potassium Organotrifluoroborates under Transition-Metal-Free Conditions. *J. Org. Chem.* **2016**, *81*, 3917-3923.

(30) Ghosh, A. K.; Chen, Z. -H.; Effenberger, K. A.; Jurica, M. S. Enantioselective Total Syntheses of FR901464 and Spliceostatin A and Evaluation of Splicing Activity of Key Derivatives. *J. Org. Chem.* **2014**, *79*, 5697-5709.

(31) Noutsias, D.; Kouridaki, A.; Vassilikogiannakis, G. Scope and Limitations of the Photooxidations of 2-(α -Hydroxyalkyl)furans: Synthesis of 2-Hydroxy-exo-brevicomine. *Org. Lett.* **2011**, *13*, 1166-1169.

(32) Kaminska, J.; Gornicka, I.; Sikora, M.; Gora, J. Preparation of Homochiral (*S*)- and (*R*)-1-(2-Furyl)ethanols by Lipase-Catalyzed Transesterification. *Tetrahedron Asym.* **1996**, *7*, 907-910.

(33) Pawlicki, M.; Latos-Grażyński, L.; Szterenber, L. 5,10,15-Triaryl-21,23-dioxacorrolole and its Isomer with a Protruding Furan Ring. *J. Org. Chem.* **2002**, *67*, 5644-5653.

(34) Martin-Matute, B.; Nevado, C.; Cárdenas, D. J. Intramolecular Reactions of Alkynes with Furans and Electron Rich Arenes Catalyzed by PtCl₂: The Role of Platinum Carbenes as Intermediates. *J. Am. Chem. Soc.* **2003**, *125*, 5757-5766.

(35) Asta, C.; Schmidt, D.; Conrad, J.; Förster-Fromme, B.; Tolasch, T. Beifuss, U. The First Enzymatic Achmatowicz Reaction: Selective Laccase-Catalyzed Synthesis of 6-Hydroxy-(2*H*)-pyran-3(6*H*)-ones and (2*H*)-pyran-2,5(6*H*)-diones. *RSC Adv.* **2013**, *3*, 19259-19263.

(36) Zhou, S.; Wu, K.-H.; Chen, C. -A. Gau, H. -M. Highly Enantioselective Arylation of Aldehydes and Ketones Using AlArEt₂(THF) as Aryl Sources. *J. Org. Chem.* **2009**, *74*, 3500-3505.

(37) Parsons, P. J.; Jones, D. R.; Padgham, A. C.; Allen, L. A. T.; Penkett, C. S.; Green, R. A.; White, A. J. P. A New Approach for the Synthesis of Highly Substituted Aromatic Rings: The Alkyne-Mediated Approach. *Chem. Eur. J.* **2016**, *22*, 3981-3984.

(38) Wei, S.; Du, H. A Highly Enantioselective Hydrogenation of Silyl Enol Ethers Catalyzed by Chiral Frustrated Lewis Pairs. *J. Am. Chem. Soc.* **2014**, *136*, 12261-12264.

(39) Ambre, R.; Yu, C. -Y.; Mane, S. B.; Yao, C. -F.; Hung, C. -H. Toward Carboxylate Group Functionalized A₄, A₂B₂, A₃B Oxaporphyrins and zinc complex of oxaporphyrins. *Tetrahedron* **2011**, *67*, 4680-4688.

(40) Yu, X.; O'Doherty, G. A. De Novo Asymmetric Synthesis and Biological Evaluation of the Trisaccharide Portion of PI-080 and Vineomycin B2. *Org. Lett.* **2008**, *10*, 4529-4532.

(41) Li, Z.; Tong, R. Catalytic Environmentally Friendly Protocol for Achmatowicz Rearrangement. *J. Org. Chem.* **2016**, *81*, 4847-4855.

(42) Kuo, Y. -H.; Shih, K. -S. Preparation of 6-Hydroxy-2*H*-pyran-3(6*H*)-one from 2-Furylcarbinol by Photooxidation. Synthesis of a Pheromone of *Vespa orientalis*. *Heterocycles* **1990**, *31*, 1941-1949.

(43) Bartlett, M. J.; Northcote, P. T.; Lein, M. Harvey, J. E. ¹³C-NMR Analysis of 3,6-Dihydro-2*H*-pyrans: Assignment of Remote Stereochemistry Using Axial Shielding Effects. *J. Org. Chem.* **2014**, *79*, 5521-5532.