


This article was downloaded by: [Duke University Libraries]

On: 17 July 2012, At: 08:59

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcar20>

NOVEL SYNTHESIS OF 1-THIOGLYCOPYRANOSSES VIA THIOIMINIUM SALTS

Takayoshi Fujihira ^a, Mitsutaka Chida ^b, Haruo Kamijo ^b, Toshio Takido ^b & Manabu Seno ^b

^a Japan Sugar Refiners' Association, Technical Department and Research Laboratory, 5-7 Sanbancho, Chiyoda-ku, Tokyo, 102-0075, Japan

^b Department of Materials Applied Chemistry, College of Science and Technology, Nihon University, 1-8-14 Kanda Surugadai, Chiyoda-ku, Tokyo, 101-8308, Japan

Version of record first published: 16 Aug 2006

To cite this article: Takayoshi Fujihira, Mitsutaka Chida, Haruo Kamijo, Toshio Takido & Manabu Seno (2002): NOVEL SYNTHESIS OF 1-THIOGLYCOPYRANOSSES VIA THIOIMINIUM SALTS, Journal of Carbohydrate Chemistry, 21:4, 287-292

To link to this article: <http://dx.doi.org/10.1081/CAR-120013495>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

JOURNAL OF CARBOHYDRATE CHEMISTRY
Vol. 21, No. 4, pp. 287–292, 2002NOVEL SYNTHESIS OF 1-THIOGLYCOPYRANOSSES
VIA THIOIMINIUM SALTSTakayoshi Fujihira,^{1,*} Mitsutaka Chida,² Haruo Kamijo,²
Toshio Takido,² and Manabu Seno²¹Japan Sugar Refiners' Association, Technical Department and Research
Laboratory, 5-7 Sanbancho, Chiyoda-ku, Tokyo 102-0075, Japan²Department of Materials Applied Chemistry, College of Science and
Technology, Nihon University, 1-8-14 Kanda Surugadai, Chiyoda-ku,
Tokyo 101-8308, Japan

ABSTRACT

Methanolysis of a glycosylthioiminium salt, which was prepared from the reaction of acetohalogenosugar with thioacetamide, afforded the corresponding per-*O*-acetylated 1,2-*trans*-1-thioglycose in good yield after fractional crystallization. This synthetic method is very convenient in operation and proceeds without loss of acetyl groups, as the reaction is carried out under mild and neutral conditions.

Key Words: Synthesis; 1-Thioglycopyranose; Thioamide; Iminium salt

INTRODUCTION

1-Thioglycopyranoses are used as starting materials for the synthesis of various 1-thioglycosides; for example, glucosinolates,^[1] carbohydrate clusters^[2] and alkyl thioglycosides.^[3] The 1-thioglycosides are useful as stable glycosyl donors for the synthesis of oligosaccharides.^[4] Tetra-*O*-acetyl-1-thio-*D*-glucopyranose is usually prepared by alkali hydrolysis of the *S*-glucopyranosyl isothiuronium bromide^[5] prepared from the reaction

*Corresponding author. Fax: +81-3-3288-3399; E-mail: fujihira@dream.com

of tetra-*O*-acetyl- α -D-glucopyranosyl bromide with thiourea. Other 1-thioglycoses are also prepared by similar methods.^[1,7–10] It should be noted that, in the reaction with thiourea, the prolonged time required for decomposition of the thioisouronium salt brings about gradual hydrolysis of acetyl groups.^[5] Very recently, the conversion of *S*-glycosyl isothiurea derivatives into thioglycosides without loss of acetyl groups was reported.^[11]


We recently described the preparation of 2-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylthio)ethaniminium bromide from the reaction of tetra-*O*-acetyl- α -D-glucopyranosyl bromide with thioacetamide.^[6] It was found in a subsequent study that the glycosylthioiminium salt affords tetra-*O*-acetyl-1-thio- β -D-glucopyranose by methanolysis. In this paper we report on this novel, simple and effective method for the synthesis of per-*O*-acetylated 1,2-*trans*-1-thioglycoses, including disaccharides.

RESULTS AND DISCUSSION

Per-*O*-acetylated glycopyranosyl bromides **1a** and **1b** were synthesized by the literature procedure^[7] and other glycopyranosyl bromides **1d–1f** were similarly prepared.^[8] Tetra-*O*-acetyl- β -D-mannopyranosyl chloride **1c** is commercially available. According to the method previously described,^[6] 2-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosylthio)ethaniminium bromide **2a** is prepared from tetra-*O*-acetyl- α -D-glucopyranosyl bromide **1a** and thioacetamide in a melt state without solvent or under reflux in dry benzene. As this thioiminium salt was gradually hydrolyzed in the presence of moisture in air, the glycosylthioiminium salt, which is an adduct of acetohalogenosugar and thioacetamide, was predicted to be easily decomposed by the addition of a nucleophilic reagent. It was found that **2a** was decomposed by the addition of methanol for a short time (ca. 10 min.) at 20°C to give tetra-*O*-acetyl-1-thio- β -D-glucopyranose. This synthetic method is applicable to other sugars **1b–1f** (see Scheme 1). In the case of monosaccharides **1a–1c**, freshly prepared and dried (anhydrous conditions) acetohalogenosugar and thioacetamide were stirred just above the melting temperature (ca. 120°C) in a dry nitrogen or argon stream. Although the starting materials (acetohalogenosugar and thioacetamide) were soluble in a warm nonpolar organic solvent (e.g., benzene), the reaction products were barely soluble. The products are mainly the thioiminium salts **2a–2c** as indicated by FAB-MS data; the fragment ion peaks of m/z 406 ($[M-Br^-]^+$) and 331 (pyranosyl ring cation) were clearly observed.^[6] In the case of disaccharides **1d–1f**, the reaction was complete (the reaction mixture became solid and viscous) in approximately 3 h under reflux conditions in a small amount of dry benzene to give glycosylthioiminium salts. The reaction intermediates from disaccharides are presumed to be thioiminium salts as with monosaccharides, although their formation could not be detected by FAB-MS under the present reaction conditions. Side reactions such as the amino-carbonyl reaction were minimized under these conditions. When the reaction of disaccharides was carried out without solvent at an elevated temperature, the amino-carbonyl reaction proceeded to give products of higher melting points. From this result, we decided to include a small amount of dry benzene as a solvent in these cases (see Experimental). As shown in Table 1, per-*O*-acetylated 1,2-*trans*-1-thioglycoses synthesized from monosaccharides are tetra-*O*-acetyl-1-thio- β -D-glucopyranose **3a** (yield 89%), tetra-*O*-acetyl-1-thio- β -D-galactopyranose **3b** (97%) and tetra-*O*-acetyl-1-thio- α -D-mannopyranose **3c** (98%), and from disaccharides, hepta-*O*-acetyl-1-thio- β -lactose **3d** (75%), hepta-*O*-acetyl-1-thio- β -mal-

SYNTHESIS OF 1-THIOGLYCOPYRANOSES

289



Scheme 1. Synthesis of per-*O*-acetylated 1,2-*trans*-1-thioglycopyranoses.

tose **3e** (73%) and hepta-*O*-acetyl-1-thio- β -cellobiose **3f** (76%). In the methanolysis of **2e** the yield of **3e** deposited from the methanol solution was 44%.

¹H NMR coupling constants from 9.6 Hz to 9.9 Hz for H-1 of reducing terminal, except for **3c**, were observed for the methanolysis products. Measurements of ¹H and ¹³C NMR of **3b** were carried out on viscous precipitates after methanol was distilled off, as the crystallization was difficult (Tables 1 and 2). Also, the molecular weights of these products (**3a–3f**) were measured by chemical ionization mass spectrometry, the fragment ion peaks of *m/z* 365 (*M*+1) and 331 (pyranose ring cation) were observed in the case of monosaccharides (**3a–3c**) and the fragment ion peaks of *m/z* 653 (*M*+1) and 619 and 332 (pyranose ring cation) were observed in the case of disaccharides (**3d–3f**). From these results, it was confirmed that per-*O*-acetylated 1,2-*trans*-1-thioglycoses were obtained by the present method.

This reaction is considered to proceed in a similar way as the reaction using thiourea. Glycosylthioiminium salt **2** is formed by the reaction of the acetohalogenosugar with thioacetamide by an S_N2 type reaction and undergoes nucleophilic attack by methanol to give per-*O*-acetylated 1,2-*trans*-1-thioglycose. In the reaction using *d*-methanol, the formation of the product **3** having a deuteriated thiol group was observed. Furthermore, evolution of hydrogen bromide was detected in this reaction and suggests the formation of acetonitrile.^[6]

The reaction temperature must to be monitored carefully in the preparation of glycosylthioiminium salts by this method in order to avoid the amino-carbonyl side reaction. It is necessary for the reaction temperature to be lower than the melting point of thioacetamide (ca. 114°C) and higher than the melting points of acetohalogenosugar. The progress of the side-reaction may be detected by observing a change in color of the reaction mixture; from initial yellow to orange color, and finally brown.



Table 1. Methanolysis^a Glycosylthioiminium Salts **2** and Physical Properties of 1-Thioglycose **3**

Sugar 1	Acetylated R	X	Reaction Temp. of 2 (°C)	Yield (%) of 3	mp (°C) [Ref.]	¹ H NMR ^b (δ, ppm) H-1 of Red. Terminal	Form of 3	CL-MS ^c (M + 1)
a	Glucopyranosyl	Br	120	89	114 ~ 115 [115] ^[5]	4.55 (<i>J</i> _{1,2} = 9.61 Hz)	β	365
b	Galactopyranosyl	Br	120	97	syrup [syrup] ^[9] ^d	4.55 (<i>J</i> _{1,2} = 9.90 Hz)	β	365
c	Mannopyranosyl	Cl	120	98	159 ~ 160 [syrup] ^[10]	4.89 (<i>J</i> _{1,2} = 1.29 Hz)	α	365
d	Lactosyl	Br	reflux ^e	75	81 ~ 82 [syrup] ^[12]	4.54 (<i>J</i> _{1,2} = 9.90 Hz)	β	653
e	Maltosyl	Br	reflux ^e	73 (44) ^f	153 ~ 154 [149 ~ 152] ^[12]	4.59 (<i>J</i> _{1,2} = 9.88 Hz)	β	653
f	Cellobiosyl	Br	reflux ^e	76	208.5 ~ 209.5 [209 ~ 212] ^[13]	4.51 (<i>J</i> _{1,2} = 9.90 Hz)	β	653

^aMethanolysis temp.: 20°C.
^bTMS standard in CDCl₃.
^cGas: isobutane.
^dmp 86.5 ~ 88.0°C (by purifying for analytical purposes).
^eSolvent: dry benzene.
^fYield deposited in methanolysis.



SYNTHESIS OF 1-THIOGLYCOPYRANOSSES

291

Table 2. ^{13}C NMR Chemical Shift Data of 1-Thioglycose **3**

3	Chemical Shift (δ , ppm, CDCl_3)											
	C-1	C-2	C-3	C-4	C-5	C-6	C-1'	C-2'	C-3'	C-4'	C-5'	C-6'
a	78.7	73.5	73.6	68.1	76.3	62.0						
b	79.1	70.8	71.5	67.2	74.9	61.4						
c	76.3	71.6	72.0	65.2	76.9	62.6						
d	78.5	73.9	73.5	70.7	77.2	58.5	101.1	69.1	71.0	66.6	62.2	60.8
e	78.2	74.3	76.1	72.6	76.5	63.0	95.6	69.9	69.3	67.9	68.6	61.4
f	78.5	73.7	73.2	76.7	77.2	61.5	100.8	71.5	67.7	72.0	72.9	62.1

The reaction of acetohalogenosugars with thiourea affords stable thioisouronium salts, from which per-*O*-acetylated 1,2-*trans*-1-thioglycose is derived by alkali hydrolysis. However, the time required for alkali hydrolysis can result in loss of acetyl groups and careful treatment of pH adjustment is always necessary to obtain a good yield. In the present synthetic method using thioacetamide instead of thiourea, the resulting thioiminium salts are easily decomposed even with alcohol, and per-*O*-acetylated 1,2-*trans*-1-thioglycose is obtained without loss of acetyl groups and also tedious pH adjustment. Furthermore, per-*O*-acetylated 1,2-*trans*-1-thioglycose is purified by fractional crystallization from a chilled alcohol solution without the need for separation by column chromatography. In conclusion, the procedure described here is a convenient method to prepare per-*O*-acetylated 1,2-*trans*-1-thioglycose in good yield under mild and neutral conditions.

EXPERIMENTAL

^1H and ^{13}C NMR spectra were recorded with a JEOL GX-400 instrument using tetramethylsilane as a standard. Chemical ionization mass spectra were recorded by direct inlet probe with a JEOL JMS-AMII 150 instrument using isobutane as a reaction gas.

General method (3a–3c). Freshly prepared and dried (anhydrous conditions) acetohalogenosugar (2.5 mmol) and thioacetamide (2.7 mmol, 0.2 g) were stirred mechanically under argon at approximately 120°C for 5 min. Dry methanol (20 mL) was added after cooling to 20°C, and the mixture was stirred for about 10 min (until the solid dissolved). Solvent was evaporated under reduced pressure, a slight amount of methanol was added again, and the mixture cooled to –18°C. The product separated by fractional crystallization was filtered and dried.

For the disaccharides (**3d–3f**), the first step of the reaction was carried out as follows. Freshly prepared and dried acetohalogenosugar (2.5 mmol) and thioacetamide (2.7 mmol, 0.2 g) in dry benzene (below 5 mL) were stirred mechanically at reflux temperature (ca. 82°C) for 3 h under argon. Molecular sieves (4 Å) were included in this reaction mixture in order to maintain anhydrous conditions. Dry methanol (20 mL) was added after cooling to 20°C, and the mixture was stirred for about 10 min (until the solid dissolved). Solvent was evaporated under reduced pressure, and a slight



amount of methanol was added again and the mixture cooled to -18°C . The product separated by fractional crystallization was filtered and dried.

ACKNOWLEDGMENT

We thank Ms. Fujita for measurements of NMR spectra.

REFERENCES

1. Gardrat, C.; Quinsac, A.; Joseph, B.; Rollin, P. Synthesis of indole glycosinolates, sugar variants of naturally occurring glucobrassicin. *Heterocycles* **1993**, 35, 1015–1027.
2. Herczegh, P.; Kovacs-Kulyassa, A.; Erdosi, G.; Kovacs, L.; Menyhart, T.; Szabo Pal, T.; Kornyei, J. Thioglycosidic carbohydrate podands. *Carbohydr. Lett.* **1998**, 3, 59–64.
3. Saito, S.; Tsuchiya, T. Synthesis of alkyl- β -D-thioglycopyranosides, a series of new nonionic detergents. *Chem. Pharm. Bull.* **1985**, 33, 503–508.
4. Fugedi, P.; Garegg, J.; Lonn, H.; Norberg, T. Thioglycosides as glycosylating agents in oligosaccharide synthesis. *Glycoconjugate J.* **1987**, 4, 97–108.
5. Horton, D. 1-Thio-D-Glucose. In *Methods in Carbohydrate Chemistry*; Whistler, R.L., Ed.; Academic Press, Inc.: New York, 1963; Vol. 2, 433–437.
6. Fujihira, T.; Takido, T.; Seno, M. Synthesis of alkyl and arylthioglycosides and thiodisaccharides via thioiminium salts in a two-phase system. *J. Mol. Catal. A: Chem.* **1999**, 137, 65–75.
7. Lemieux, R.U. Acylglycosyl Halides. In *Methods in Carbohydrate Chemistry*; Whistler, R.L., Ed.; Academic Press, Inc.: New York, 1963; Vol. 2, 221–222.
8. Ravindranathan Kartha, K.P.; Jennings, H.J. A simplified, one-pot preparation of acetobromosugars from reducing sugars. *J. Carbohydr. Chem.* **1990**, 9, 777–781.
9. Frgala, J.; Cerny, M.; Stanek, J. Preparation of alkyl 1-thio- β -D-galactopyranosides substituted in the alkyl with reactive groups. *Collect. Czech. Chem. Commun.* **1975**, 40, 1411–1425.
10. Matta, K.L.; Girotra, R.N.; Barlow, J.J. Synthesis of *p*-nitrobenzyl and *p*-nitrophenyl 1-thioglycopyranosides. *Carbohydr. Res.* **1975**, 43, 101–109.
11. Ibatullin, F.M.; Selivanov, S.I.; Shavva, A.G. A general procedure for conversion of *S*-glycosyl isothiurea derivatives into thioglycosides, thiooligosaccharides and glycosyl thioesters. *Synthesis* **2001**, 419–422.
12. Stanek, J.; Sindlerova, M.; Cerny, M. Derivatives of D-thioxylopyranose and of some reducing 1-deoxy-1-thiodisaccharides. *Collect. Czech. Chem. Commun.* **1965**, 30, 297–303.
13. Durette, P.L.; Shen, T.Y. Synthesis of 1-thio- β -cellobiosides. *Carbohydr. Res.* **1978**, 67, 484–490.

Received July 9, 2001

Accepted April 4, 2002