


# PdCl<sub>2</sub>/NaI-Catalyzed Homodimeric Coupling-Cyclization Reaction of 2,3-Allenols: An Efficient Synthesis of 4-(1',3'-Dien-2'-yl)-2,5-dihydrofuran Derivatives

Youqian Deng,<sup>†</sup> Yihua Yu,<sup>‡</sup> and Shengming Ma<sup>\*,†,§</sup>

Laboratory of Molecular Recognition and Synthesis, Department of Chemistry, Zhejiang University, Hangzhou 310027, Zhejiang, People's Republic of China, Shanghai Key Laboratory of Functional Magnetic Resonance Imaging, Department of Physics, East China Normal University, Shanghai 200062, People's Republic of China, and State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai 200032, People's Republic of China

masm@mail.sioc.ac.cn

Received October 29, 2007



A PdCl<sub>2</sub>/NaI-catalyzed homodimeric coupling-cyclization reaction of 2,3-allenols was observed to provide an efficient route to 4-(1',3'-dien-2'-yl)-2,5-dihydrofuran derivatives. By using the easily available optically active starting materials, 2,5-dihydrofurans with high enantiopurity may be prepared. A Pd(II)-catalyzed mechanism was also discussed.

## Introduction

The transition metal-catalyzed cyclizative dimerization reaction of two functionalized allenes is attractive for synthetic organic chemists due to the issues of chirality transfer, diversity, and substituent-loading capability of allenes.<sup>1,2</sup> The first palladium-catalyzed homodimerization reaction of 1,2-allenyl ketones has been reported by Hashmi et al., which led to the formation of monocyclic 3-(3'-oxo-1'-alkenyl)-substituted furan derivatives.<sup>3,4</sup> The reaction of 1,2-allenyl ketones would also afford 2-(3'-oxo-1'-alkenyl)-substituted furan derivatives when AuCl<sub>3</sub> was used as the catalyst.<sup>5</sup> The palladium-catalyzed macrocyclization reaction of 1,ω-bis(1,2-allenylketone)s was also reported.<sup>6</sup> We have reported the homodimerization reaction of 2,3-allenoic acids affording bibutenolides, in which both

allenes were cyclized.<sup>7</sup> Recently, Hashmi et al. reported the AuCl<sub>3</sub>-catalyzed reaction of 2,3-allenols, which afforded bis-(2,5-dihydrofuran)s as a byproduct in low yield.<sup>8</sup> On the other

\* Address correspondence to this author. Phone: 86-21-549-25147. Fax: 86-21-641-67510.

<sup>†</sup> Zhejiang University.

<sup>‡</sup> East China Normal University.

<sup>§</sup> Chinese Academy of Sciences.

(1) (a) Patai, S. *The Chemistry of Ketenes, Allenes, and Related Compounds*; John Wiley & Sons: New York, 1980; Part 1. (b) Schuster, H. F.; Coppola, G. M. *Allenols in Organic Synthesis*; John Wiley & Sons: New York, 1984. (c) Landor, S. R. *The Chemistry of Allenes*; Academic Press: New York, 1982; Vols. 1–3. (d) Krause, N.; Hashmi, A. S. K., Eds. *Modern Allene Chemistry*; Wiley-VCH: Weinheim, Germany, 2004; Vols. 1 and 2.

(2) For reviews and accounts, see: (a) Yamamoto, Y.; Radhakrishnan, U. *Chem. Soc. Rev.* **1999**, 28, 199. (b) Larock, R. C. *J. Organomet. Chem.* **1999**, 576, 111. (c) Grigg, R.; Sridharan, V. *J. Organomet. Chem.* **1999**, 576, 65. (d) Zimmer, R.; Dinesh, C. U.; Nandan, E.; Khan, F. A. *Chem. Rev.* **2000**, 100, 3067. (e) Hashmi, A. S. K. *Angew. Chem., Int. Ed.* **2000**, 39, 3590. (f) Reissig, H.-U.; Hormuth, S.; Schade, W.; Okala Amombo, M.; Watanabe, T.; Pulz, R.; Hausherr, A.; Zimmer, R. *J. Heterocycl. Chem.* **2000**, 37, 597. (g) Ma, S.; Li, L. *Synlett* **2001**, 1206. (h) Reissig, H.-U.; Schade, W.; Okala Amombo, M.; Pulz, R.; Hausherr, A. *Pure Appl. Chem.* **2002**, 74, 175. (i) Ma, S. Carbopalladation of Allenes. In *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; Wiley-Interscience: New York, 2002; p 1491. (j) Ma, S. *Acc. Chem. Res.* **2003**, 36, 701. (k) Ma, S. *Chem. Rev.* **2005**, 105, 2829. (l) Ma, S. Pd-Catalyzed two- or three-component cyclization of functionalized allenes. In *Topics in Organometallic Chemistry*; Tsuji, J., Ed.; Springer-Verlag: Heidelberg, Germany, 2005; pp 183–210.

(3) (a) Hashmi, A. S. K. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 1581. (b) Hashmi, A. S. K.; Ruppert, T. L.; Knöfel, T.; Bats, J. W. *J. Org. Chem.* **1997**, 62, 7295.

(4) For a most recent report from Alcaide's group, see: Alcaide, B.; Almendros, P.; Martínez del Campo, T. *Eur. J. Org. Chem.* **2007**, 2844.

(5) Hashmi, A. S. K.; Schwarz, L.; Choi, J.-H.; Frost, T. M. *Angew. Chem., Int. Ed.* **2000**, 39, 2285.

(6) Hashmi, A. S. K.; Schwarz, L.; Bolte, M. *Eur. J. Org. Chem.* **2004**, 1923.

(7) (a) Ma, S.; Yu, Z. *Org. Lett.* **2003**, 5, 1507. (b) Ma, S.; Yu, Z.; Gu, Z. *Chem. Eur. J.* **2005**, 11, 2351.

hand, 2,5-dihydrofurans, an important class of heterocyclic compounds, are useful intermediates for organic synthesis<sup>9</sup> and common structural units in many natural products.<sup>10</sup> These compounds are usually prepared via cyclization of 2,3-allenols,<sup>11–16</sup> a RCM reaction,<sup>17</sup> Ag(I)-catalyzed rearrangement-cyclization of 4-hydroxypropargyl esters,<sup>18</sup> dehydration of *cis*-2-alken-1,4-diols,<sup>19</sup> palladium-catalyzed reaction of cyclic alkynyl carbonates with electron-deficient alkenes,<sup>20</sup> Prins reaction of terminal alkene and formaldehyde,<sup>21</sup> reaction of oxazirconacyclopentenes with propynoates,<sup>22</sup> reaction of 1,4-dithio-1,3-dienes with aldehydes,<sup>23</sup> tungsten-promoted intramolecular annulation of propargyl bromides with ketones and aldehydes,<sup>24</sup> and Au(I)-catalyzed rearrangement of butynediol monobenzoates.<sup>25</sup> In this paper, we wish to report an efficient synthesis of 4-(1',3'-dien-2'-yl)-2,5-dihydrofuran derivatives via the PdCl<sub>2</sub>/

**TABLE 1.** Pd(II)-Catalyzed Homodimeric Coupling-Cyclization Reaction of 2,3-Allenol (**1a**)

entry	solvent	additive (equiv)	time (h)	yield of <b>2a</b> <sup>a</sup> (%)
1	DMA	none	24	trace
2	DMA	NaCl (0.5)	12	22 <sup>b,d</sup>
3	DMA	LiCl (0.5)	12	25 <sup>b</sup>
4	DMA	NaI (0.1)	5	42
5	DMA	NaI (0.3)	5	80
6	DMA	NaI (0.5)	5	86
7	DMA	NaI (0.5)	21	75 <sup>b,c,e</sup>
8	DMA	NaI (1.0)	5	81
9	DMA	KI (0.5)	5	85
10	DMA	I <sub>2</sub> (0.25)	0.5	66
11	DMA	I <sub>2</sub> (0.05)	3	42
12	DMA	CH <sub>3</sub> I (10)	1.5	86
13	THF	NaI (0.5)	17	60 <sup>b,f</sup>
14	acetone	NaI (0.5)	17	47 <sup>b,g</sup>
15	CH <sub>3</sub> CN	NaI (0.5)	17	44 <sup>b,h</sup>

<sup>a</sup> Isolated yield. <sup>b</sup> Determined by <sup>1</sup>H NMR analysis using 1,3,5-trimethylbenzene as the internal standard. <sup>c</sup> 2 mol % PdCl<sub>2</sub> was used. <sup>d</sup> 10% of **1a** was recovered. <sup>e</sup> 16% of **1a** was recovered. <sup>f</sup> 11% of **1a** was recovered. <sup>g</sup> 40% of **1a** was recovered. <sup>h</sup> 33% of **1a** was recovered.

NaI-catalyzed homodimeric coupling-cyclization reaction of 2,3-allenols, in which one molecule of the 2,3-allenol was cyclized while the other formed the 1,3-diene unit and helped to regenerate the catalytically active Pd(II) species.<sup>26</sup>

## Results and Discussion

**PdCl<sub>2</sub>/NaI-Catalyzed Homodimeric Coupling-Cyclization Reaction of 2,3-Allenols.** The homodimeric coupling-cyclization reaction of 2,3-allenol (**1a**) was chosen to establish the protocol. Some representative results are listed in Table 1. From Table 1, it was observed that under the catalysis of PdCl<sub>2</sub>, the reaction of **1a** afforded a trace amount of 3-(1-cyclohexylidenemethylvinyl)-1-oxaspiro[4.5]dec-3-ene (**2a**) (Table 1, entry 1). When NaCl or LiCl was applied as the additive, the reaction afforded **2a** in 22% and 25% yields, respectively (Table 1, entries 2 and 3). Surprisingly, when NaI<sup>7b</sup> was applied as the additive, the reaction afforded **2a** in higher yields (Table 1, entries 4–8). Addition of KI<sup>7b</sup> (Table 1, entry 9), I<sub>2</sub><sup>7b</sup> (Table 1, entries 10 and 11), or CH<sub>3</sub>I<sup>7a</sup> (Table 1, entry 12) is also effective for this reaction. Among the solvents tested, DMA is the best. In THF, acetone, and CH<sub>3</sub>CN, the yields were lower and some of **1a** was recovered (Table 1, entries 13–15). In conclusion, the best results were obtained when the reaction was conducted using 5 mol % PdCl<sub>2</sub> and 0.5 equiv of NaI in DMA leading to a 86% yield of product **2a** (Table 1, entry 6).

Some typical results are summarized in Tables 2 and 3. Various substituted 2,3-allenols that bear alkyl or aryl groups were successfully used to afford 4-(1',3'-dien-2'-yl)-2,5-dihydrofuran derivatives in moderate to good yields. Furthermore, it is important to note that a high stereoselectivity for the

(26) This reaction was first observed in this group during the study of the PdCl<sub>2</sub>-catalyzed cyclization of propadienyl cyclohexanol in the presence of allylic bromide, see ref 15b.

(8) Hashmi, A. S. K.; Carmen Blanco, M.; Fischer, D.; Bats, J. W. *Eur. J. Org. Chem.* **2006**, 1387.

(9) (a) Marshall, J. A.; Pinney, K. G. *J. Org. Chem.* **1993**, *58*, 7180. (b) Marshall, J. A.; Yu, B. *J. Org. Chem.* **1994**, *59*, 324. (c) Yamada, O.; Ogasawara, K. *Synlett* **1995**, 427. (d) Kim, S.; Kim, K. H. *J. Chem. Soc., Perkin Trans. 1* **1997**, 1095. (e) Paolucci, C.; Musiani, L.; Venturelli, F.; Fava, A. *Synthesis* **1997**, 1415. (f) Shieh, S.-J.; Fan, J.-S.; Chandrasekharan, M.; Liao, F.-L.; Wang, S.-L.; Liu, R.-S. *Organometallics* **1997**, *16*, 3987. (g) Doyle, M. P.; Forbes, D. C.; Protopopova, M. N.; Stanley, S. A.; Vasbinder, M. M.; Xavier, K. R. *J. Org. Chem.* **1997**, *62*, 7210. (h) Crotti, P.; Bussolo, V. D.; Favero, L.; Pineschi, M. *Tetrahedron* **1997**, *53*, 1417. (i) Bauer, T. *Tetrahedron* **1997**, *53*, 4763.

(10) (a) Kupchan, S. M.; Davies, V. H.; Fujita, T.; Cox, M. R.; Restivo, R. J.; Bryan, R. F. *J. Org. Chem.* **1973**, *38*, 1853. (b) Semple, J. E.; Wang, P. C.; Lysenko, Z.; Joullie, M. M. *J. Am. Chem. Soc.* **1980**, *102*, 7505.

(11) For a related recent review, see ref 1d.

(12) For the electrophilic cyclization of 2,3-allenols, see: (a) Beaulieu, P. L.; Morisset, V. M.; Garratt, D. G. *Tetrahedron Lett.* **1980**, *21*, 129. (b) Smadja, W. *Chem. Rev.* **1983**, *83*, 263. (c) Bridges, A. J.; Thomas, R. D. *J. Chem. Soc., Chem. Commun.* **1984**, 694. (d) Whitby, R.; Kocienski, P. *Tetrahedron Lett.* **1987**, *28*, 3619. (e) Marshall, J. A.; Wang, X. *J. Org. Chem.* **1990**, *55*, 2995. (f) Marshall, J. A.; Wang, X. *J. Org. Chem.* **1991**, *56*, 4913.

(13) For the base-mediated cyclization of 2,3-allenols, see: (a) Hoff, S.; Brandsma, L.; Arens, J. F. *Recl. Trav. Chim. Pays-Bas* **1969**, *88*, 609. (b) Brasholz, M.; Rössig, H.-U. *Angew. Chem., Int. Ed.* **2007**, *46*, 1634. (c) Katritzky, A. R.; Verin, S. V. *J. Heterocycl. Chem.* **1995**, *32*, 323. (d) Magnus, P.; Albaugh-Robertson, P. J. *Chem. Soc., Chem. Commun.* **1984**, 805.

(14) For the metal-catalyzed cycloisomerization reaction, see: Ag<sup>+</sup>: (a) Olsson, L. I.; Claesson, A. *Synthesis* **1979**, 743. (b) Nikam, S. S.; Chu, K. H.; Wang, K. K. *J. Org. Chem.* **1986**, *51*, 745. (c) Marshall, J. A.; Sehon, C. A. *J. Org. Chem.* **1995**, *60*, 5966. (d) Marshall, J. A.; Yu, R. H.; Perkins, J. F. *J. Org. Chem.* **1995**, *60*, 5550. Hg<sup>2+</sup>: (a) Gelin, R.; Gelin, S.; Albrand, M. *Bull. Soc. Chim. Fr.* **1972**, 1946. (b) Walkup, R. D.; Park, G. *Tetrahedron Lett.* **1987**, *28*, 1023. Au<sup>3+</sup>: Hoffmann-Röder, A.; Krause, N. *Org. Lett.* **2001**, *3*, 2537.

(15) For the Pd-catalyzed cyclization reaction of 2,3-allenols in the presence of allylic halides, see: (a) Ma, S.; Gao, W. *Tetrahedron Lett.* **2000**, *41*, 8933. (b) Ma, S.; Gao, W. *J. Org. Chem.* **2002**, *67*, 6104.

(16) For the Pd-Cu catalyzed, domino cyclization of 2,3-allenols with acrylate, alkynes, and boronic acids, see: Alcaide, B.; Almendros, P.; Rodríguez-Acebes, R. *Chem. Eur. J.* **2005**, *11*, 5708.

(17) Carda, M.; Castillo, E.; Rodríguez, S.; Uriel, S.; Marco, J. A. *Synlett* **1999**, 1639.

(18) (a) Shigemasa, Y.; Yasui, M.; Ohrai, S.; Sasaki, M.; Sashiwa, H.; Saimoto, H. *J. Org. Chem.* **1991**, *56*, 910. (b) Saimoto, H.; Yasui, M.; Ohrai, S.; Oikawa, H.; Yokoyama, K.; Shigemasa, Y. *Bull. Chem. Soc. Jpn.* **1999**, *72*, 279.

(19) Sato, F.; Kanbara, H.; Tanaka, Y. *Tetrahedron Lett.* **1984**, *25*, 5063.

(20) Darcel, C.; Bruneau, C.; Albert, M.; Dixneuf, P. H. *Chem. Commun.* **1996**, 919.

(21) Talipov, R. F.; Starikov, A. S.; Gorina, I. A.; Akmanova, N. A.; Safarov, M. G. *Zh. Org. Khim.* **1993**, *29*, 1024.

(22) Xi, C.; Kotora, M.; Takahashi, T. *Tetrahedron Lett.* **1999**, *40*, 2375.

(23) Chen, J.; Song, Q.; Li, P.; Guan, H.; Jin, X.; Xi, Z. *Org. Lett.* **2002**, *4*, 2269.

(24) Shieh, S.-J.; Tang, T.-C.; Lee, J.-S.; Lee, G.-H.; Peng, S.-M.; Liu, R.-S. *J. Org. Chem.* **1996**, *61*, 3245.

(25) Buzas, A.; Istrate, F.; Gagosz, F. *Org. Lett.* **2006**, *8*, 1957.


SCHEME 1. Homodimeric Coupling-Cyclization Reaction of Optically Active 2,3-Allenols (**1g**)

TABLE 2. Homodimeric Coupling-Cyclization Reaction of 2,3-Allenols

Reaction scheme for Table 2: Homodimeric Coupling-Cyclization Reaction of 2,3-Allenols. The reaction conditions are 5 mol % PdCl<sub>2</sub>, 0.5 equiv NaI, DMA, 80 °C. The starting material **1** is a 2,3-allenol with substituents R<sup>1</sup> and R<sup>2</sup>. The product **2** is a 2,5-dihydrofuran derivative with substituents R<sup>1</sup> and R<sup>2</sup>.

entry	R <sup>1</sup>	R <sup>2</sup>	R <sup>2</sup>	time (h)	isolated yield of <b>2</b> (%)
1	H	(CH <sub>2</sub> ) <sub>5</sub> ( <b>1a</b> )	CH <sub>3</sub> ( <b>1b</b> )	5	86 ( <b>2a</b> )
2	H	CH <sub>3</sub>	CH <sub>3</sub> ( <b>1b</b> )	5	66 ( <b>2b</b> )
3	H	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub> ( <b>1c</b> )	4	85 ( <b>2c</b> )
4	H	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	<i>n</i> -C <sub>4</sub> H <sub>9</sub> ( <b>1d</b> )	10	92 ( <b>2d</b> )
5	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	H	H ( <b>1e</b> )	12	84 ( <b>2e</b> )
6	<i>n</i> -C <sub>7</sub> H <sub>15</sub>	H	H ( <b>1f</b> )	12	87 ( <b>2f</b> )

TABLE 3. Stereoselective Homodimeric Coupling-Cyclization Reaction of 2,3-Allenols


Reaction scheme for Table 3: Stereoselective Homodimeric Coupling-Cyclization Reaction of 2,3-Allenols. The reaction conditions are 5 mol % PdCl<sub>2</sub>, 0.5 equiv NaI, DMA, 80 °C. The starting material **1** is a 2,3-allenol with substituents R<sup>1</sup> and R<sup>2</sup>. The product **2** is a 2,5-dihydrofuran derivative with substituents R<sup>1</sup> and R<sup>2</sup>.

entry	R <sup>1</sup>	R <sup>2</sup>	time (h)	yield of <b>2</b> <sup>a</sup> (%)	<i>E</i> / <i>Z</i> ratio of <b>2</b> <sup>b</sup>
1	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	CH <sub>3</sub> ( <b>1g</b> )	13	83 ( <b>2g</b> )	≥95:5
2	<i>n</i> -C <sub>4</sub> H <sub>9</sub>	C <sub>2</sub> H <sub>5</sub> ( <b>1h</b> )	14	86 ( <b>2h</b> )	≥96:4
3	Ph	<i>n</i> -C <sub>4</sub> H <sub>9</sub> ( <b>1i</b> )	11	75 ( <b>2i</b> )	>99:1 <sup>c</sup>
4	CO <sub>2</sub> CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub> ( <b>1j</b> )	17	58 ( <b>2j</b> )	95:5 <sup>c</sup>
5	allyl	CH <sub>3</sub> ( <b>1k</b> )	1.5	61 ( <b>2k</b> )	>99:1

<sup>a</sup> Isolated yield. <sup>b</sup> *E*/*Z* was determined by <sup>1</sup>H NMR analysis. <sup>c</sup> *Z*/*E* ratio.

formation of the C=C bond carrying R<sup>1</sup> and R<sup>2</sup> groups was observed affording the products (*E*)-**2** when secondary 2,3-allenols were applied (Table 3). The stereochemistry of these products was determined by the NOESY study of (*E*)-**2h**.

In addition, by using the optically active 2,3-allenols *R*-(+)-**1g** (97% ee) and *S*-(-)-**1g** (98% ee),<sup>27</sup> the reaction occurred smoothly to give the product *R*-**2g** in 86% yield and 97% ee and *S*-**2g** in 88% yield and 99% ee, respectively (Scheme 1). These results indicated that no racemization was observed under the standard reaction conditions.




SCHEME 2. PdX<sub>2</sub>-Catalyzed Homodimeric Self-Coupling Cyclization Reaction of 2,3-Allenol (**1a**)

X = Cl,	24 h,	trace
X = Br,	16 h,	65%
X = I,	5 h,	90%

To clarify the mechanism of this reaction, three control experiments were conducted (Scheme 2). It should be noted that when 5 mol % PdBr<sub>2</sub> or PdI<sub>2</sub> was used, the reaction proceeded even in the absence of NaI affording **2a** in 65% or 90% yields, respectively, indicating the possible in situ formation of PdI<sub>2</sub> under the current reaction conditions.

On the basis of these experimental findings, it was proposed that the interaction of **1g** with PdI<sub>2</sub>, which might be produced in situ from PdCl<sub>2</sub> and NaI, would form 2,5-dihydrofuranyl palladium intermediate **M1** via cyclic oxypalladation. Then regioselective carbopalladation of a second molecule of **1g** with **M1** would highly regioselectively form the  $\pi$ -allylic palladium intermediate **M2**. Subsequent *trans*- $\beta$ -hydroxide elimination<sup>28–31</sup> would afford **2g** and IPd<sup>+</sup>[OH<sup>−</sup>]. Finally, IPd<sup>+</sup>[OH<sup>−</sup>] was converted to the catalytically active species PdI<sub>2</sub> by its reaction with HI generated in the first step (Scheme 3). It should be noted when R<sup>1</sup>  $\neq$  H, R<sup>2</sup>  $\neq$  H, and R<sup>3</sup> = H, the intermediate **M3** was more favored for its thermodynamic stability over **M4**. **M3** would afford *E*-isomers highly stereoselectively through a *trans*- $\beta$ -hydroxide elimination (Scheme 4).

(28) Ma, S.; Gu, Z. *J. Am. Chem. Soc.* **2005**, *127*, 6182.(29) (a) Harrington, P. J.; Hegedus, L. S.; McDaniel, K. F. *J. Am. Chem. Soc.* **1987**, *109*, 4335. (b) Francis, J. W.; Henry, P. M. *Organometallics* **1991**, *10*, 3498. (c) Saito, S.; Hara, T.; Takahashi, N.; Hirai, M.; Moriwake, T. *Synlett* **1992**, 237. (d) Ma, S.; Lu, X. *J. Organomet. Chem.* **1993**, *447*, 305.(30) (a) Kimura, M.; Horino, Y.; Mukai, R.; Tanaka, S.; Tamaru, Y. *J. Am. Chem. Soc.* **2001**, *123*, 10401. (b) Ozawa, F.; Okamoto, H.; Kawagishi, S.; Yamamoto, S.; Minami, T.; Yoshifuji, M. *J. Am. Chem. Soc.* **2002**, *124*, 10968. (c) Manabe, K.; Kobayashi, S. *Org. Lett.* **2003**, *5*, 3241. (d) Kabalka, G. W.; Dong, G.; Venkataiah, B. *Org. Lett.* **2003**, *5*, 893. (e) Yoshida, M.; Gotou, T.; Ihara, M. *Chem. Commun.* **2004**, 1124.(31) For the stereoselectivity of  $\beta$ -heteroatom elimination, see: (a) Frost, C. G.; Howarth, J.; Williams, J. M. J. *Tetrahedron: Asymmetry* **1992**, *3*, 1089. (b) Daves, G. D., Jr. *Acc. Chem. Res.* **1990**, *23*, 201. (c) Zhu, G.; Lu, X. *Organometallics* **1995**, *14*, 4899. (d) Zhang, Z.; Lu, X.; Xu, Z.; Zhang, Q.; Han, X. *Organometallics* **2001**, *20*, 3724. (e) Alcaide, B.; Almendros, P.; Martínez del Campo, T. *Angew. Chem., Int. Ed.* **2006**, *45*, 4501.(27) Xu, D.; Li, Z.; Ma, S. *Chem. Eur. J.* **2002**, *8*, 5012.


**SCHEME 3. Possible Mechanism of the Homodimeric Coupling-Cyclization Reaction of 1g**

**SCHEME 4. A Rationale for the Stereoselectivity Observed**

**SCHEME 5. The Diels–Alder Reaction of 2a with 3**


Furthermore, the Diels–Alder reaction of **2a** with maleamide **3** afforded diastereoisomeric polycyclic compounds **4aa** and **4ab** in 81% and 8% yields, respectively (Scheme 5). The structure of the major product **4aa** was further established by the X-ray diffraction study (Figure 1).

In conclusion, we have developed a homodimeric coupling-cyclization reaction of 2,3-allenols using  $\text{PdCl}_2/\text{NaI}$  as the catalyst, which provides an efficient route to 4-(1',3'-dien-2'-yl)-2,5-dihydrofuran derivatives. By using the optically active 2,3-allenols, no racemization was observed under the standard reaction conditions, affording optically active products in good yields. On the basis of control experiment, it was found that the addition of NaI is very important for this transformation. Due to the easy availability of the starting allenols and usefulness of 2,5-dihydrofuran, this method will be useful in organic synthesis. Further studies on the scope, mechanism, and synthetic applications of this reaction are being carried out in our laboratory.

**Experimental Section**

**Synthesis of 3-(1-Cyclohexylidenemethylvinyl)-1-oxaspiro[4.5]-dec-3-ene (2a): Typical Procedure.** To a mixture of  $\text{PdCl}_2$  (8.8 mg, 5 mol %), 0.050 mmol) and NaI (75.1 mg, 0.50 mmol) were



**FIGURE 1.** ORTEP structure of **4aa**.

added **1a** (136.8 mg, 0.99 mmol) and DMA (2 mL). After being stirred at 80 °C for 5 h the reaction was complete as monitored by TLC and the resulting mixture was cooled to room temperature and quenched by 10 mL of water. The mixture was extracted with  $\text{Et}_2\text{O}$  ( $3 \times 25$  mL). The combined organic layer was washed with a saturated aqueous solution of  $\text{Na}_2\text{S}_2\text{O}_3$  and brine. The product solution was dried over anhydrous  $\text{Na}_2\text{SO}_4$ . Evaporation and column chromatography on silica gel (petroleum ether/diethyl ether = 20:1) afforded **2a** (109.3 mg, 86%): liquid;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  5.78 (s, 1H), 5.74 (s, 1H), 4.89 (s, 1H), 4.86 (s, 1H), 4.73 (d,  $J = 2.0$  Hz, 2H), 2.20 (t,  $J = 5.6$  Hz, 2H), 2.16 (t,  $J = 5.6$  Hz, 2H), 1.72–1.28 (m, 16H).

**Synthesis of Polycyclic Compound 4aa.** To maleamide **3** (519.3 mg, 3.0 mmol) were added **2a** (387.8 mg, 1.5 mmol) and toluene (15 mL). The mixture was refluxed at 130 °C for 5 h. After the reaction was complete as monitored by TLC, it was cooled to room temperature, evaporated, and analyzed by 300 MHz  $^1\text{H}$  NMR spectroscopic analysis with  $\text{CH}_2\text{Br}_2$  (105  $\mu\text{L}$ , 1.5 mmol) as the internal standard, which showed that the diastereomeric ratio of **4aa/4ab** is 91/9. Column chromatography on silica gel (petroleum ether/ethyl acetate = 8:1) afforded 54.1 mg (8%) of **4ab** and 524.0 mg (81%) of **4aa**. **4ab**: oil;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.52–7.43 (m, 2H), 7.43–7.34 (m, 1H), 7.30 (d,  $J = 7.5$  Hz, 2H), 5.55 (s, 1H), 4.26 (dd,  $J_1 = 27.9$  Hz,  $J_2 = 13.8$  Hz, 2H), 3.08–2.95 (m, 2H), 2.80–2.69 (m, 1H), 2.50–2.40 (m, 1H), 2.25–1.90 (m, 6H), 1.80–1.40 (m, 15H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  178.00, 177.95, 143.5, 137.3, 131.7, 129.1, 128.5, 126.3, 124.8, 119.7, 83.7, 66.4, 48.9, 40.6, 40.3, 37.0, 36.4, 30.7, 30.1, 29.1, 28.5, 27.7, 26.4, 25.6, 22.8, 21.7; MS ( $m/z$ ) 431 ( $\text{M}^+$ , 3.21), 333 (100); IR (neat,  $\text{cm}^{-1}$ ) 2929, 2853, 1779, 1716, 1599, 1500, 1447, 1380, 1178; HRMS calcd for  $\text{C}_{28}\text{H}_{33}\text{NO}_3$  ( $\text{M}^+$ ) 431.2460, found 431.2450. **4aa**: solid, mp 178–179 °C (ethyl acetate);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.46–7.27 (m, 3H), 7.13 (d,  $J = 8.1$  Hz, 2H), 5.44 (s, 1H), 4.21 (s, 2H), 3.54–3.44 (m, 1H), 3.27 (t,  $J = 8.1$  Hz, 1H), 2.78 (d,  $J = 14.7$  Hz, 1H), 2.55–2.38 (m, 2H), 2.40–2.30 (m, 1H), 2.20–2.00 (m, 4H), 2.08–1.30 (m, 15H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  178.4, 175.5, 143.7, 139.7, 131.9, 128.8, 128.3, 126.4, 125.1, 119.7, 82.9, 66.1, 49.9, 41.7, 41.1, 36.8, 36.1, 32.1, 31.7, 30.5, 28.3, 27.5, 26.3, 25.6, 23.0; MS ( $m/z$ ) 431 ( $\text{M}^+$ , 7.89), 333 (100); IR (neat,  $\text{cm}^{-1}$ ) 2928, 2853, 1775, 1713, 1598, 1500, 1446, 1382. Elemental analysis calcd for  $\text{C}_{28}\text{H}_{33}\text{NO}_3$ : C, 77.93; H, 7.71; N, 3.25. Found: C, 77.74; H, 7.81; N, 2.99. Crystal data for **4aa**:  $\text{C}_{28}\text{H}_{33}\text{NO}_3$ , MW = 431.55, monoclinic, space group  $P2(1)/c$ , final  $R$  indices [ $I > 2\sigma(I)$ ],  $R_1 = 0.0688$ ,  $wR_2 = 0.1807$ ,  $a = 16.051(6)$  Å,  $b = 13.024(5)$  Å,  $c = 11.211(4)$  Å,  $\alpha = 90^\circ$ ,  $\beta = 97.815(8)^\circ$ ,  $\gamma = 90^\circ$ ,  $V = 2322.0(15)$  Å<sup>3</sup>, crystal size  $0.490 \times 0.362 \times 0.178$  mm<sup>3</sup>,  $T = 293(2)$  K,  $Z = 4$ , reflections collected/unique 13674/5282 ( $R_{\text{int}} = 0.0967$ ), 5282



data, 1 restraint, 302 parameters. Supplementary crystallographic data have been deposited at the Cambridge Crystallographic Data Center as CCDC 651148.

**Acknowledgment.** Financial support from the National Natural Science Foundation of China (20332060), Major State Basic Research and Development Program (2006CB806105), and Cheung Kong Scholar Program is greatly appreciated. This work was conducted at Zhejiang University. We thank Mr.

Guangke He in this group for reproducing the results presented in entries 1 and 4 of Table 2 and entries 1 and 4 of Table 3.

**Supporting Information Available:** Experimental procedures, copies of  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of all compounds, and the CIF file of compound **4aa**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO702332M