

# Photoreaction of Benzoic Acid with Sodium Hypochlorite in Aqueous Alkali


Yoshiro Ogata\* and Kohtaro Tomizawa

Department of Applied Chemistry, Faculty of Engineering, Nagoya University, Chikusa-ku, Nagoya, Japan

The photoreaction of benzoic acid with sodium hypochlorite in aqueous alkali ( $\text{pH} \geq 12$ ) has been studied. At a low initial ratio of  $[\text{ClO}^-]/[\text{PhCOO}^-]$ , *e.g.* 0.1, hydroxylation and chlorination at the aromatic ring occur simultaneously with *ipso*-substitution of the carboxylate group to give hydroxy- and chlorobenzoic acids together with phenol. The product distribution depends on the wavelength of light, which implies the dependence on the concentration of active species generated from  $\text{ClO}^-$  and on the light stability of products. At comparable concentrations of  $\text{ClO}^-$  and  $\text{PhCOO}^-$ , the products initially formed from the photoreaction react further with  $\text{ClO}^-$  in the dark to give polychlorinated derivatives. The initial steps for the photoreaction are discussed on the basis of the reactivity of the active species,  $\text{O}(^3\text{P})$ ,  $\text{O}(^1\text{D})$ ,  $\text{O}^\cdot$ , and  $\text{Cl}^\cdot$ , generated by photolysis of  $\text{ClO}^-$ .

Sodium hypochlorite can be used for selective oxidation and halogenation of organic compounds,<sup>1</sup> and for oxidative removal of pollutants.<sup>2</sup> Hypochlorite ion ( $\text{ClO}^-$ ) under irradiation delivers highly active atomic oxygen,  $\text{O}(^3\text{P})$  and  $\text{O}(^1\text{D})$ , along with  $\text{O}^\cdot$  depending on the wavelength used.<sup>3,4</sup> The gas-phase reactions of  $\text{O}(^3\text{P})$  were reported.<sup>5</sup>

As we have reported,<sup>6a</sup> the reaction of benzoic acid with  $\text{HO}^\cdot$  radical from irradiated  $\text{H}_2\text{O}_2$  under acidic conditions is mainly initiated by addition of  $\text{HO}^\cdot$  to the ring and by hydrogen atom abstraction from  $\text{COOH}$  as a minor process [equation (1a)].<sup>6a</sup> On the other hand, in the reaction of  $\text{PhCOO}^-$  with  $\text{HO}^\cdot$  under neutral conditions, *ipso*-attack of  $\text{HO}^\cdot$  on  $\text{PhCOO}^-$  [equation (1b)] together with addition of  $\text{HO}^\cdot$  to the ring are observed.<sup>6b</sup>



The present study reports that the title reaction leads preferentially to hydroxylation rather than chlorination. The reaction mechanism is discussed.

## Results and Discussion

For an examination of the dependence of the reactivity of active species on the wavelength, a mixture of  $\text{PhCOONa}$  (0.01M) and  $\text{NaClO}$  (0.001–0.200M) in aqueous alkali ( $\text{pH} \geq 12$ ) was photolysed at 254 and  $>350$  nm. The re-

maining  $\text{ClO}^-$  after irradiation was quickly reduced with aqueous  $\text{Na}_2\text{S}_2\text{O}_3$ . The reaction products observed by g.l.c. and g.l.c.-mass spectrometry were composed of hydroxybenzoic acids ( $p \gtrsim o > m$ ), chlorobenzoic acids ( $o \gtrsim p > m$ ), phenol, and chlorophenols.

The decomposed percentage of benzoic acid with  $\text{ClO}^-$  decomposed and the product yields based on the decomposed benzoic acid are listed in Table 1, where the yields of hydroxybenzoic acids and phenol include those of their chlorinated compounds. Both hydroxybenzoic acids and phenol react rapidly with  $\text{ClO}^-$  in the dark to give their chloro-derivatives.

As shown in Table 1, the conversion of  $\text{PhCOO}^-$  based on decomposed  $\text{ClO}^-$ , *i.e.* the ability of  $\text{ClO}^-$  to decompose  $\text{PhCOO}^-$ , is high when the ratio of  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0$  is low, *e.g.*, at  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0 = 0.01$ , the efficiency for the decomposition of  $\text{PhCOO}^-$  is 42% at 254 nm and 37% at  $>350$  nm, while at  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0 = 1.0$ , 24% at 254 nm and 19% at  $>350$  nm. This decrease of efficiency at the higher ratio of 1.0 may be ascribed to the consumption of  $\text{ClO}^-$  by the spontaneous decomposition of  $\text{ClO}^-$  induced by active species such as atomic O, Cl, and organic radicals before reacting with  $\text{PhCOO}^-$ .

The time dependence of product yield at  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0 = 0.1$  is shown in Table 3. The decomposition ratio of  $\text{PhCOO}^-$  versus  $\text{ClO}^-$  decreased with increasing irradiation time, *i.e.*  $\text{ClO}^-$  is photodecomposed spontaneously and by the reaction with  $\text{PhCOO}^-$ ; it can also react with hydroxylated products in the dark. Hence, as the reaction proceeds, the yield of hydroxybenzoic acid decreases, while that of chlorohydroxybenzoic acid increases, especially by  $>350$  nm irradiation. The yield of chlorinated compounds at 254

Table 1. Product yields of photoreaction of  $\text{PhCOOH}$  with  $\text{NaClO}$  in aqueous alkali ( $\text{pH} \geq 12$ ) at 20–21 °C<sup>a</sup>

Ratio <sup>b</sup>	$\lambda/\text{nm}$	$\text{PhCOOH}$ decomposed (%) <sup>c</sup>	Product yield (%) <sup>d</sup>			
			$\text{HOC}_6\text{H}_4\text{COOH}$	$\text{ClC}_6\text{H}_4\text{COOH}$	PhOH	Others
0.1	254	42.3	60 (5)	~1	<1	
0.5	254	30.2	64 (23)	1	1 (1)	
1.0	254	24.1	42 (37)	3	2 (2)	<i>e</i>
0.1	$>350$	36.8	71 (31)	5	5 (4)	<i>e</i>
0.5	$>350$	27.3	65 (41)	5	7 (7)	<i>e</i>
1.0	$>350$	19.1	64 (58)	4	6 (6)	<i>e</i>

<sup>a</sup> Initial concentration of benzoic acid  $[\text{PhCOOH}]_0$  was 0.01M. Irradiation time: 120 min at 254 nm, 40 min at  $>350$  nm.

<sup>b</sup>  $[\text{ClO}^-]/[\text{PhCOO}^-]_0$ . <sup>c</sup> Based on  $\text{ClO}^-$  decomposed. <sup>d</sup> Based on  $\text{PhCOOH}$  decomposed: the parentheses indicate chlorinated compound derived from the dark reaction with  $\text{ClO}^-$ . <sup>e</sup> Dichloro-derivatives were detected by g.l.c.

nm was lower than that at  $>350$  nm, which may be explained by secondary photolysis, *i.e.* chlorinated aromatic compounds are decomposed in aqueous alkali alone by absorbing the 254 nm irradiation, but they are stable to light of wavelength  $>350$  nm which is not absorbed. For example, chlorohydroxybenzoic acids were *ca.* 50% decomposed by 2 h irradiation at 254 nm; this photolysis was accelerated in the presence of  $\text{PhCOO}^-$ . On the other hand, no photolysis was observed at  $>350$  nm. Both chlorobenzoic acids and chlorophenols in aqueous alkali were decomposed by 254 nm light, *e.g.*, *ca.* 30% for chlorobenzoic acid and 10% for chlorophenol for 2 h irradiation. These percentages are lower than that of chlorohydroxybenzoic acid.

The yields of chlorinated products increase as the concentration of  $\text{ClO}^-$  increases (Table 1); this trend is explicable by the easy chlorination of hydroxylated aromatic compounds with  $\text{ClO}^-$  in the dark.

Benzene and phenylbenzoic acid were other products detected. Also the presence of  $\text{CO}_2$  was confirmed by the  $\text{Ba}(\text{OH})_2$  test. At high ratio of  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0$ , *e.g.*, over 20,  $\text{PhCOO}^-$  was completely decomposed to give  $\text{CO}_2$  together with acetic and oxalic acids and unknown products. The Figure shows the percentage formation of  $\text{CO}_2$  from  $\text{PhCOO}^-$  decomposed. The formation of  $\text{CO}_2$  was observed even at the initial stage. Hence  $\text{CO}_2$  is formed not only *via* complete oxidation, but also *via* substitution of the carboxylate group by radicals or an electron transfer from  $\text{PhCOO}^-$  to radicals.

**Mechanism of Initial Stage.**—Benzoate ion has  $\lambda_{\text{max}}$  at 224 nm ( $\epsilon$  8 350) and  $\text{ClO}^-$  has  $\lambda_{\text{max}}$  at 292 nm ( $\epsilon$  390). The 254 nm light is absorbed by both  $\text{PhCOO}^-$  and  $\text{ClO}^-$ , while the  $>350$  nm light is absorbed only by  $\text{ClO}^-$  because there is no u.v. absorption of  $\text{PhCOO}^-$  at  $>300$  nm.


**Table 2.** Isomeric distribution of dark reaction products of hydroxylated compounds with  $\text{ClO}^-$  in aqueous alkali ( $\text{pH} \geq 12$ ) at  $20^\circ\text{C}$  <sup>a</sup>

Hydroxylated compound	Yield (%) <sup>b</sup>	Isomeric distribution
2- $\text{HOC}_6\text{H}_4\text{COOH}$	87	3-Cl-2- $\text{HOC}_6\text{H}_3\text{COOH}$ (40), 5-Cl-2- $\text{HOC}_6\text{H}_3\text{COOH}$ (60)
3- $\text{HOC}_6\text{H}_4\text{COOH}$	82	2-Cl-3- $\text{HOC}_6\text{H}_3\text{COOH}$ (4), 4-Cl-3- $\text{HOC}_6\text{H}_3\text{COOH}$ (46), 6-Cl-3- $\text{HOC}_6\text{H}_3\text{COOH}$ (44)
4- $\text{HOC}_6\text{H}_4\text{COOH}$	89	3-Cl-4- $\text{HOC}_6\text{H}_3\text{COOH}$ (85), 3,5-Cl <sub>2</sub> -4- $\text{HOC}_6\text{H}_2\text{COOH}$ (15)
PhOH	93	2-Cl- $\text{C}_6\text{H}_4\text{OH}$ (80), 4-Cl- $\text{C}_6\text{H}_4\text{OH}$ (20)

<sup>a</sup> Initial concentration of hydroxy compound and  $\text{ClO}^-$ : 0.1M. Reaction time: 1 h for 3-, 4-hydroxybenzoic acids and phenol; 2.5 h for 2-hydroxybenzoic acid. <sup>b</sup> Based on  $\text{ClO}^-$  decomposed.


Active species generated on photolysis of aqueous  $\text{NaClO}$  were reported to be  $\text{O}(^3\text{P})$ ,  $\text{O}^{\cdot-}$ ,  $\text{O}(^1\text{D})$ ,  $\text{Cl}^\cdot$ , and  $\text{HO}^\cdot$ , the last one being derived from the reaction of water with  $\text{O}(^1\text{D})$  and  $\text{O}^{\cdot-}$ .<sup>4,5</sup> For a basic solution of pH 12, *ca.* 50% of  $\text{HO}^\cdot$  would be present as  $\text{O}^{\cdot-}$  because the  $\text{pK}_a$  of  $\text{HO}^\cdot$  is 11.9.<sup>7</sup> Atomic chlorine is less reactive and has a longer life than atomic oxygen of  $\text{HO}^\cdot$ .<sup>8</sup>

The quantum yields for three types of photodecomposition of  $\text{ClO}^-$  [equation (2)] are  $\phi_a$  0.08,  $\phi_b$  0.278, and  $\phi_c$  0.133 at 254 nm and  $\phi_a$  0.28 and  $\phi_b$  0.08 at 365 nm in aqueous alkali.<sup>4</sup> Hence the 254 nm photolysis of  $\text{ClO}^-$  should generate



$\text{O}(^3\text{P})$ ,  $\text{O}^{\cdot-}$ ,  $\text{O}(^1\text{D})$ , and  $\text{Cl}^\cdot$  as active species, while  $>350$  nm photolysis should give  $\text{O}(^3\text{P})$ ,  $\text{O}^{\cdot-}$ , and  $\text{Cl}^\cdot$ , but no  $\text{O}(^1\text{D})$ , although the distribution of active species may be different from that reported by Buxton.<sup>4</sup> The concentration of  $\text{Cl}^\cdot$  at 254 nm ( $\phi$  0.278) must be higher than that at  $>350$  nm ( $\phi$  0.08) based on the available quantum yield.<sup>4</sup>

Chlorobenzoic acid may be a primary product formed directly by the addition of atomic chlorine whereas chloro-








**Figure.** Plots of yield of  $\text{CO}_2$  versus percentage of decomposed  $\text{PhCOOH}$  at various ratios of  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0$  for the photo-reaction of  $\text{PhCOO}^-$  with  $\text{ClO}^-$ . Yield (%) of  $\text{CO}_2$  was calculated on the basis of the stoichiometry:  $\text{PhCOOH} \rightarrow 7 \text{CO}_2$ . (O) 254 nm light irradiation; (●)  $>350$  nm light irradiation.  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0$  25 (curve a), 20 (curve b), 10 (curve c).

**Table 3.** Time dependence of product yields of photoreaction of  $\text{PhCOOH}$  with  $\text{NaClO}$  in aqueous alkali ( $\text{pH} \geq 12$ ) <sup>a</sup>

$\lambda/\text{nm}$	$t/\text{min}$	$\text{ClO}^-$ decomposed (%)	$10^2 \text{PhCOOH}$ decomposed (mmol)	Product yield (%) <sup>b</sup>		
				$\text{HOC}_6\text{H}_4\text{COOH}$	$\text{ClC}_6\text{H}_4\text{COOH}$	PhOH
254	30	21	3.04	64 (3)	3	2 (1)
254	60	38	5.22	65 (2)	1	2 (2)
254	120	71	7.40	60 (5)	~1	<1
>350	10	28	3.38	78 (24)	5	6 (4)
>350	20	54	5.29	74 (29)	6	6 (5)
>350	40	92	7.20	71 (31)	5	5 (4)

<sup>a</sup>  $[\text{ClO}^-]_0/[\text{PhCOO}^-]_0$  0.1,  $[\text{PhCOO}^-]_0$  0.01M, volume of the solution is 200 ml. <sup>b</sup> Based on  $\text{PhCOOH}$  decomposed. Parentheses indicate chlorinated compounds derived from the dark reaction with  $\text{ClO}^-$ .



hydroxybenzoic acids and chlorophenols would be formed *via* the dark reaction of ring-hydroxylated compounds with NaOCl.

Hydroxybenzoic acids may be formed *via* addition of HO $\cdot$  or O( $^3$ P) to the aromatic ring followed by a hydrogen atom abstraction [equation (4)] in a manner similar to chlorination [equation (3)], where R $\cdot$  is any radical.

Since the quantum yield for the formation of Cl $\cdot$  ( $\phi_b$  0.278) is smaller than that of HO $\cdot$  ( $\phi_b + \phi_c$  0.411) as shown in equation (2), the yield of chlorobenzoic acid at 254 nm is much lower than that of hydroxybenzoic acids even if the direct photolysis of chlorobenzoic acid is taken into account. The lower yield of chlorobenzoic acid may also be ascribed to the lower reactivity of Cl $\cdot$  than those of HO $\cdot$  or O( $^3$ P),<sup>8</sup> because hydrogen abstraction and addition to the aromatic ring are similar in that the driving force for both reactions is pairing of odd electrons. Probably, Cl $\cdot$  may mainly act as a hydrogen atom abstractor R $\cdot$  from the adducts [equations (3) and (4)], but does not add to the aromatic ring.

Two pathways are probable for the formation of phenol: (a) *ipso*-attack of O( $^3$ P),<sup>5b</sup> but not HO $\cdot$ ,<sup>6b</sup> on the ring carbon bearing COO $^-$  [equation (5)] as reported for the reaction of toluene with O( $^3$ P)<sup>5b</sup> and (b) electron transfer of PhCOO $^-$  to radicals followed by decarboxylation and coupling with HO $\cdot$  [equation (6)].

The fact that the yield of phenol at >350 nm is higher than at 254 nm even at the initial stage of the photolysis agrees with the higher concentration of O( $^3$ P) at >350 nm than at 254 nm.<sup>4</sup> Hence O( $^3$ P) contributes largely to the formation of phenol. The formation of traces of phenylbenzoic acid and benzene implies the formation of phenyl radical as an intermediate. Since there is no direct photodecarboxylation, an electron transfer from PhCOO $^-$  to HO $\cdot$  and/or Cl $\cdot$  is conceivable. Unfortunately, the control runs for this electron transfer failed in a photoreaction of PhCOO $^-$  with H $_2$ O $_2$  or Cl $_2$  because of the rapid and spontaneous decomposition of H $_2$ O $_2$  even in the dark and also the rapid conversion of Cl $_2$  into ClO $^-$  in aqueous alkali of pH 12.

Since the yield of phenol parallels with the generation of O( $^3$ P) at various wavelengths and since traces of phenylbenzoic acid and benzene are formed, phenol may be formed *via* equations (5) and (6), *i.e.* *ipso*-attack of O( $^3$ P) on the

COO $^-$  group and one-electron transfer, the former being the main path for phenol formation.

Primary products thus formed, *i.e.* hydroxy- and chlorobenzoic acids and phenol, undergo further reactions with ClO $^-$  in the dark, direct photolysis, and/or reactions with active species formed from ClO $^-$  under irradiation, giving various products.

As shown in the Figure, the yield of CO $_2$  at a high ratio of [ClO $^-$ ] $_0$ /[PhCOO $^-$ ] $_0$ , *e.g.* 20, increases with the time of irradiation. As stated before, highly oxidised products, *e.g.* polyhydroxylated products, may be oxidised with evolution of CO $_2$  to degradation products and finally to acetic and oxalic acids, *etc.*

## Experimental

**Materials.**—Aqueous NaClO (*ca.* 1M) was prepared by introducing gaseous Cl $_2$  into aqueous NaOH (2.1M) at 0  $^\circ$ C and the concentration of ClO $^-$  was determined by iodometry. Benzoic acid was purified by recrystallisation from water, m.p. 121–122  $^\circ$ C. *o*- and *p*-chlorobenzoic acid were prepared by the KMnO $_4$  oxidation of the corresponding chlorotoluenes, m.p. 141–142  $^\circ$ C for the *ortho*- and 241–243  $^\circ$ C for the *para*-isomer.<sup>9</sup> Other reagents were of guaranteed grade.

**Apparatus.**—U.v. spectra were recorded of a Hitachi 124 spectrophotometer. M.p.s were measured by a Yanagimoto micro m.p. apparatus. pH was measured by a Hitachi-Horiba M7 pH meter. G.l.c. analyses were performed on a Yanagimoto G 180 gas chromatograph with flame ionisation detector using three sorts of columns (PEG 20M, KG-02, and Bentone 34). G.l.c.-mass spectral analyses were conducted with a JEOL LMS D-300 mass spectrometer. The measurement of the fluorescence of PhCOO $^-$  was done by a Hitachi 650-10S fluorescence spectrophotometer. A Halos low-pressure 60 W Hg lamp and a Phillips super-pressure 500 W Hg lamp were used as light sources.

**Photoreaction of Benzoate Ion with Hypochlorite Ion.**—A mixture of benzoic acid (0.01M) and sodium hypochlorite (0.001–0.200M) in aqueous NaOH (200 ml; pH  $\geq$  12) under nitrogen was irradiated using a Halos low-pressure 60 W

Hg lamp through a quartz filter in a quartz cell (254 nm irradiation) or a Phillips super-pressure 500 W Hg lamp through a naphthalene filter (50 mm; 0.5M) in a Pyrex cell (>350 nm irradiation). The concentration of  $\text{ClO}^-$  was measured by iodometry.

Unchanged  $\text{ClO}^-$  in a 10 ml portion was quickly decomposed by adding excess of aqueous  $2\text{M-Na}_2\text{S}_2\text{O}_3$  and the solution was then acidified with aqueous  $\text{H}_2\text{SO}_4$ . This work-up suppressed the further oxidation of hydroxylated products and the evolution of  $\text{Cl}_2$  on acidification. The solution thus treated was extracted with ether ( $3 \times 20$  ml), dried ( $\text{Na}_2\text{SO}_4$ ), and esterified by ethereal diazomethane. The reaction products were estimated and identified by g.l.c. and g.l.c.-mass spectral comparison with authentic samples. Another portion was acidified and  $\text{CO}_2$  was expelled by  $\text{N}_2$  bubbling and estimated by absorption in aqueous  $\text{Ba(OH)}_2$  followed by acidimetry with aqueous  $\text{HCl}$ .

**Dark Reaction.**—*o*-, *m*-, and *p*-hydroxybenzoic acids and phenol (0.01M) in aqueous alkali were oxidised by  $\text{ClO}^-$  (0.01M) within 1 h except for *o*-hydroxybenzoic acid. Work-up and identification were similar to those described in the photoreaction and products were identified by g.l.c. and g.l.c.-mass spectroscopy. Oxidation products and yields are shown in Table 2.

#### Acknowledgements

This study was supported by the Japanese Ministry of Education, Grant in Aid No. 543,024. We thank Mr. Y. Fujitani

for assistance with preliminary experiments and Toa Gosei Chem. Ind. Co. for a gift of chlorine.

#### References

- (a) S. K. Chakrabarty, 'Oxidation in Organic Chemistry,' ed. W. S. Trahanovsky, Academic Press, New York, 1978, Part C, ch. 5; (b) L. F. Fieser and M. Fieser, 'Reagents for Organic Syntheses,' Wiley, New York, vols. 1—6.
- Y. Kojima, *PPM*, 1973, No. 10, 42.
- (a) Y. Ogata, T. Suzuki, and K. Takagi, *J. Chem. Soc., Perkin Trans. 2*, 1978, 562; (b) Y. Ogata, T. Suzuki, and K. Takagi, *ibid.*, 1979, 1715; (c) M. Nakamura and Y. Ogata, *Bull. Chem. Soc. Jpn.*, 1977, **50**, 2396.
- G. V. Buxton and M. S. Subhani, *J. Chem. Soc., Faraday Trans. 1*, 1972, **68**, 958.
- (a) J. J. Havel and K. H. Chan, *J. Am. Chem. Soc.*, 1975, **97**, 5800; (b) J. S. Gaffney, R. Atkinson, and J. N. Pitts, Jr., *ibid.*, 1976, **98**, 1828.
- (a) Y. Ogata, K. Tomizawa, and Y. Yamashita, *J. Chem. Soc., Perkin Trans. 2*, 1980, 616; (b) G. W. Klein, K. Bhatia, V. Madhavan, and R. H. Schuler, *J. Phys. Chem.*, 1975, **79**, 1767.
- P. Neta, M. Z. Hoffman, and M. Simic, *J. Phys. Chem.*, 1972, **76**, 847.
- J. K. Kochi, 'Free Radicals,' Wiley, New York, 1973, vol. 1.
- H. T. Clarke and E. R. Taylor, *Org. Synth.*, 1943, Coll. Vol. 2, 135.

Received 23rd June 1983; Paper 3/1073