


This article was downloaded by: [The Aga Khan University]

On: 23 October 2014, At: 02:35

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

PREPARATION OF ALUMINA-SUPPORTED DIMETHYLAMMONIUM CHLOROCHROMATE (DMCC) AND ITS USE IN THE OXIDATION OF ALCOHOLS AND BENZOINS

Gui-Sheng Zhang ^a, Qi-Zeng Shi ^a, Mi-Feng Chen ^a & Kun Cai ^a

^a Department of Chemistry, Henan Normal University, Xinxiang, 453002, PR CHINA

Published online: 09 Feb 2009.

To cite this article: Gui-Sheng Zhang, Qi-Zeng Shi, Mi-Feng Chen & Kun Cai (1998) PREPARATION OF ALUMINA-SUPPORTED DIMETHYLAMMONIUM CHLOROCHROMATE (DMCC) AND ITS USE IN THE OXIDATION OF ALCOHOLS AND BENZOINS, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 30:2, 215-218, DOI: [10.1080/00304949809355282](https://doi.org/10.1080/00304949809355282)

To link to this article: <http://dx.doi.org/10.1080/00304949809355282>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

4-(*p*-Chlorophenyl)-3,6,8-tricyano-2-hydroxy-5,7-diphenyl-1H-quinoline Hydrochloride (11). A mixture of **7b** (0.01 mol, 4.49 g) and ammonium acetate (0.01 mol) in acetic acid (2 mL) was treated with ethyl cyanoacetate (0.01 mol, 1.13 g). The reaction mixture was refluxed for 3 h and allowed to cool at room temperature, then poured into ice-cold water, and 2-3 drops of conc. HCl was added. The solid was collected and recrystallized from a mixture of ethanol and dimethylformamide (1:1).

Acknowledgements. This work was financed by the University of Kuwait, Research Project SC 082. We are grateful to the University of Kuwait for financial support and to the General Facility Project in the Chemistry Department for analytical and spectral measurements.

REFERENCES

1. B. Basil and R. H. Kenneth, *Ger. Patent*, **2**, 627, 210 (1976); *Chem. Abstr.*, **86**, 139625b (1977).
2. W. J. Welstead Jr. and H. W. Moran, *US Patent*, 4,126, 635 (1978); *Chem. Abstr.*, **90**, 719244 (1979).
3. W. Zecher and F. Krohnke, *Chem. Ber.*, **94**, 690, 698, 707 (1961).
4. F. Krohnke and W. Zecher, *Angew. Chem.*, **74**, 811 (1962); *Angew. Chem. Int.*, **1**, 626 (1962).
5. F. Krohnke, *Synthesis*, **1** (1976).

PREPARATION OF ALUMINA-SUPPORTED DIMETHYLAMMONIUM CHLOROCHROMATE (DMCC) AND ITS USE IN THE OXIDATION OF ALCOHOLS AND BENZOINS


Submitted by
(07/26/97)

Gui-Sheng Zhang*, Qi-Zeng Shi, Mi-Feng Chen and Kun Cai

Department of Chemistry, Henan Normal University
Xinxiang 453002, P. R. CHINA

Chromium compounds have played an important role in the formulation of reagents adsorbed on inert inorganic supports as oxidants in organic synthesis.¹⁻⁷ These reagents oxidize a wide variety of alcohols to carbonyl compounds under mild reaction conditions and afford the products in high yields. However, they have the disadvantage of being photosensitive and unstable and can be stored for only several weeks under vacuum in the dark. We now report that dimethylammonium chlorochromate ($\text{Me}_2\text{NH}_2\text{CrO}_3\text{Cl}$, DMCC) adsorbed on alumina (DMCC/alumina) is a new reagent

suitable for oxidation of alcohols, benzoin and also sensitive compounds such as allyl alcohol and cinnamic alcohol to the corresponding carbonyl compounds. In addition, DMCC/alumina is more stable than the above reagents¹⁻⁷ and may be kept for several months in air at room temperature without losing its activity. The reagent is not flammable and is stable at temperatures below 135°; it will change to a black solid slowly at temperatures above 135°. The reagent is easily prepared by the addition of alumina to a solution of dimethylammonium chlorochromate in water and followed by rotary evaporation to dryness. The reaction is performed by simple stirring of a mixture of the alcohol and the oxidant in cyclohexane at a suitable temperature. The product is isolated by filtration of the reagent and removal of the solvent by distillation. The results obtained from the oxidation of eleven alcohols and five benzoin are reported in Table 1. The oxidation of other functional groups with this reagent is under investigation.



- a) R = Ph, R' = H b) R = Ph, R' = CH₃ c) R = PhCH=CH, R' = H d) R = 2-furyl, R' = H
 e) R = CH₂=CH, R' = H f) R = R' = Ph g) R = R' = -(CH₂)₅- h) R = R' = -(CH₂)₄-
 i) R = PhCH₂, R' = H j) R = CH₃(CH₂)₈, R' = H k) R = CH₃(CH₂)₆, R' = H
 l) R = Ph, R' = C₆H₅CO m) R = *p*-CH₃C₆H₄, R' = *p*-CH₃C₆H₄CO
 n) R = *p*-CH₃OC₆H₄, R' = *p*-CH₃OC₆H₄CO o) R = *p*-ClC₆H₄, R' = *p*-ClC₆H₄CO
 p) R = 2-furyl, R' = 2-furoyl

EXPERIMENTAL SECTION

Mps were determined on a micro-melting point apparatus and are uncorrected. IR spectra were measured on a Nicolet FT-IR spectrometer. The ¹H NMR spectra were obtained on a Varian EM-360 A (60MHz) spectrometer in CDCl₃ with TMS as the internal standard.

Preparation of DMCC/Alumina.- To a solution of chromium trioxide (10g, 0.1mol) in water (10mL) was added dimethylamine hydrochloride (8.2g, 0.1mol) over 15min at 40°. The mixture was then cooled until a yellow-orange solid formed and was then reheated to 40° to give a solution. Alumina (60g, 100-200 mesh) was then added to the solution with stirring at 40°. After evaporation on a rotary evaporator at 60°, the orange solid was dried in vacuum at 50° for 1 h. The reagent may be kept for several months in air at room temperature without losing its activity.

Oxidation of Benzyl Alcohol to Benzaldehyde. Typical Procedure.- To a solution of benzyl alcohol (1.4g, 13 mmol) in cyclohexane (15mL) was added 20.3g (26mmol) of DMCC/alumina in one portion. After stirring for 2 h at 30°, the orange solid, which became deep brown, was then filtered and washed with ether. The combined filtrates were evaporated and the residue was distilled *in vacuo* to afford 1.25g (91%) of benzaldehyde, bp. 62°/10 mmHg. Compounds **2b-2k** were obtained in a similar fashion (Table 1).

Oxidation of Benzoin to Benzil. Typical Procedure.- To a solution of benzoin (1g, 4.7mmol) in cyclohexane (15mL), the reagent (7.4g, 9.4mmol) was added and the mixture was stirred for 12 h at

60°. The orange solid, which became deep brown, was then removed by filtration and washed several times with ether. The combined filtrates were evaporated and the residue was recrystallized from EtOH to yield 0.89g (90%) of the product **2l** as a yellow needles, mp. 94-96°. IR (KBr): 1655, 1590, 1580, 795, 720, 690, 680 cm⁻¹; ¹H NMR (CDCl₃, 60 MHz): δ 7.15-7.90 (m, Ar-H).

Compounds 2m-2p were obtained in the same way; **2m**: IR (KBr):1655, 1590, 1565, 1450, 875, 835, 755, 730, 680 cm⁻¹; ¹H NMR (CDCl₃, 60 MHz): δ 2.40 (6H, s, 2CH₃), 7.28, 7.80 (8H, dd, *J* = 8Hz, Ar-H). **2n**: IR (KBr):1670, 1600, 1510, 880, 830 cm⁻¹; ¹H NMR (CDCl₃, 60 MHz): δ 3.80 (6H, s, 2CH₃), 6.90, 7.76 (8H, dd, *J* = 8Hz, Ar-H). **2o**: IR (KBr): 1650, 1590, 875, 820, 730 cm⁻¹; ¹H NMR (CDCl₃, 60 MHz): δ 7.41, 7.72 (8H, dd, *J* = 8.7Hz, Ar-H). **2p**: IR (KBr): 1650, 1400, 1290, 1030, 940, 810, 760 cm⁻¹; ¹H NMR (CDCl₃, 60 MHz): δ 6.60 (2H, m), 7.66 (4H, m).

TABLE 1. Oxidation of Alcohols with DMCC/alumina

Alcohols (1)	Temp. (°C)	Time (h)	Ratio of oxidant/1	Yield ^{a,b} (%)	mp. or bp./mmHg of 2 Found(°C)	lit.(°C)
1a	25-30	2	2	93(91)	61/10	62/10 ⁸
	55-60	1.5	2	94(89)		
1b	25-30	2	3	94(90)	83.5/12	83/12 ⁸
	55-60	1	3	96(89)		
1c	25-30	3	2	89(80)	130/20	130/20 ⁸
	55-60	2	2	87(78)		
1d	25-30	3	2	80(60)	68/20	68/20 ⁸
	55-60	2	2	70(40)		
1e	25-30	2	2	80 ^c		
1f	55-60	4	3	(84)	46.5-49	47-49 ⁸
1g	55-60	3	3	90(86)	48/15	47/15 ⁸
1h	55-60	3	3	75(65)	129/760	130/760 ⁸
1i	55-60	3	2	89(60)	78/10	78/10 ⁸
1j	55-60	3	2	90(81)	94.5/12	94/12 ⁸
1k	55-60	3	2	86(80)	71/20	72/20 ⁸
1l	60	12	2	(90)	94-96	92-94 ⁹
1m	60	13	2	(82)	102-104	103-104 ⁹
1n	58	14	2	(78)	132-134	130-133 ⁹
1o	56	11	2	(85)	195-197	195-196 ¹⁰
1p	55	13	2	76)	162-164	164-165 ¹¹

a) Yields determined by glc., unless otherwise noted. b) Figures in parentheses are yields of isolated product. All compounds isolated had identical spectral characteristics with the corresponding authentic samples. c) Determined as 2,4-DNP

Acknowledgment.- The Project 29577273 was supported by The National Natural Science Foundation of China.

REFERENCES

1. Y. S.Cheng, W. L.Liu and S. H. Chen, *Synthesis*, 223 (1980).
2. J. D. Lou and Y.Y.Wu, *Synth. Commun.*, **17**, 1717 (1987).
3. J. D. Lou, *ibid.*, **19**, 1841 (1989).
4. J. S. Filippo Jr and C. I. Chern, *J. Org. Chem.*, **42**, 2182 (1977).
5. J. M. Lalancette, G. Rollin and P. Dumas, *Can. J. Chem.*, **50**, 3058 (1972).
6. R. P. Singh, H. N. Subbarao and S. Dev, *Tetrahedron*, **35**, 1789 (1979).
7. E. Santaniello, F. Ponti and A. Manzocchi, *Synthesis*, 534 (1978).
8. J. Buckingham and S. M. Donaghy, "Heilbron's Dictionary of Organic Compounds". 5th ed. New York, (1982).
9. T. L. Ho, *Synthesis*, 697 (1972).
10. M. Gomberg and F. J. Van Natta, *J. Am. Chem. Soc.*, **51**, 2241 (1929).
11. A. Citterio, *Tetrahedron Lett.*, **30**, 2701 (1978).

**A SIMPLE REGIOSELECTIVE NUCLEAR
BROMINATION OF ALKYL PHENYL ETHERS**

Submitted by
(08/13/97)

Shigetaka Yakabe, Masao Hirano* and Takashi Morimoto*

*Department of Applied Chemistry, Faculty of Technology
Tokyo University of Agriculture and Technology
Koganei, Tokyo 184, JAPAN*

Oxohalogen acids and their organic and inorganic derivatives have been extensively employed as versatile, mild oxidizing and halogenating reagents in organic syntheses.¹ Sodium chlorite (NaClO_2 , **1**) has enjoyed an important position in carbohydrate chemistry² and is also known as a specific reagent for the oxidation of aldehydes to carboxylic acids.³ However, there has been no synthetically useful reactions other than the above oxidations. Indeed, reaction of aromatic hydrocarbons and of alcohols with aqueous NaClO_2 gave complex mixtures of products ascribed to the competing oxidation and chlorination.⁴ Our recent investigation into *cleaner* syntheses using *in situ*