

Direct Oxidation of Allylic Methylene to Carbonyl¹

By B. W. FINUCANE and J. B. THOMSON*

(Department of Chemistry, University College, Dublin 4, Ireland)


Summary Triterpene 12-enes with various substituents in ring A, taraxeryl acetate, and cholesteryl acetate are oxidised, essentially quantitatively, to the corresponding $\alpha\beta$ -unsaturated ketones by irradiated solutions of *N*-bromosuccinimide in moist solvents.

It has been reported² that 11-oxo-urs-12-en-3 β -yl acetate (II) is formed in *ca.* 80% crude yield by treatment of α -amyrin acetate (I) with *N*-bromosuccinimide in aqueous dioxan. In our hands, the reaction as described gave very variable yields (up to 60%) of the required ketone (II) together with a number of by-products which will be described elsewhere. However, we find that when the reaction mixture is irradiated with visible light, reproducible—essentially quantitative—yields of $\alpha\beta$ -unsaturated ketones are obtained from a number of compounds containing an allylic methylene group.

Thus, a solution of the olefin (1 mole) in a water-miscible, inert solvent† containing water (1–10%) is stirred (1 hr.) at room temperature with freshly crystallised *N*-bromosuccinimide (2.5 moles) and finely divided calcium carbonate (*ca.* 2 moles) while being irradiated with visible light. The mixture is filtered, diluted with an excess of water, and the product extracted into ether or chloroform.

In this way, α -amyrin acetate (I) gives 98% α -amyrenonyl acetate (II), m.p. 289–290°, $[\alpha]_D +96^\circ$; β -amyrin acetate yields 100% β -amyrenonyl acetate, m.p. 268–269°, $[\alpha]_D +102^\circ$; and taraxeryl acetate (III) affords 96% 16-oxo-taraxeryl acetate (IV), m.p. 251–252°, $[\alpha]_D +92^\circ$. Similar

yields of $\alpha\beta$ -unsaturated ketones are obtained from α - and β -amyrene, their 3-oxo- and 3 β -benzoyloxy-derivatives, and, surprisingly, their 3 β -ols (t.l.c. of the crude product shows only traces of 3,11-diketone). In the case of cholesteryl acetate the crude product (96%) shows traces of impurities on t.l.c. but is probably pure enough for most purposes. Crystallisation from chloroform–methanol gives 81% 7-oxocholesteryl acetate, m.p. 159–160°, $[\alpha]_D -98^\circ$.



Satisfactory analytical data were obtained for (IV); other products were compared with authentic samples.

We thank Biorex Laboratories Ltd. for financial support.

(Received, August 25th, 1969; Com. 1293.)

† *e.g.* Dioxan, tetrahydrofuran, glyme, or diglyme. The volume of the solution is adjusted so that the reaction mixture is translucent.

¹ Presented in part at the XXII Annual Irish Universities Chemistry Research Colloquium, New University of Ulster, Coleraine, May, 1969. Previous paper in this series: B. W. Finucane and J. B. Thomson, *Chem. Comm.*, 1969, 380.

² S. Corsano and G. Piancatelli, *Ann. Chim. (Italy)*, 1965, 55, 742.