

ARTICLE

NMR-Derived Models of Amidopyrine and Its Metabolites in Complexes with Rabbit Cytochrome P450 2B4 Reveal a Structural Mechanism of Sequential N-Dealkylation

Arthur G. Roberts,^{*,†} Sara E. A. Sjögren,[‡] Nadezda Fomina,[†] Kathy T. Vu,[§] Adah Almutairi,[†] and James R. Halpert[†]

[†]The Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, 9500 Gilman Drive, #0703, La Jolla, California 92093-0703, United States

^{*}Lund University, BMC F12, S-221 84 Lund, Sweden

⁹University of Gothenburg, Sahlgrenska Academy, Box 400, 405 30 Gothenburg, Sweden

Supporting Information

ABSTRACT: To understand the molecular basis of sequential N-dealkylation by cytochrome P450 2B enzymes, we studied the binding of amidopyrine (AP) as well as the metabolites of this reaction, desmethylamidopyrine (DMAP) and aminoantipyrine (AAP), using the X-ray crystal structure of rabbit P450 2B4 and two nuclear magnetic resonance (NMR) techniques: saturation transfer difference (STD) spectroscopy and longitudinal (T_1) relaxation NMR. Results of STD NMR of AP and its metabolites bound to P450 2B4 were similar, suggesting that they occupy similar niches within the enzyme's active site. The model-dependent



dent relaxation rates (R_M) determined from T_1 relaxation NMR of AP and DMAP suggest that the N-linked methyl is closest to the heme. To determine the orientation(s) of AP and its metabolites within the P450 2B4 active site, we used distances calculated from the relaxation rates to constrain the metabolites to the X-ray crystal structure of P450 2B4. Simulated annealing of the complex revealed that the metabolites do indeed occupy similar hydrophobic pockets within the active site, while the N-linked methyls are free to rotate between two binding modes. From these bound structures, a model of N-demethylation in which the N-linked methyl functional groups rotate between catalytic and noncatalytic positions was developed. This study is the first to provide a structural model of a drug and its metabolites complexed to a cytochrome P450 based on NMR and to provide a structural mechanism for how a drug can undergo sequential oxidations without unbinding. The rotation of the amide functional group might represent a common structural mechanism for N-dealkylation reactions for other drugs such as the local anesthetic lidocaine.

The mammalian cytochrome P450 (P450) 2B enzymes are responsible for the detoxification and elimination of a wide range of drugs from coumarins to amphetamines.¹ P450 catalyzed N-dealkylation is one of the main reaction pathways of these enzymes and occurs through a single electron oxidation or hydrogen abstraction mechanism.²⁻⁴ Drugs such as the antidepressant sertraline and the anorectic drug benzphetamine are N-dealkylated in a single step,^{4,5} while the local anesthetic lidocaine is sequentially N-dealkylated in a multistep process.⁶ The latter pathway leads to complicated metabolic profiles and distorted kinetics that often stymie predictions of in vitro and in vivo metabolism.^{7,8}

Studies of sequential P450-mediated N-dealkylation reactions have successfully identified and characterized intermediates within a number of metabolic pathways.⁹ Kinetic isotope effects,² redox potential,¹⁰ and radical probes^{3,11} have highlighted possible reaction mechanisms for the individual steps. X-ray crystal structures of P450 2B4¹ and P450 2B6¹² have helped to explain the broad substrate specificity of these enzymes and have also shown that these enzymes have considerable plasticity. Unfortunately, these X-ray crystal studies were limited to tight binding ligands, so the relationship between the X-ray crystal structures and weakly bound N-dealkylated P450 2B substrates remains less clear.

Therefore, the nonsteroidal anti-inflammatory drug (NSAID) amidopyrine (AP) was analyzed by NMR and used as a model compound for the N-dealkylation reactions. This drug was selected because it has been studied extensively in a variety of P450 2B enzymes and because the products and N-dealkylated intermediates can be easily obtained or synthesized.¹³⁻¹⁶ Additionally, AP and its metabolites have low affinities for the enzyme¹⁷⁻¹⁹ and are therefore likely to be in fast exchange with the P450 2B active site, which was a requirement for the NMR experiments.

Received:November 9, 2010Revised:January 31, 2011Published:February 08, 2011



 a (A) Amidopyrine (AP), (B) desmethylamidopyrine (DMAP), and (C) aminoantipyrine (AAP) are shown. The International Union of Pure and Applied Chemistry (IUPAC) numbering of the pyrazolone and the phenyl ring are shown in panel B.

Scheme 1 illustrates the N-demethylation reactions of AP by P450 2B enzymes.¹³ AP (Scheme 1A) is demethylated oxidatively to DMAP, which is shown in Scheme 1B. Because of weak binding, DMAP can either be released from the enzyme²⁰ or undergo a second demethylation resulting in aminoantipyrine [AAP (Scheme 1C)]. After being released from the P450 active site, AAP is acetylated by the cytosolic arylamine *N*-acetyltransferase (NAT2) to form acetylaminoantipyrine²¹ and eliminated.

The interactions of AP, DMAP, and AAP, which will be termed AP and its metabolites, with the rabbit P450 2B4 isoform were examined for the following reasons. First, the P450 2B4 enzyme has been the focus of a number of biochemical and biophysical studies (e.g., ref 22). Of the mammalian P450s, the most ligand-bound X-ray crystal structures have been determined for this enzyme (e.g., ref 23). Unlike the truncated constructs, which are available for most other mammalian P450s, the truncated variant of P450 2B4²⁴ used in this study is monomeric and soluble, making it more amenable to NMR analysis.

The interaction of AP and its metabolites with P450 2B4 was investigated using two NMR techniques: saturation transfer difference (STD) and longitudinal (T_1) relaxation. Distances calculated from the T_1 relaxation were used to constrain molecules within the active site of P450 2B4. A simulated annealing simulation of the complex was performed to ascertain the timedependent positions of the metabolites. This combination of results provided, for the first time, the basis of a structural model of P450 2B-catalyzed N-dealkylation reactions.

MATERIALS AND METHODS

Materials. Unless otherwise specified, all the chemicals were purchased from Sigma-Aldrich (St. Louis, MO). 4-Aminoantipyrine hydrochloride was purchased from Acros Organic (Geel, Belgium). The materials used for the protein purification were the same as described previously.^{12,23}

Synthesis of Desmethylamidopyrine (DMAP). The synthesis of DMAP is shown in Scheme S1 of the Supporting Information. The products of the reactions were analyzed by ¹H and ¹³C NMR using a JEOL 500 MHz NMR spectrometer (JEOL Ltd., Tokyo, Japan) and a 100 MHz Varian NMR spectrometer (Varian, Inc., Palo Alto, CA), respectively. First, the compound 4-aminoantipyrine hydrochloride (1.916 g, 8 mmol, 1) was dissolved in 80 mL of CH₃Cl, and diisopropyl ethylamine (1.8 mL, 8 mmol) was added dropwise. When the starting material was completely dissolved (3 min), acetic anhydride (1.51 mL, 16 mmol) and NiCl₂ (0.056 g, 0.24 mmol) were added. The reaction progress was monitored by thin-layer

chromatography (TLC) with a mobile phase comprised of a 9:1 mixture of ethyl acetate and methanol for 10 min. After the reaction was completed, the product was dried under vacuum. The acetylated product **2** was purified by flash chromatography, using a CombiFlash Companion System (Teledyne Isco, Inc., Lincoln, NE) on silica gel with ethyl acetate and methanol gradient. The yield of the resulting white powder was 1.916 g (98.5%): ¹H NMR (500 MHz, CDCl₃) δ (s, 1H), 7.48–7.43 (m, 2H), 7.36 (dd, *J* = 1.1, 8.4 Hz, 2H), 7.30 (dt, *J* = 1.1, 8.5 Hz, 1H), 3.08 (s, 3H), 2.21 (s, 3H), 2.07 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.86, 161.16, 152.19, 134.68, 129.08, 126.61, 123.97, 113.21, 35.35, 21.04, 13.89.

Compound 2 (0.98 g, 4 mmol) and NaH (60% oil by weight, 0.288 g, 4.8 mmol) were placed in a two-neck round-bottom flask under nitrogen, and dry THF was added. The reaction mixture was cooled to 0 °C, and methyliodide (0.3 mL, 4.8 mmol) was added. The reaction was brought to ambient temperature overnight. Solvent was removed under vacuum, and the methylated product **3** was purified by flash chromatography using a Combi-Flash Companion System on silica gel with an ethyl acetate and methanol gradient. The yield of **3**, which was a yellow oil, was 0.89 g (86%): ¹H NMR (100 MHz, DMSO- d_6) δ 7.53–7.94 (m, 2H), 7.37–7.75 (m, 3H), 3.13 (s, 3H), 2.96 (s, 3H), 2.22 (s, 3H), 1.83 (s, 3H); ¹³C NMR (400 MHz, DMSO- d_6) δ 170.86, 161.16, 152.59, 134.68, 128.08, 126.61, 123.97, 113.21, 135.35, 121.04, 13.99, 10.08.

Compound 3 (0.89 g, 3.4 mmol) was refluxed in 30 mL of water containing 0.57 mL (6.8 mmol) of HCl overnight. The solution was neutralized with NaHCO₃ and lyophilized. DMAP (4) was purified by flash chromatography using CombiFlash Companion System on a C-18 column with a H₂O and methanol gradient. The yield of 4 was 0.63 g (85%) and the purity of the resulting yellow oil was >95% as determined by NMR: ¹H NMR (500 MHz, CDCl₃) δ 7.49–7.40 (m, 4H), 7.24 (ddd, *J* = 1.4, 2.9, 7.5 Hz, 1H), 2.86 (s, 3H), 2.82 (s, 3H), 2.25 (s, 3H).

Protein Expression and Purification. The N-terminally truncated P450 2B4 with an H226Y mutation to prevent dimerization²⁵ was expressed in *Escherichia coli* TOPP3 cells as described previously.²³ Protein purification was conducted essentially as described in ref 12. The P450 concentration was measured using reduced P450 with CO (CO-reduced) difference spectra and a molar extinction coefficient of 91 mM⁻¹ cm^{-1.26}. The purity of the enzyme judged from sodium dodecyl sulfate—polyacrylamide gel electrophoresis (SDS—PAGE) and A_{417}/A_{280} ratios was >95%.

Binding Titrations of AP, DMAP, and AAP to P450 2B4. The UV–visible spectra of P450 2B4 in the presence of AP and its metabolites were measured with a S2000 single-channel charged-coupled device (CCD) rapid scanning spectrometer (Ocean

Optics, Inc., Dunedin, FL) with L7893 deuterium and a halogen light source with a fiber-optic cable (Hamamatsu, Inc., Bridgewater, NJ).²⁷ All titrations were performed in 500 mM potassium phosphate buffer (KP_i) (pH 7.4). Because of interference in the absorbance spectra from DMAP, titrations with the metabolite were performed at P450 2B4 concentrations of \geq 20 μ M in 0.5 or 0.1 cm path length cuvettes. The concentration-dependent titration curves were fit with the scientific analysis package Igor Pro 6.1 (Wavemetrics, Inc., Lake Oswego, OR) to the equation for the equilibrium of bimolecular association:²⁸

$$A = A_{\max} \frac{[E] + [L] + K_{\rm D} - \sqrt{([E] + [L] + K_{\rm D})^2 - 4[E][L]}}{2[E]} + \text{offset}$$
(1)

where A is the absorbance amplitude in units of molar extinction coefficient (μ M⁻¹ cm⁻¹), A_{max} is the maximal absorbance amplitude in units of molar extinction coefficient (μ M⁻¹ cm⁻¹), [E] is the enzyme concentration, [L] is the ligand concentration, K_D is the dissociation constant, and offset is a constant that compensates for small errors in the absorbance measurement but was always close to 0. In addition to fitting the titrations, the absolute absorbance spectrum of P450 2B4 with saturating concentrations of the AP and its metabolites was fit by a least-squares method using software written in the Python programming language (version 2.6) to a linear combination of low-spin (LS), LS with imidazole (LS_{imidazole}), high-spin (HS), and ferric P420 (P420) heme spectral standards as described previously.^{29,30} The LS_{imidazole} spectral standard represents the model heme spectrum for a nitrogenous ligand coordinated to the heme.

Longitudinal (T_1) Relaxation and Saturation Transfer Difference (STD) NMR of AP and Its Metabolites Bound to P450 2B4. NMR experiments were performed on a 600 MHz Bruker Avance III instrument (Bruker Daltonics Inc., Billerica, MA) with a three-channel (¹H, ¹³C, and ¹⁵N) cryoprobe at 25 °C. Relaxation delays of 15 and 30 s were used for the oxidized and reduced anaerobic samples, respectively.

 T_1 relaxation measurements were employed to orient metabolites of AP in the active site of P450 2B4 as described previously (e.g., refs 31 and 32). It was measured using a 180° inversion recovery pulse followed by WATER suppression by GrAdient Tailored Excitation (WATERGATE)³³ or excitation sculpting.¹² Details of the inversion recovery experiment are provided in ref 34. The inversion recovery pulse program and parameters were obtained from G. Wagner's pulse sequence library at http://gwagner.med.harvard.edu/intranet/nmr/wwwpslib/ GW pulse relax.html. For these experiments, P450 2B4 was dialyzed once against 500 mM KP_i (pH 7.4) and then twice against deuterated 500 mM KP_i (pH 7.4). NMR samples contained each of the molecules at 1.0 mM and 40 $\mu\mathrm{M}$ dialyzed P450 2B4 in 500 KP_i (pH 7.4) with 99% D_2O as the lock solvent. Paramagnetic samples with the P450 2B4 heme in the Fe³⁺ state were prepared by purging the sample under a flow of argon in a sealed 5 mm NMR tube for 2 h. Diamagnetic samples with the P450 2B4 heme in the Fe²⁺ state were obtained by gently bubbling the sample under CO for \sim 2 min and then reducing it with dithionite prior to purging with argon for 2 h. For both paramagnetic and diamagnetic samples, a BD anaerobic indicator strip (BD Biosciences, San Jose, CA) was used to ensure an anaerobic environment of the samples. Samples that became aerobic during the experiment,

as judged by the anaerobic indicator strip, were excluded from the analysis. Shifts in the NMR peaks of AP and its metabolites in the presence of oxidized and reduced P450 2B4 showed that the binding of these ligands was in fast exchange (see Results and Discussion and Figure S2 of the Supporting Information).

STD NMR has previously been used to probe the interactions of ligands with proteins by selective excitation of the protein that is transferred to the ligand.^{35,36} In this study, the technique was used to map the interactions of AP and its metabolites with P450 2B4. A saturation transfer pulse sequence was used³⁵ with a WATERGATE or excitation sculpting^{33,37} pulse sequence following the train of 50 ms saturation pulses. Details of the pulse sequence and parameters can be found in ref 35. Diamagnetic samples were prepared as described above, except that the samples contained 500 mM KP_i (pH 7.4) with 10% D₂O. The percent of STD (%STD) transfer was calculated by dividing the amplitude of the STD NMR peaks against the analogous peaks in the WATERGATE NMR spectrum under the same conditions. The %STD was monitored over time and fit to a single-exponential function in Igor Pro 6.1.

Processing, Background Subtraction, and Deconvolution of NMR Spectra. NMR data were processed using the NMR processing software package iNMR (http://www.inmr.net). NMR peaks of AP and its metabolites were initially assigned using the ¹H proton NMR spectrum of AP in CDCl₃ from the Spectral DataBase System for organic compounds (SDBS) and using NMR Predictor.³⁸ The final NMR peak assignments were determined using the NMR peak positions, the integrated intensities of the NMR peaks, and the splitting of the NMR peaks. The NMR spectra were then translated into ASCII text format and imported into the scientific analysis package Igor Pro 6.1. The background and baseline were subtracted from the NMR spectra using software written in the Python programming language and integrated into Igor Pro 6.1. The NMR spectra were then fit to a linear combination of Gaussian, Lorentzian, or Voigt line shapes $^{39-42}$ using the multiple-peak fitting package of Igor Pro 6.1 and a least-squares fitting program written in Python using the numpy Python module. These fitted curves were used as standards for the singular-value decomposition (SVD) analysis.

Singular-Value Decomposition (SVD) Analysis for the T_1 Relaxation and STD NMR Experiments. SVD analysis with standards was used to measure the amplitudes of the NMR peaks from the T_1 relaxation and STD NMR experiments.^{43–46} The SVD algorithm was written in Python using the numpy Python module and integrated into Igor Pro 6.1. This method gauged the peak amplitude more accurately than measuring the peak amplitude directly or integrating the peak. The SVD-derived amplitudes for the T_1 relaxation and STD experiments were fit to a single-exponential function in Igor Pro 6.1. The derived fitting parameters provided the %STD_{max} for the STD experiments and the relaxation rates for the T_1 relaxation experiments.

Calculating the Apparent Distances (r_{app}) from the T_1 Paramagnetic Relaxation Rates (R_P). The T_1 paramagnetic relaxation rate (R_P) was experimentally determined by measuring the differences between the NMR longitudinal relaxation rate between oxidized P450, where the heme was in the Fe³⁺ state, and reduced P450, where the heme was in the Fe²⁺ state. This relationship can be represented mathematically by the equation $R_P = R_{Fe^{3+}} - R_{CO-Fe^{2+}}$, where R_P is the paramagnetic relaxation rate, $R_{Fe^{3+}}$ is the relaxation rate of the oxidized P450, and $R_{CO-Fe^{2+}}$ is the relaxation rate of CO-reduced P450. The

Tab	le 1	. I	Paramagnetic	Relaxation	Rates and	Calcu	lated	Distances"
-----	------	-----	--------------	------------	-----------	-------	-------	------------

peak	$R_{\rm Fe^{3+}} (\rm s^{-1})$	$R_{\rm CO-Fe^{2+}}(s^{-1})$	$R_{\rm P}~({ m s}^{-1})$	$R_{\rm M} ({\rm s}^{-1})^{b,c}$	r_{app}^{c} (Å)	$r_{\min}^{c,d}$ (Å)							
1.0 mM Amidonyrine and 40 µM D450 2B4													
2'/6'	0.633 ± 0.004	0.379 ± 0.004	0.254 ± 0.006	38.11 ± 1.55	10.88 ± 0.07	9.69 ± 0.07							
3'/5'	0.901 ± 0.008	0.415 ± 0.007	0.486 ± 0.011	72.89 ± 2.91	9.76 ± 0.06	8.70 ± 0.06							
4′	0.826 ± 0.014	0.392 ± 0.011	0.434 ± 0.017	32.57 ± 1.70	9.95 ± 0.09	9.95 ± 0.09							
1-CH3	0.758 ± 0.011	0.654 ± 0.013	0.104 ± 0.017	23.40 ± 3.95	12.62 ± 0.36	10.51 ± 0.30							
4-CH3	1.205 ± 0.015	0.971 ± 0.009	0.234 ± 0.017	105.28 ± 8.54	11.02 ± 0.15	8.18 ± 0.11							
5-CH ₃	0.893 ± 0.016	0.758 ± 0.017	0.135 ± 0.023	30.43 ± 5.37	12.08 ± 0.36	10.06 ± 0.30							
1.0 mM Desmethylamidopyrine and 40 μ M P450 2B4													
2'/6'	0.507 ± 0.003	0.360 ± 0.003	0.147 ± 0.005	30.81 ± 4.51	11.96 ± 0.29	10.65 ± 0.26							
3'/5'	0.675 ± 0.004	0.390 ± 0.002	0.285 ± 0.005	59.83 ± 8.60	10.71 ± 0.26	9.54 ± 0.23							
4′	0.606 ± 0.008	0.351 ± 0.004	0.255 ± 0.009	26.78 ± 3.95	10.91 ± 0.27	10.91 ± 0.27							
1-CH ₃	0.699 ± 0.007	0.623 ± 0.010	0.075 ± 0.012	23.74 ± 5.12	13.36 ± 0.48	11.13 ± 0.40							
4-CH ₃	0.832 ± 0.004	0.581 ± 0.005	0.251 ± 0.007	79.14 ± 11.49	10.93 ± 0.26	9.10 ± 0.22							
5-CH3	0.904 ± 0.011	0.710 ± 0.015	0.194 ± 0.019	61.09 ± 10.51	11.42 ± 0.33	9.51 ± 0.27							
1.0 mM Aminoantipyrine and 40 μ M P450 2B4													
2'/6'	0.687 ± 0.004	0.400 ± 0.004	0.286 ± 0.006	87.43 ± 7.39	9.31 ± 0.13	8.29 ± 0.12							
3'/5'	0.899 ± 0.006	0.417 ± 0.004	0.483 ± 0.007	147.19 ± 12.28	8.53 ± 0.12	7.60 ± 0.11							
4′	0.737 ± 0.009	0.406 ± 0.006	0.331 ± 0.011	50.51 ± 4.48	9.09 ± 0.13	9.09 ± 0.13							
1-CH ₃	0.701 ± 0.003	0.572 ± 0.015	0.130 ± 0.015	59.25 ± 8.53	10.63 ± 0.25	8.85 ± 0.22							
5-CH ₃	0.943 ± 0.005	0.652 ± 0.012	0.290 ± 0.013	132.76 ± 12.50	9.29 ± 0.15	7.74 ± 0.12							

^{*a*} Abbreviations: $R_{Fe^{3+}}$, paramagnetic relaxation rate of oxidized P450 2B4; $R_{CO-Fe^{2+}}$, paramagnetic relaxation rate of reduced P450 2B4 with CO; R_{P} , paramagnetic relaxation rate (i.e., $R_{P} = R_{Fe^{3+}} - R_{CO-Fe^{2+}}$). The distances were calculated with a τ_{C} of 3 × 10⁻¹⁰ s, which is the average value of P450 species under variety of conditions.^{31,49}. ^{*b*} R_{M} was calculated using eq 4. ^{*c*} The K_{D} values were obtained from the UV—visible absorbance in Figure 2, and the spin state was determined by least-squares fitting in Figure 1. The average and the standard deviation were from three independent experiments. ^{*d*} The minimal distance calculated on the basis of the chemical equivalence.

٦

model-dependent relaxation rate ($R_{\rm M}$) is dependent on the fraction of ligand bound to the protein and can be expressed mathematically as $R_{\rm P} = \alpha R_{\rm M}$, where α is the fraction of ligand bound with respect to total ligand concentration ([L]) and $R_{\rm M}$ is the model-dependent relaxation rate. For a single-ligand binding model, α is equal to [E]/($K_{\rm D}$ + [L]). The paramagnetic relaxation rate is related to distance by the Solomon–Bloembergen equation (SI units):^{34,47–50}

$$R_{\rm P} - \alpha \frac{2}{15} \left(\frac{\mu_0}{4\pi} \right) \frac{\gamma_{\rm N}^2 g_{\rm e}^2 \mu B^2 S(S+1)}{r_{\rm app}^6}$$

Г

$$\left[\frac{\tau_{\rm C}}{1 + (\omega_{\rm N} - \omega_{\rm E})^2 \tau_{\rm C}^2} + \frac{3\tau_{\rm C}}{1 + \omega_{\rm N}^2 \tau_{\rm C}^2} + \frac{6\tau_{\rm C}}{1 + (\omega_{\rm N} - \omega_{\rm E})^2 \tau_{\rm C}^2}\right]$$
(2)

where μ_0 is the magnetic permeability of free space, γ_N is the nuclear gyromagnetic ratio, g_e is the electronic g factor, μ_B is the Bohr magneton, r_{app} is the apparent time-averaged electron–nuclear distance, τ_C is the correlation time for the nuclear–electron interaction vector, which is dominated by the electron spin relaxation rates, S is the electronic spin quantum number, and ω_N and ω_E are the radial frequencies of the nucleus and electron, respectively. The S(S+1) term accounts for the fractional spin state of the P450. For mixed spin state systems, ⁵¹ S(S+1) is equal to $8.75f_{HS} + 0.75f_{LS}$, where f_{HS} and f_{LS} are the fractions of high-spin (HS) and low-spin (LS) species, respectively.

Equation 2 neglects the chemical equivalence of the protons with the implicit assumption that all protons associated with an NMR peak are exactly the same distance from the heme. Chemical equivalence has been treated previously for parallel and perpendicular orientations of P450-bound aromatic functional groups^{51,54} but never generalized for P450s. The relationship between the r_{app} and the distances of the individual protons is shown below, with *n* and r_n representing the number and the distances of those chemically equivalent protons, respectively.

$$r_{\rm app}^{-6} = \frac{1}{n} \sum_{0}^{n} r_n^{-6} \cong \frac{1}{n} r_{\rm min}^{-6}$$
(3)

 r_{\min} is the minimum distance of a proton from a group of chemically equivalent protons. In eq 3, r_{\min}^{-6} is approximately equal to the sum of r_n^{-6} , because longer distances make a negligible contribution to the sum. The relationship between $R_{\rm P}$ and $R_{\rm M}$ is simply

$$R_{\rm P} = \frac{1}{n} \, \alpha R_{\rm M} \tag{4}$$

Effect of Ligand Dynamics on R_P , r_{app} , and r_{avg} . Because of ligand dynamics and the fast exchange requirement for these measurements, the distances calculated from this analysis are not likely to represent absolute distances, but time-averaged distances weighted toward the shortest distance. In this study, the time averaging of distances of mobile nuclei and paramagnetic species was accomplished using the ensemble approach.^{46,55,56} The time-dependent ensemble averaging equations of R_P , R_M , and r_{app} are shown in eq 5, where Δt_n is the fraction of time, t is the total time, and f_t is the fraction of time at an individual



Figure 1. Least-squares fitting of the absolute absorbance spectra of P450 2B4 with AP, DMAP, AAP, and BP. The spectra (gray lines), fits (thick black lines), and standards (dotted lines) are shown with (A) no ligands, (B) 256μ M BP, (C) 15 mM AP, (D) 24 mM DMAP, and (E) 15 mM AAP. The P450 2B4 concentration was 74 μ M for the spectra with DMAP and 5 μ M for the rest of the samples. The high-spin (HS), low-spin (LS), ferric P420 (P420), and LS with imidazole (LS_{imidazole}) UV-visible heme absorbance standards are labeled.

distance. The left part of eq 5 shows that $R_{\rm p}$ is linearly related to the individual relaxation rates, whereas $r_{\rm app}$ has r^6 dependence in the right part. This will skew $r_{\rm app}$ toward the shortest distance, whereas $R_{\rm p}$ and $R_{\rm M}$ will remain linear. For a molecule that fluctuates, the average distance ($r_{\rm avg}$, eq 6) is quite different from $r_{\rm app}$, which is shown below. Like $R_{\rm p}$ and $R_{\rm M}$, $r_{\rm avg}$ is linear with respect to the individual distances (r_n) in contrast to $r_{\rm app}$. To demonstrate the effect of time averaging on $r_{\rm app}$, a simulation of $r_{\rm app}$, $r_{\rm avg}$ and $R_{\rm p}$ is presented in the Supporting Information.

$$R_{\rm P} = \alpha R_{\rm M} = \sum_{0}^{n} \frac{\Delta t_n}{t} R_n = \sum_{0}^{n} f_t R_n \propto r_{\rm app}^{-6}$$
$$= \sum_{0}^{n} \frac{\Delta t_n}{t} r_n^{-6} \sum_{0}^{n} f_t r_n^{-6}$$
(5)

$$r_{\rm avg} = \sum_{0}^{n} \frac{\Delta t_n}{t} r_n = \sum_{0}^{n} f_t r_n \tag{6}$$

Molecular Docking Using Distance-Restrained Simulated Annealing of AP and Its Metabolites to P450 2B4. Distancerestrained simulated annealing, which is a type of molecular dynamics (MD) simulation, is a tool commonly employed in the refinement of NMR structures and complexes.^{57,58} The technique was used here to determine the preferred orientations of the molecules in the active site of the X-ray crystal structure of P450 2B4 (Protein Data Bank entry 1SUO⁵⁹) using the distances calculated from the R_P values as described for other systems.^{60–62} Residues that were missing from the X-ray crystal structure were added using the homology modeling program Modeler⁶³ and the amino acid sequence of the truncated construct of P450 2B4.⁵⁹ AP and its metabolites were each then manually docked into the X-ray crystal structure of P450 2B4 without ligands using the PyMOL Molecular Graphics System, version 1.2r2 (Schrödinger, LLC). Distance-restrained simulated annealing and energy minimization of P450 2B4 and the substrates were performed with Groningen MAchine for Chemical Simulation (GROMACS), version 4.07.64 The complexes of AP and its metabolites and P450 2B4 were simulated with the GROMOS 96 53a6 force field.⁶⁵ A derivation of the force field parameters and charges of the heme with a spherical iron in the GROMOS 53a6 force field are described in ref 66. The particle-mesh Ewald⁶⁷ method was used for electrostatics with positional restraints placed on the α carbon backbone during the simulated annealing. The molecules were restrained to P450 2B4 using the NMR calculated distances from Table 1. A time constant of 50 ps was applied to the distance restraint to allow the molecule to move in the active site during the simulation, because the calculated distances reflect multiple orientations of the substrates in the P450 2B4 active site. Prior to the simulation, the molecule was energy minimized using the steepest descent method to eliminate van der Waals contact. The system was then heated to 800 K and cooled to 300 K in 150 ps and allowed to equilibrate for 100 ps at 300 K. Snapshots of the molecules bound to P450 2B4 were taken during the last 50 ps of the MD simulation and used to show the binding modes of the molecules.

RESULTS AND DISCUSSION

UV–Visible Spectroscopy of P450 2B4 in the Presence of AP and Its Metabolites. The P450 heme absorbance spectrum is sensitive to ligand binding and solution conditions and is composed of overlapping spectra of the HS $(+^{5}/_{2})$ and LS



Figure 2. Analysis of UV—visible absorbance changes of P450 2B4 in the presence of (A) BP, (B) AP, (C) DMAP, and (D) AAP. Difference spectra and binding curves of the P450 2B4 heme with BP, DMAP, and AAP are shown in the left and right panels, respectively. Panel B shows the effect of AP on the K_D of BP (\blacksquare) with an extrapolated line used to estimate the K_D . The concentration of P450 2B4 was 5 μ M in the presence of AP and AAP. Because of interference from the DMAP absorbance, the UV—visible spectra were recorded with 20 μ M P450 2B4 and 74 μ M P450 at <24 and 24 mM DMAP, respectively. The binding curves in the right panels were made by monitoring the maximum heme absorbance change as a function of ligand concentration.

 $(+^{1}/_{2})$ states of the heme. The fraction of LS and HS was estimated using heme absorbance standards as described in Materials and Methods.^{29,30} The binding affinity was determined directly by monitoring ligand-induced heme absorbance changes with absorbance difference spectra^{26,68} or indirectly by analyzing competition with benzphetamine (BP).⁶⁹

Figure 1 shows the absolute absorbance spectra of P450 2B4 with fits to absorbance standards in the presence of a saturating amount of AP, DMAP, AAP, or BP. In all cases, the correlation coefficient was >0.997, indicating very accurate fits. The concentration of P420 by fitting was <20% in all the samples. In the absence of ligands, the %LS and %HS determined from the absolute absorbance spectrum in Figure 1A were 77 and 23%, respectively. In the presence of a saturating amount of BP (Figure 1B), %HS increased by 20% to 43%. The increases in %HS generally indicate displacement of the heme-bound water.⁷⁰ In contrast, AP induces very little change in the spin state of the P450 2B4 heme. The %LS increased by <2% to 79%. It is unclear from the analysis whether the small increase in %LS is due to stabilization of a heme-bound water, due to weak interaction with AP, or merely an artifact of fitting. The effects of



Figure 3. 1D ¹H NMR spectrum of 1.0 mM DMAP in 99% D_2O . The peaks are labeled according to Scheme 1B.

a saturating amount of DMAP on the P450 2B4 heme absorbance were more dramatic. The fit of the absolute absorbance spectrum revealed an increase of 11% in the HS state to 34%, indicating displacement of heme-bound water. The best fits of the absolute absorbance spectrum of P450 2B4 in the presence of the product AAP were obtained with LS, $LS_{imidazole}$, and HS standards. Because of similarities between the $LS_{imidazole}$ and P420 spectral standards, the P420 concentration could not be accurately measured in this case. Therefore, the %LS should be considered approximate. %LS and %LS_{imidazole} were approximately 41% each, indicating an incomplete shift to 82% LS by weak interaction of the ligand⁷¹ or stabilization of the hemebound water complex.⁷²

Figure 2 shows the UV—visible analysis of P450 2B4 in the presence of AP, DMAP, AAP, or probe substrate BP. BP caused an increase at 386 nm and a decrease at 421 nm in the difference spectra (Figure 2A) that were characteristic of a type I absorbance difference spectra.⁶⁸ The resulting binding curve was fit to eq 1 with a K_D of $34 \pm 10 \ \mu$ M and a ΔA_{max} of $0.014 \pm 0.003 \ \mu$ M⁻¹ cm⁻¹, which is in line with previously estimated K_D values for BP of $40-121 \ \mu$ M.^{4,22,73}

In contrast, the absorbance changes of P450 2B4 in the presence of AP were quite minute (Figure 1C), so the binding of AP was analyzed indirectly using BP as a probe substrate. Figure 2B shows the effect of AP on the K_D of BP. Increasing the AP concentration causes a linear increase in the K_D of BP to 1.7 ± 0.2 mM at 64 mM AP. By linear extrapolation through the points, a K_D of 2.0 \pm 0.1 mM was estimated, which correlates well with K_M values obtained from AP kinetics of ~ 2 mM.^{17,18}

The absorbance difference spectra of P450 2B4 in the presence of DMAP are shown in the left panel of Figure 2C. DMAP induced a type I absorbance change with increases and decreases at 383 and 419 nm, respectively.⁶⁸ Fitting the binding curve to eq 1 gives a K_D and a ΔA_{max} of 3.2 ± 0.6 mM and 0.015 ± 0.008 μ M⁻¹ cm⁻¹, respectively, very similar to values determined previously.¹⁹

The difference spectra of P450 2B4 in the presence of AAP showed a decrease at 411 nm and an increase at 434 nm that were characteristic of type II absorbance difference spectra.⁶⁸ A $K_{\rm D}$ of 2.6 ± 0.4 mM and a $\Delta A_{\rm max}$ of 0.0080 ± 0.0002 μ M⁻¹ cm⁻¹ were estimated from fitting the binding curve.

NMR Assignment of AP and Its Metabolites. Figure 3 shows the one-dimensional (1D) ¹H NMR spectrum of DMAP showing seven well-defined NMR peaks. Three NMR peaks, two triplets and one doublet, in the aromatic region of the spectrum



Figure 4. Saturation transfer difference (STD) of AP, DMAP, and AAP with P450 2B4. (A) STD NMR spectrum of 2B4 with AP. (B) %STD transfer of the 4-CH₃ NMR peak as a function of saturation time. (C) % STD_{max} vs proton peaks for AP (black), DMAP (gray), and AAP (white).

between 7 and 8 ppm were assigned to the phenyl ring of DMAP. One triplet was centered at 7.60 ppm (J = 7.4 Hz), while the other triplet was centered at 7.55 ppm (J = 7.3 Hz) with half the peak area of the first triplet. From the splitting and the relative area of the NMR peak centered at 7.6 ppm, it was assigned to the protons at positions 3' and 5' on the phenyl ring. This triplet is most likely a doublet of doublets, because inequivalent nuclei flank the protons at these positions. By process of elimination, the peak at 7.55 ppm was assigned to the proton at position 4'. The remaining protons at positions 2' and 6' of the phenyl ring were then assigned to the doublet centered at 7.39 ppm (J = 7.4Hz). In contrast, the methyl groups of the pyrazolone ring formed singlets in the NMR spectrum. The peak areas of these singlets were virtually identical. No additional peaks reflecting contamination from the amidopyrine metabolites were observed. From the NMR spectrum of AP in the SBDS database, the

methyl groups at positions 1 and 5 were assigned to 3.14 and 2.30 ppm, respectively. The remaining methyl at position 4, which is metabolized by P450 2B4, was assigned to the NMR peak centered at 2.64 ppm and is upfield of the analogous peak in the AP NMR spectrum that was centered at 2.68 ppm in this study (data not shown) and at 2.784 ppm in the SDBS database. The amide proton of the desmethylamido group was absent from the NMR spectrum because of exchange broadening. NMR spectra of AP and AAP were virtually identical, with small deviations in the peak positions, so they were given the same relative assignments (data not shown).

Saturation Transfer Difference (STD) NMR To Probe the Interaction of AP and Its Metabolites to Diamagnetic P450 2B4. Saturation transfer difference (STD) NMR has been used to probe ligand—protein interaction for drug screening, epitope mapping, and fragment-based drug design for ligands with K_D values in the tens of nanomolar to millimolar range.^{35,74,75} The efficiency and the degree of saturation transfer depend on the orientation and position of the ligand with respect to the protein. In this study, STD NMR was used to probe the interaction of the molecules with CO-reduced P450 2B4 and was complementary to the T_1 relaxation measurements.

Figure 4 shows the analysis of AP, DMAP, and AAP by STD NMR. The STD NMR spectrum of DMAP with a 2 s saturation is shown in Figure 4A. The STD NMR spectrum looked distinctly different from the ¹H proton NMR spectrum shown in Figure 3. These differences indicated specific interactions between DMAP and P450 2B4. Most noticeably, the methyl NMR peaks numbered 1, 4, and 5 were no longer of equal amplitude. The N-linked methyl at position 1 had the strongest STD signal over the other two peaks. To determine the maximal amount of saturation, %STD was monitored as a function of time. Figure 4B shows %STD for the NMR peak of the N-linked methyl at position 4. Because the maximal %STD (%STD_{max}) was not achieved with a 2 s saturation and a longer saturation could potentially damage the cryoprobe, the %STD_{max} was estimated by fitting to a single-exponential function. The function fits to the curve with a saturation transfer rate of $\sim 0.5 \text{ s}^{-1}$ and a STD_{max} of $6.3 \pm 0.3\%$.

Figure 4C shows the $\$STD_{max}$ values for the NMR peaks of AP, DMAP, and AAP. The relative amplitudes of $\$STD_{max}$ were similar for all the ligands, suggesting that they occupy similar niches within the P450 2B4 active site. The $\$STD_{max}$ was highest for the N-linked methyl at position 1 on the pyrazolone ring for all the molecules with values between 30 and 40%. In contrast, the other methyls had the weakest interaction with a $\$STD_{max}$ of <12.6%. The saturation transfer between the phenyl ring protons and P450 2B4 was intermediate with a $\$STD_{max}$ between 12 and 20%.

 T_1 Relaxation NMR Was Used To Estimate the Orientation of AP and Its Metabolites in the P450 2B4 Active Site. A waterfall representation of individual 1D proton NMR spectra that comprise the T_1 relaxation NMR experiment (Figure 5) shows that with short inversion recovery times the peaks are positive but decrease as the inversion recovery time is increased. At long inversion recovery times, the absolute amplitude is similar to that at short inversion recovery times, but in the opposite direction. In Figure 5B, the 1D proton NMR spectrum with an inversion recovery time of 10 s was fit by the least-squares method to NMR peak standards (e.g., ref 76). In all cases, the residuals from the data and the fit were minute and the correlation coefficients were ≥ 0.98. Using the simulated peaks and the



Figure 5. NMR relaxation of 1 mM DMAP in the presence of 40 μ M reduced P450 2B4. (A) Waterfall graph of the 1D proton NMR spectra at various inversion recovery times with the background and baseline subtracted. (B) 1D proton NMR spectra at an inversion recovery time of 10 s (10 s) with least-squares fit (fit) and residual (res). (C) SVD analysis of the N-linked methyl peak at position 4 as a function of inversion recovery time.

NMR spectra, the inversion recovery time was analyzed using the SVD method and fit to a single-exponential function in Figure 5C with a χ^2 of 6.4 \times 10⁻⁴.

Table 1 shows the relaxation rates and calculated distances that were determined from the SVD analysis. The first three columns show the $R_{\text{Fe}^{3+}}$, $R_{\text{CO}-\text{Fe}^{2+}}$, and the difference, which is R_{P} . Because of chemical equivalence, R_{M} values in column 4 were calculated from the R_{P} using eq 4. The rates varied as much as 5-fold between protons. Using the Solomon–Bloembergen equation (eq 2) and treatment described in Materials and Methods, r_{app} and r_{min} were calculated and are shown in the last two columns. Taken together, the distances can vary as much as 4 Å depending on the orientation of the ligands in the active site. Because of the r^6 distance dependence, chemical equivalence, and time averaging, the r_{app} from the relaxation will be less dispersed and less sensitive to orientation than R_{M} .

Because of the higher sensitivity of R_M with respect to the orientation and position of AP and its metabolites, the R_M of these molecules was analyzed in Figure 6 with respect to the 4' proton of the phenyl ring. Positive and negative values indicate that the proton is closer and farther from the heme than the 4' proton, respectively. Large differences in $\%\Delta R_M$ for some of the protons on the phenyl and pyrazolone rings suggest that at least one axis of each of these groups was perpendicular to the heme. Consistent with metabolism of the N-linked methyls of AP and DMAP, this functional group was closest to the heme with relaxation rates that were 50–200% greater than R_M of protons



Figure 6. Relative percent differences in the R_M ($\%\Delta R_M$) of AP (black), DMAP (gray), and AAP (white). $\%\Delta R_M$ was calculated with respect to the 4' proton of the phenyl ring.

on the phenyl ring. For these molecules, the $\%\Delta R_M$ values of the 2'/6' proton and the methyl at position 1 were also similar, indicating that they were relatively parallel to the 4' proton. In contrast, the $\%\Delta R_M$ of the 3'/5' proton and the methyl at position 5 of DMAP was parallel like AAP, while the methyl at position 5 was farther from the heme for AP. Unfortunately, because of chemical equivalence and time averaging, this analysis provided no information about how these molecules are specifically sequestered in the P450 2B4 active site or its relationship to the STD NMR.

Models of AP and Its Metabolites Bound to P450 2B4. To determine the orientation of AP and its metabolites within the active site, we restrained the molecules in the active site of P450 2B4 using the distances calculated from the R_P values. Simulated annealing of the complex was used to identify binding site(s) of these molecules.

The MD snapshots of the last 50 ps at 300 K are shown in Figure 7 for AP, DMAP, and AAP. In all cases, they occupy similar niches within the P450 2B4 active site, which correlates well with the similarities of the ligands' STD NMR spectra. The phenyl group fits into a crevasse formed by seven hydrophobic residues: V104, F108, I114, F115, L238, F297, and V367. The phenyl groups of chlorophenyl imidazole inhibitors in the X-ray crystal structures of P450 2B4 and 2B6 interact with similar residues but lie perpendicular to the orientation of the phenyl group in the AP and its metabolites.^{12,59,77} Mutating V367 causes significant changes in the binding affinity and metabolic profiles of testosterone and BP.^{22,78} The methyl at position 1 on the pyrazolone ring binds in a hydrophobic pocket formed by I101, V104, I209, and V477. This is the same pocket occupied by chlorines of the chlorophenyl imidazole inhibitor-bound X-ray crystal structures.^{12,59,77^t} The methyl at position 5 interacts with a wider hydrophobic pocket formed by residues F206, I209, V477, and G478. The V477F mutant has strong effects on the regioselectivity of the enzyme toward testosterone.⁷⁸ The oxygen at position 3 of the pyrazolone does not directly interact with any residues but forms a hydrogen bond with water in the active site for all three ligands.

The most significant differences in the interactions of these molecules with P450 2B4 were at the site of metabolism. In the AP-bound model (Figure 7A), the dimethylamido group at position 4 on the pyrazolone ring rotates in the MD simulation. During the rotation, one of the methyls interacts with three residues (i.e., A298, I363, and the methyl of T302) that are near



Figure 7. Distance-restrained model of (A) AP, (B) DMAP, and (C) AAP to P450 2B4. The left column shows the MD snapshots of these molecules. To differentiate the protein residues from the ligand, the carbons are colored green and orange, respectively. The heme is colored red. For the sake of clarity, only one of the MD snapshots of P450 2B4 is shown and residues I101, V477, and G478 are hidden from view. In the right column are cartoon representations of the figures that are shown on the left. The heme is labeled and shown as a circle and a line. Hydrogen bonds are represented by dotted lines.

the heme. The I363 residue is the least conserved of these residues. The I363A mutation, which is found in rat P450 2B2,⁷⁹ increased the rate of metabolism of the enzyme toward testo-sterone and altered its stereo- and regioselectivity toward the ligand.⁷⁸ On the other hand, mutating T302 to an alanine inhibited dealkylation of BP and 7-ethoxy-4-trifluoromethyl coumarin.^{22,78} The other methyl interacts with F206 and the hydrophobic section of the side chain of E301 with the carboxylic acid functional group facing away from molecule. These methyls form a diagonal with respect to the heme plane, where the amidomethyl bond is positioned within ~7 Å.

The methyl of the desmethylamido group of DMAP also rotates freely in the MD simulation (Figure 7B). Like AP, the methyl interacted with A298, T302, and I363. In this orientation, the amide of DMAP formed hydrogen bonds with the carbonyl backbone of F297 and the hydroxyl of T302. The 4-CPI molecule in the P450 2B6 X-ray crystal structure has similar interactions with residues within the P450 active site.¹² The imidazole functional group of 4-CPI in the X-ray crystal structure of P450 2B6 is hydrogen bonded to the carbonyl backbone of A297 and the hydroxyl of T302.¹²

The amide functional group of AAP remains relatively fixed during the MD simulation (Figure 7C). The group forms a hydrogen bond with the oxygen on the pyrazolone functional group. AAP either interacted weakly with the heme or formed hydrogen bonds with a heme-bound water, which was consistent with the ligand-induced shift of the P450 2B4 heme to LS (see Figures 1E and 2C).

Mechanism of AP Metabolism. On the basis of the distancerestrained models presented in Figure 7, a mechanism for AP metabolism is proposed in Scheme 2. AP and its metabolites occupy very similar hydrophobic niches within the P450 2B4 active site. The N-linked methyls of the dimethylamido group rotate freely in AP in Scheme 2A between noncatalytic and catalytic positions. The methyl in the noncatalytic position interacts with F206 and E301, while the other methyl binds to a hydrophobic pocket formed by A298, I363, and T302 in a catalytic position, where it is demethylated to DMAP. Because DMAP binds weakly, the molecule can migrate out of the active site. If the molecule remains bound or rebinds, DMAP can assume a catalytically inactive orientation with the N-linked methyl of DMAP in Scheme 2B interacting with F206 and E301. Rather than unbinding and rebinding into a catalytically active orientation, which is energetically unfavored and inefficient, the methyl rotates, until it binds to the hydrophobic pocket near the heme in Scheme 2C. To stabilize the molecule, the Scheme 2. Cartoon Model of AP Metabolism^a



^{*a*} The diagram shows (A) AP, (B) DMAP in a catalytically inactive position, (C) DMAP in a catalytically active position, and (D) the product AAP. Thick lines represent hydrophobic crevasses and pockets of P450 2B4, while dashed lines represent hydrogen bonds between residues of P450 2B4 and DMAP. The catalytic and noncatalytic positions of the methyl are labeled in panel A. The heme is labeled and is shown as a circle and a line.

amide of DMAP forms hydrogen bonds with the backbone of F297 and a hydroxyl of T302, and the molecule is demethylated again to form AAP in Scheme 2D. The amide of AAP is stabilized by a hydrogen bond to the oxygen of the pyrazolone ring. Both the hydrogen and the oxygen interact weakly with either the heme or an intervening heme-bound water. AP or the intermediate DMAP exchanges with AAP, and the process is repeated. This model may also explain the sequential metabolism of tighter binding ligands such as lidocaine⁶ by allowing the rotation and metabolism of the N-linked alkyl groups without unbinding.

ASSOCIATED CONTENT

Supporting Information. Results of a simulation of r_{avgy} , r_{app} , and R_p of a dynamic phenyl molecule near a heme (Figure S1), Figure S2 that was used to determine fast exchange conditions and shows the difference in NMR peak positions of DMAP in the presence of oxidized and reduced P450 2B4 (details of the simulation and analysis of Figure S2 are in the results of the Supporting Information), additional force field parameters that were used for the MD simulations and the energy minimization to ligate C436 to the heme (Table S1), and the organic synthesis scheme of DMAP (Scheme S1). This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

*The Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, 9500 Gilman Drive, #0703, La Jolla, CA 92093-0703. Telephone: (858) 822-7804. Fax: (858) 246-0089. E-mail: a1roberts@ucsd.edu.

Funding Sources

This study was supported by National Institutes of Health Grant ES003619 and by King Abdulaziz City for Science and Technology (KACST) in Saudi Arabia.

ACKNOWLEDGMENT

We thank Jill L. Hartmann for editing the manuscript. N.F. thanks KACST for a postdoctoral fellowship.

ABBREVIATIONS

AAP, aminoantipyrine; AP, amidopyrine; BP, benzphetamine; P450, cytochrome P450; P450 2B4, cytochrome P450 2B4; P450 2B6, cytochrome P450 2B6; CCD, charge-coupled device; DMAP, desmethylamidopyrine; HIV, human immunodeficiency virus; HS, high-spin; IUPAC, International Union of Pure and Applied Chemistry; KP_i, potassium phosphate; LS, low-spin; NAT2, arylamine *N*acetyltransferase; NMR, nuclear magnetic resonance; NSAID, nonsteroidal anti-inflammatory drug; P420, ferric P420; R_{M} , modeldependent relaxation rate; R_P , paramagnetic relaxation rate; SDBS, Spectral DataBase System for organic compounds; STD, saturation transfer difference; T_1 relaxation, longitudinal T_1 relaxation; TLC, thin-layer chromatography; WATERGATE, WATER suppression by GrAdient Tailored Excitation.

REFERENCES

(1) Zhao, Y., and Halpert, J. R. (2007) Structure-function analysis of cytochromes P450 2B. *Biochim. Biophys. Acta* 1770, 402–412.

(2) Guengerich, F. P., Yun, C. H., and Macdonald, T. L. (1996) Evidence for a 1-electron oxidation mechanism in N-dealkylation of N, N-dialkylanilines by cytochrome P450 2B1. Kinetic hydrogen isotope effects, linear free energy relationships, comparisons with horseradish peroxidase, and studies with oxygen surrogates. J. Biol. Chem. 271, 27321–27329.

(3) Bhakta, M., Hollenberg, P. F., and Wimalasena, K. (2005) Evidence for a hydrogen abstraction mechanism in P450-catalyzed *N*-dealkylations. *Chem. Commun.*, 265–267.

(4) Schwarze, W., Blanck, J., Ristau, O., Janig, G. R., Pommerening, K., Rein, H., and Ruckpaul, K. (1985) Spin state control of cytochrome P-450 reduction and catalytic activity in a reconstituted P-450 LM2 system as induced by a series of benzphetamine analogues. *Chem.-Biol. Interact.* 54, 127–141.

(5) Kobayashi, K., Ishizuka, T., Shimada, N., Yoshimura, Y., Kamijima, K., and Chiba, K. (1999) Sertraline N-demethylation is catalyzed by multiple isoforms of human cytochrome P-450 in vitro. *Drug Metab. Dispos.* 27, 763–766.

(6) Hanna, I. H., Roberts, E. S., and Hollenberg, P. F. (1998) Molecular basis for the differences in lidocaine binding and regioselectivity of oxidation by cytochromes P450 2B1 and 2B2. *Biochemistry* 37, 311–318.

(7) Houston, J. B., and Kenworthy, K. E. (2000) In vitro-in vivo scaling of CYP kinetic data not consistent with the classical Michaelis-Menten model. *Drug Metab. Dispos. 28*, 246–254.

(8) Ito, K., Iwatsubo, T., Kanamitsu, S., Nakajima, Y., and Sugiyama, Y. (1998) Quantitative prediction of in vivo drug clearance and drug interactions from in vitro data on metabolism, together with binding and transport. *Annu. Rev. Pharmacol. Toxicol.* 38, 461–499.

(9) Crewe, H. K., Notley, L. M., Wunsch, R. M., Lennard, M. S., and Gillam, E. M. (2002) Metabolism of tamoxifen by recombinant human cytochrome P450 enzymes: Formation of the 4-hydroxy, 4'-hydroxy and *N*-desmethyl metabolites and isomerization of trans-4-hydroxytamoxifen. *Drug Metab. Dispos.* 30, 869–874.

(10) Macdonald, T. L., Gutheim, W. G., Martin, R. B., and Guengerich, F. P. (1989) Oxidation of substituted *N*,*N*-dimethylanilines by cytochrome P-450: Estimation of the effective oxidation-reduction potential of cytochrome P-450. *Biochemistry 28*, 2071–2077.

(11) Bhakta, M. N., and Wimalasena, K. (2002) Microsomal P450catalyzed N-dealkylation of N,N-dialkylanilines: Evidence for a C_{α} -H abstraction mechanism. J. Am. Chem. Soc. 124, 1844–1845. (12) Gay, S. C., Shah, M. B., Talakad, J. C., Maekawa, K., Roberts, A. G., Wilderman, P. R., Sun, L., Yang, J. Y., Huelga, S. C., Hong, W. X., Zhang, Q., Stout, C. D., and Halpert, J. R. (2010) Crystal structure of a cytochrome P450 2B6 genetic variant in complex with the inhibitor 4-(4-chlorophenyl)imidazole at 2.0-Å resolution. *Mol. Pharmacol.* 77, 529–538.

(13) Agundez, J. A., Martinez, C., and Benitez, J. (1995) Metabolism of aminopyrine and derivatives in man: In vivo study of monomorphic and polymorphic metabolic pathways. *Xenobiotica* 25, 417–427.

(14) Geisslinger, G., Bocker, R., and Levy, M. (1996) High-performance liquid chromatographic analysis of dipyrone metabolites to study their formation in human liver microsomes. *Pharm. Res.* 13, 1272–1275.

(15) Vlahov, V., Badian, M., Verho, M., and Bacracheva, N. (1990) Pharmacokinetics of metamizol metabolites in healthy subjects after a single oral dose of metamizol sodium. *Eur. J. Clin. Pharmacol.* 38, 61–65.

(16) Levy, M., Zylber-Katz, E., and Rosenkranz, B. (1995) Clinical pharmacokinetics of dipyrone and its metabolites. *Clin. Pharmacokinet.* 28, 216–234.

(17) Kanaeva, I. P., Nikityuk, O. V., Davydov, D. R., Dedinskii, I. R., Koen, Y. M., Kuznetsova, G. P., Skotselyas, E. D., Bachmanova, G. I., and Archakov, A. I. (1992) Comparative study of monomeric reconstituted and membrane microsomal monooxygenase systems of the rabbit liver. II. Kinetic parameters of reductase and monooxygenase reactions. *Arch. Biochem. Biophys.* 298, 403–412.

(18) Gram, T. E., Wilson, J. T., and Fouts, J. R. (1968) Some characteristics of hepatic microsomal systems which metabolize aminopyrine in the rat and rabbit. *J. Pharmacol. Exp. Ther.* 159, 172–181.

(19) Bast, A., and Noordhoek, J. (1980) Inhibition of aminopyrine demethylation and binding to cytochrome P-450 by its main metabolites in rat liver microsomes. *Br. J. Pharmacol. 68*, 121P–122P.

(20) Noda, A., Tsubone, N., Mihara, M., Goromaru, T., and Iguchi, S. (1976) Formation of 4-formylaminoantipyrine as a new metabolite of aminopyrine. II. Enzymatic demethylation and oxidation of aminopyrine and 4-monomethylaminoantipyrine. *Chem. Pharm. Bull.* 24, 3229–3231.

(21) Agundez, J. A., Carrillo, J. A., Martinez, C., and Benitez, J. (1995) Aminopyrine metabolism in man: The acetylation of aminoantipyrine cosegregates with acetylation of caffeine. *Ther. Drug Monit. 17*, 1–5.

(22) Muralidhara, B. K., Sun, L., Negi, S., and Halpert, J. R. (2008) Thermodynamic fidelity of the mammalian cytochrome P450 2B4 active site in binding substrates and inhibitors. *J. Mol. Biol.* 377, 232–245.

(23) Gay, S. C., Sun, L., Maekawa, K., Halpert, J. R., and Stout, C. D. (2009) Crystal structures of cytochrome P450 2B4 in complex with the inhibitor 1-biphenyl-4-methyl-1H-imidazole: Ligand-induced structural response through α -helical repositioning. *Biochemistry* 48, 4762–4771.

(24) Scott, E. E., Spatzenegger, M., and Halpert, J. R. (2001) A truncation of 2B subfamily cytochromes P450 yields increased expression levels, increased solubility, and decreased aggregation while retaining function. *Arch. Biochem. Biophys.* 395, 57–68.

(25) Scott, E. E., He, Y. A., Wester, M. R., White, M. A., Chin, C. C., Halpert, J. R., Johnson, E. F., and Stout, C. D. (2003) An open conformation of mammalian cytochrome P450 2B4 at 1.6-Å resolution. *Proc. Natl. Acad. Sci. U.S.A. 100*, 13196–13201.

(26) Omura, T., and Sato, R. (1964) The carbon monoxide-binding pigment of liver microsomes. II. Solubilization, purification, and properties. *J. Biol. Chem.* 239, 2379–2385.

(27) Fernando, H., Halpert, J. R., and Davydov, D. R. (2006) Resolution of multiple substrate binding sites in cytochrome P450 3A4: The stoichiometry of the enzyme-substrate complexes probed by FRET and Job's titration. *Biochemistry* 45, 4199–4209.

(28) Morrison, J. F. (1969) Kinetics of the reversible inhibition of enzyme-catalysed reactions by tight-binding inhibitors. *Biochim. Biophys. Acta* 185, 269–286.

(29) Davydov, D. R., Deprez, E., Hoa, G. H., Knyushko, T. V., Kuznetsova, G. P., Koen, Y. M., and Archakov, A. I. (1995) Highpressure-induced transitions in microsomal cytochrome P450 2B4 in solution: Evidence for conformational inhomogeneity in the oligomers. *Arch. Biochem. Biophys.* 320, 330–344. (30) Renaud, J. P., Davydov, D. R., Heirwegh, K. P., Mansuy, D., and Hui Bon Hoa, G. H. (1996) Thermodynamic studies of substrate binding and spin transitions in human cytochrome P-450 3A4 expressed in yeast microsomes. *Biochem. J.* 319 (Part 3), 675–681.

(31) Regal, K. A., and Nelson, S. D. (2000) Orientation of caffeine within the active site of human cytochrome P450 1A2 based on NMR longitudinal (T_1) relaxation measurements. *Arch. Biochem. Biophys.* 384, 47–58.

(32) Hummel, M. A., Gannett, P. M., Aguilar, J. S., and Tracy, T. S. (2004) Effector-mediated alteration of substrate orientation in cytochrome P450 2C9. *Biochemistry* 43, 7207–7214.

(33) Piotto, M., Saudek, V., and Sklenar, V. (1992) Gradient-tailored excitation for single-quantum NMR spectroscopy of aqueous solutions. *J. Biomol. NMR 2*, 661–665.

(34) Kowalewski, J., and Mäler, L. (2006) Nuclear spin relaxation in liquids: Theory, experiments, and applications, Taylor & Francis, New York.

(35) Mayer, M., and Meyer, B. (2001) Group epitope mapping by saturation transfer difference NMR to identify segments of a ligand in direct contact with a protein receptor. *J. Am. Chem. Soc.* 123, 6108–6117.

(36) Meyer, B., and Peters, T. (2003) NMR spectroscopy techniques for screening and identifying ligand binding to protein receptors. *Angew. Chem.* 42, 864–890.

(37) Parella, T., Adell, P., Sánchez-Ferrando, F., and Virgili, A. (1998) Effective multiple-solvent suppression scheme using the excitation sculpting principle. *Magn. Reson. Chem.* 36, 245–249.

(38) Aires-de-Sousa, J., Hemmer, M. C., and Gasteiger, J. (2002) Prediction of ¹H NMR chemical shifts using neural networks. *Anal. Chem.* 74, 80–90.

(39) Bruce, S. D., Higinbotham, J., Marshall, I., and Beswick, P. H. (2000) An analytical derivation of a popular approximation of the Voigt function for quantification of NMR spectra. *J. Magn. Reson.* 142, 57–63.

(40) Kielkopf, J. F. (1973) New approximation to the Voigt function with applications to spectral-line profile analysis. *J. Opt. Soc. Am.* 63, 987–995.

(41) Marshall, I., Bruce, S. D., Higinbotham, J., MacLullich, A., Wardlaw, J. M., Ferguson, K. J., and Seckl, J. (2000) Choice of spectroscopic lineshape model affects metabolite peak areas and area ratios. *Magn. Reson. Med.* 44, 646–649.

(42) Marshall, I., Higinbotham, J., Bruce, S., and Freise, A. (1997) Use of Voigt lineshape for quantification of in vivo ¹H spectra. *Magn. Reson. Med.* 37, 651–657.

(43) Stoica, P., Selen, Y., Sandgren, N., and Van Huffel, S. (2004) Using prior knowledge in SVD-based parameter estimation for magnetic resonance spectroscopy: The ATP example. *IEEE Trans. Biomed. Eng. 51*, 1568–1578.

(44) Chen, H., Van Huffel, S., van Ormondt, D., and de Beer, R. (1996) Parameter estimation with prior knowledge of known signal poles for the quantification of NMR spectroscopy data in the time domain. J. Magn. Reson., Ser. A 119, 225–234.

(45) Viereck, N., Nørgaard, L., Bro, R., and Engelsen, S. B. (2006) Chemometric Analysis of NMR Data. In *Modern Magnetic Resonance* (Webb, G. A., Ed.) pp 1833–1843, Springer, Dordrecht, The Netherlands.

(46) Gay, S. C., Roberts, A. G., Maekawa, K., Talakad, J. C., Hong, W. X., Zhang, Q., Stout, C. D., and Halpert, J. R. (2010) Structures of cytochrome P450 2B4 complexed with the antiplatelet drugs ticlopidine and clopidogrel. *Biochemistry* 49, 8709–8720.

(47) Villafranca, J. J. (1989) Paramagnetic probes of macromolecules. *Methods Enzymol.* 177, 403–413.

(48) Solomon, I. (1955) Relaxation processes in a system of two spins. *Phys. Rev.* 99, 559–565.

(49) Modi, S., Primrose, W. U., Boyle, J. M., Gibson, C. F., Lian, L. Y., and Roberts, G. C. (1995) NMR studies of substrate binding to cytochrome P450 BM3: Comparisons to cytochrome P450cam. *Biochemistry* 34, 8982–8988.

(50) Dwek, R. A. (1973) Nuclear Magnetic Resonance (N.M.R.) in Biochemistry: Applications to Enzyme Systems, Oxford University Press, New York.

(51) Cameron, M. D., Wen, B., Roberts, A. G., Atkins, W. M., Campbell, A. P., and Nelson, S. D. (2007) Cooperative binding of acetaminophen and caffeine within the P450 3A4 active site. *Chem. Res. Toxicol.* 20, 1434–1441.

(52) Tsalkova, T. N., Davydova, N. Y., Halpert, J. R., and Davydov, D. R. (2007) Mechanism of interactions of α -naphthoflavone with cytochrome P450 3A4 explored with an engineered enzyme bearing a fluorescent probe. *Biochemistry* 46, 106–119.

(53) Davydov, D. R., Fernando, H., and Halpert, J. R. (2006) Variable path length and counter-flow continuous variation methods for the study of the formation of high-affinity complexes by absorbance spectroscopy. An application to the studies of substrate binding in cytochrome P450. *Biophys. Chem.* 123, 95–101.

(54) Hummel, M. A., Gannett, P. M., Aguilar, J., and Tracy, T. S. (2008) Substrate proton to heme distances in CYP2C9 allelic variants and alterations by the heterotropic activator, dapsone. *Arch. Biochem. Biophys.* 475, 175–183.

(55) Clore, G. M., and Iwahara, J. (2009) Theory, practice, and applications of paramagnetic relaxation enhancement for the characterization of transient low-population states of biological macromolecules and their complexes. *Chem. Rev.* 109, 4108–4139.

(56) Iwahara, J., Schwieters, C. D., and Clore, G. M. (2004) Ensemble approach for NMR structure refinement against ¹H paramagnetic relaxation enhancement data arising from a flexible paramagnetic group attached to a macromolecule. *J. Am. Chem. Soc.* 126, 5879–5896.

(57) Brunger, A. T., Adams, P. D., and Rice, L. M. (1997) New applications of simulated annealing in X-ray crystallography and solution NMR. *Structure* 5, 325–336.

(58) Clore, G. M., and Schwieters, C. D. (2002) Theoretical and computational advances in biomolecular NMR spectroscopy. *Curr. Opin. Struct. Biol.* 12, 146–153.

(59) Scott, E. E., White, M. A., He, Y. A., Johnson, E. F., Stout, C. D., and Halpert, J. R. (2004) Structure of mammalian cytochrome P450 2B4 complexed with 4-(4-chlorophenyl)imidazole at 1.9 Å resolution: Insight into the range of P450 conformations and coordination of redox partner binding. *J. Biol. Chem.* 279, 27294–27301.

(60) Knight, J. L., Mekler, V., Mukhopadhyay, J., Ebright, R. H., and Levy, R. M. (2005) Distance-restrained docking of rifampicin and rifamycin SV to RNA polymerase using systematic FRET measurements: Developing benchmarks of model quality and reliability. *Biophys. J.* 88, 925–938.

(61) Yue, S. Y. (1990) Distance-constrained molecular docking by simulated annealing. *Protein Eng. 4*, 177–184.

(62) Constantine, K. L. (2001) Evaluation of site-directed spin labeling for characterizing protein-ligand complexes using simulated restraints. *Biophys. J.* 81, 1275–1284.

(63) Eswar, N., Eramian, D., Webb, B., Shen, M. Y., and Sali, A. (2008) Protein structure modeling with MODELLER. *Methods Mol. Biol.* 426, 145–159.

(64) Van Der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A. E., and Berendsen, H. J. (2005) GROMACS: Fast, flexible, and free. *J. Comput. Chem.* 26, 1701–1718.

(65) Oostenbrink, C., Soares, T. A., van der Vegt, N. F., and van Gunsteren, W. F. (2005) Validation of the 53A6 GROMOS force field. *Eur. Biophys. J.* 34, 273–284.

(66) Gunsteren, W. F. v., Billeter, S. R., Eising, A. A., Hunenberger, P. H., Kruger, P., Mark, A. E., Scott, W. R. P., and Tironi, I. G. (1996) *Biomolecular Simulation: The GROMOS96 Manual and User Guide*, Verlag der Fachvereine Hochschulverlag AG an der ETH Zurich, Zurich.

(67) Darden, T., York, D., and Pedersen, L. (1993) Particle mesh Ewald: An N-log(N) method for Ewald sums in large systems. *J. Chem. Phys.* 98, 10089–10092.

(68) Jefcoate, C. R. (1978) Measurement of substrate and inhibitor binding to microsomal cytochrome P-450 by optical-difference spectroscopy. *Methods Enzymol.* 52, 258–279.

(69) Miller, G. P., and Guengerich, F. P. (2001) Binding and oxidation of alkyl 4-nitrophenyl ethers by rabbit cytochrome P450 1A2: Evidence for two binding sites. *Biochemistry* 40, 7262–7272.

(70) Koymans, L., Donne-op den Kelder, G. M., Koppele Te, J. M., and Vermeulen, N. P. (1993) Cytochromes P450: Their active-site structure and mechanism of oxidation. *Drug Metab. Rev.* 25, 325–387.

(71) Locuson, C. W., Hutzler, J. M., and Tracy, T. S. (2007) Visible spectra of type II cytochrome P450-drug complexes: Evidence that "incomplete" heme coordination is common. *Drug Metab. Dispos.* 35, 614–622.

(72) Segall, M. D., Payne, M. C., Ellis, W., Tucker, G. T., and Boyes, N. (1998) Evidence for stabilization of the low-spin state of cytochrome P450 due to shortening of the proximal heme bond. *Chem. Res. Toxicol. 11*, 962–966.

(73) Bridges, A., Gruenke, L., Chang, Y. T., Vakser, I. A., Loew, G., and Waskell, L. (1998) Identification of the binding site on cytochrome P450 2B4 for cytochrome b_5 and cytochrome P450 reductase. *J. Biol. Chem.* 273, 17036–17049.

(74) Gharbi-Benarous, J., Bertho, G., Evrard-Todeschi, N., Coadou, G., Megy, S., Delaunay, T., Benarous, R., and Girault, J. P. (2004) Epitope mapping of the phosphorylation motif of the HIV-1 protein Vpu bound to the selective monoclonal antibody using TRNOESY and STD NMR spectroscopy. *Biochemistry* 43, 14555–14565.

(75) Claasen, B., Axmann, M., Meinecke, R., and Meyer, B. (2005) Direct observation of ligand binding to membrane proteins in living cells by a saturation transfer double difference (STDD) NMR spectroscopy method shows a significantly higher affinity of integrin $\alpha_{IID}\beta_3$ in native platelets than in liposomes. *J. Am. Chem. Soc.* 127, 916–919.

(76) Ammann, L. P., Merritt, M., Sagalowsky, A., and Nurenberg, P. (2006) Peak-finding partial least squares for the analysis of ¹H NMR spectra. *Chemom. Intell. Lab. Syst.* 20, 231–238.

(77) Zhao, Y., Sun, L., Muralidhara, B. K., Kumar, S., White, M. A., Stout, C. D., and Halpert, J. R. (2007) Structural and thermodynamic consequences of 1-(4-chlorophenyl)imidazole binding to cytochrome P450 2B4. *Biochemistry* 46, 11559–11567.

(78) Hernandez, C. E., Kumar, S., Liu, H., and Halpert, J. R. (2006) Investigation of the role of cytochrome P450 2B4 active site residues in substrate metabolism based on crystal structures of the ligand-bound enzyme. *Arch. Biochem. Biophys.* 455, 61–67.

(79) Mizukami, Y., Sogawa, K., Suwa, Y., Muramatsu, M., and Fujii-Kuriyama, Y. (1983) Gene structure of a phenobarbital-inducible cytochrome P-450 in rat liver. *Proc. Natl. Acad. Sci. U.S.A.* 80, 3958–3962.