

Communication

# Nickel-catalysed three-component connection reaction of a phenyl group, conjugated dienes, and aldehydes: stereoselective synthesis of (*E*)-5-phenyl-3-penten-1-ols and (*E*)-3-methyl-5-phenyl-3-penten-1-ols

Kazufumi Shibata, Masanari Kimura, Keisuke Kojima, Shuji Tanaka, Yoshinao Tamaru \*

Department of Applied Chemistry, Faculty of Engineering, Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan

Received 25 September 2000; accepted 10 November 2000

This manuscript is dedicated to Professor Jean Normant on the occasion of his 65th birthday for his outstanding contribution to the development of organometallic chemistry

## Abstract


In the presence of 10 mol% of  $\text{Ni}(\text{acac})_2$ ,  $\text{Ph}_2\text{Zn}$  reacts with 1,3-butadiene and aldehydes at room temperature to give 1-alkyl and 1-aryl substituted (*E*)-5-phenyl-3-penten-1-ols (**3**) in good yields. Under similar conditions, the three components of  $\text{Ph}_3\text{BZnEt}_2$ , isoprene, and aldehydes combine with each other to furnish 1-alkyl and 1-aryl substituted (*E*)-3-methyl-5-phenyl-3-penten-1-ols (**5**) in good yields. © 2001 Elsevier Science B.V. All rights reserved.

**Keywords:** Dienes; Nickel; Organoboranes; Organozincs; Multi-component connection

1,3-Dienes have long been utilised as one of the most versatile building blocks in organic synthesis, especially in the field of transition-metal catalysed reactions for the synthesis of complex molecules [1]. Since the isoprene C-5 unit is found abundantly in natural products, the 1,4-difunctionalisation reaction of isoprene with different kinds of reactants is particularly useful, provided the stereochemistry of the tri-substituted double bond (*E* or *Z*) as well as the regiochemistry (head or tail) of the products can be regulated appropriately.



Recently, we have developed novel nickel-catalysed three-component connection reactions of organometallic reagents, 1,3-dienes, and carbonyl compounds that furnish a variety of 4-penten-1-ols (via homoallylation of carbonyls with dienes) [2,3] and 3-penten-1-ols (via allylation of carbonyls with dienes) [4,5] with high regio- and stereoselectivity.

In the latter nickel-catalysed allylative three-component connection reactions, 1,3-butadiene underwent facile reaction with  $\text{Me}_2\text{Zn}$  and aldehydes at room temperature to furnish (*E*)-3-hexen-1-ols (**2**) in excellent yields (Eq. (1)). Isoprene also reacted regioselectively with  $\text{Me}_2\text{Zn}$  at the C-4-position and with aldehydes at the C-1 position to give 3-methyl-3-hexen-1-ols (**4**) in good yields (Eq. (2)) [4]. The reaction, however, displayed poor stereoselectivity, giving rise to mixtures of (*E*)- and (*Z*)-**4** in the ratios of ca. 3:1. The stereoselectivity was improved greatly by the use of  $\text{Me}_3\text{B}$  in place of  $\text{Me}_2\text{Zn}$ , where the (*E*)-**4** isomers were produced either exclusively (for alkyl aldehydes) or highly selectively (> 13:1, for aromatic aldehydes) (Eq. (3)) [5].



\* Corresponding author. Tel./fax: +81-95-8479008.

E-mail address: tamaru@net.nagasaki-u.ac.jp (Y. Tamaru).



Here we would like to report that the three-component connection reactions of Ph–1,3-butadiene–aldehydes and Ph–isoprene–aldehydes proceed nicely and furnish (*E*)-5-phenyl-3-penten-1-ols (**3**) and (*E*)-3-methyl-5-phenyl-3-penten-1-ols (**5**), respectively, in high yields and with excellent stereoselectivity under the catalysis of a nickel complex.

We expected that the three-component connection reaction of Ph<sub>2</sub>Zn–dienes–aldehydes would be rather problematic, since, as compared with Me<sub>2</sub>Zn, Ph<sub>2</sub>Zn is so reactive that it might possibly react with aldehydes without incorporating dienes as the reaction partner [6]. Accordingly, we undertook preliminary experiments for the reaction of Ph<sub>2</sub>Zn with benzaldehyde in the absence and in the presence of a nickel catalyst (Table 1). In fact, Ph<sub>2</sub>Zn reacted rather slowly with benzaldehyde at room temperature (runs 1 and 2). However, in the presence of 10 mol% of Ni(acac)<sub>2</sub> [acac = acetylacetonate], it underwent facile reaction. The reaction was complete within 20 min at room temperature and provided benzhydrol in a quantitative yield (run 3).

Despite these somewhat discouraging preliminary results, to our pleasant surprise, the three-component connection reaction did proceed nicely for the reaction with 1,3-butadiene and aldehydes (runs 1–4, Table 2). Except for the reaction with dihydrocinnamaldehyde (run 2, Table 2), the direct phenylation of aldehydes presented little difficulty and the expected products **3** were obtained in reasonable yields and with excellent

*E*-selectivity. No (*Z*)-**3** isomers were detected. Furthermore, 1-substituted 9-phenylnona-3,7-dien-1-ols, the type of compounds obtained as the major side products in the reaction of Me<sub>2</sub>Zn–1,3-butadiene–aldehydes [4], were not formed.


For the reaction with isoprene, on the other hand, the phenylation of aldehydes turned out to be a very serious problem, especially for the reactions with reactive aldehydes (benzaldehyde and alkyl aldehydes that do not possess substituents at the α-position) (runs 5 and 6, Table 2). In these cases, the expected products **5a** and **5b** were produced only as the minor products. An attempt to minimise **6a** by adding slowly the Ph<sub>2</sub>Zn solution into the solution of benzaldehyde, isoprene, and Ni(acac)<sub>2</sub> over 1 h was unsuccessful, and **5a** and **6a** were obtained in 38 and 39% yields, respectively. For the reaction with less reactive secondary and tertiary alkyl aldehydes, the expected products **5c** and **5d** were obtained in moderate yields (runs 7 and 8).

In order to circumvent the competitive phenylation of aldehydes, we next examined Ph<sub>3</sub>B as the phenylating agent, since our preliminary study indicated that Ph<sub>3</sub>B was unreactive toward benzaldehyde even in the presence of nickel catalysts (runs 4–6, Table 1).

Although Ni(acac)<sub>2</sub> served nicely as the catalyst for the connection reaction of Me<sub>3</sub>B–isoprene–aldehydes (Eq. (3)) [5], it turned out to be completely ineffective for the version of Ph<sub>3</sub>B–isoprene–aldehydes and no **5** formed at all. In sharp contrast to this, Ni(cod)<sub>2</sub> [cod = cyclooctadiene] promoted the reaction effectively (Table 3). In the presence of 10 mol% of Ni(cod)<sub>2</sub>, the reaction proceeded smoothly at room temperature and furnished

Table 1

Reaction of Ph<sub>2</sub>Zn and Ph<sub>3</sub>B with benzaldehyde in the presence and in the absence of nickel catalysts <sup>a</sup>



| Run | Phenylating agent  | Nickel catalyst       | Reaction time (h) | Benzhydrol (% yield) <sup>b</sup> |
|-----|--------------------|-----------------------|-------------------|-----------------------------------|
| 1   | Ph <sub>2</sub> Zn | None                  | 5                 | 81                                |
| 2   | Ph <sub>2</sub> Zn | None                  | 0.3               | 65                                |
| 3   | Ph <sub>2</sub> Zn | Ni(acac) <sub>2</sub> | 0.3               | 96                                |
| 4   | Ph <sub>3</sub> B  | None                  | 36                | 0 <sup>c</sup>                    |
| 5   | Ph <sub>3</sub> B  | Ni(acac) <sub>2</sub> | 10                | 0 <sup>c</sup>                    |
| 6   | Ph <sub>3</sub> B  | Ni(cod) <sub>2</sub>  | 10                | 0 <sup>c</sup>                    |

<sup>a</sup> A mixture of benzaldehyde (1 mmol), Ph<sub>2</sub>Zn (1.2 mmol), and a nickel catalyst (0.1 mmol, if indicated) in dry THF (5 ml) was stirred at room temperature under N<sub>2</sub>.

<sup>b</sup> Yields refer to the isolated yields of benzhydrol by means of column chromatography over silica gel.

<sup>c</sup> No formation of benzhydrol was confirmed by means of TLC and GLPC.

Table 2

Synthesis of 5-phenyl-3-penten-1-ols (**3**) and 3-methyl-5-phenyl-3-penten-1-ols (**5**) via the three component reaction of  $\text{Ph}_2\text{Zn}$ , 1,3-dienes, and aldehydes in the presence of  $\text{Ni}(\text{acac})_2^a$

| Run | Aldehyde | Diene | Reaction time (h) | Product <b>5</b> (% yield) <sup>b</sup> [ <i>E:Z</i> ] <sup>c</sup> | Product <b>6</b> (% yield) <sup>b</sup> |
|-----|----------|-------|-------------------|---------------------------------------------------------------------|-----------------------------------------|
| 1   |          |       | 0.3               | <br><b>3a</b> (61) [only <i>E</i> ]                                 | <br><b>6a</b> (23)                      |
| 2   |          |       | 0.3               | <br><b>3b</b> (34) [only <i>E</i> ]                                 | <br><b>6b</b> (39)                      |
| 3   |          |       | 0.3               | <br><b>3c</b> (68) [only <i>E</i> ]                                 | <br><b>6c</b> (7)                       |
| 4   |          |       | 3                 | <br><b>3d</b> (70) [only <i>E</i> ]                                 | <br><b>6d</b> (0) <sup>d</sup>          |
| 5   |          |       | 0.3               | <br><b>5a</b> (27) [only <i>E</i> ]                                 | <b>6a</b> (59)                          |
| 6   |          |       | 0.3               | <br><b>5b</b> (16) [only <i>E</i> ]                                 | <b>6b</b> (57)                          |
| 7   |          |       | 0.3               | <br><b>5c</b> (46) [7:1]                                            | <b>6c</b> (21)                          |
| 8   |          |       | 3                 | <br><b>5d</b> (46) [20:1]                                           | <b>6d</b> (0) <sup>d</sup>              |

<sup>a</sup> A mixture of an aldehyde (1 mmol), 1,3-butadiene (4 mmol) or isoprene (4 mmol),  $\text{Ph}_2\text{Zn}$  [1.2 mmol, prepared from  $\text{PhMgBr}$  (2.4 mmol) and  $\text{ZnCl}_2$  (1.2 mmol)], and  $\text{Ni}(\text{acac})_2$  (0.1 mmol) in dry THF was stirred at ambient temperature under  $\text{N}_2$ . <sup>b</sup> Yields refer to the isolated spectroscopically homogeneous materials. <sup>c</sup> The *E,Z*-ratios were determined by means of  $^1\text{H}$  NMR (400 MHz) and  $^{13}\text{C}$  NMR (100 MHz). <sup>d</sup> The alcohol could not be detected.

**5** in reasonable yields, irrespective of the kinds of aldehydes. The stereoselectivity, however, was not satisfactory: In most cases, (*E*)- and (*Z*)-**5** were produced in the ratios of ca. 10:1.

The ratios of (*E*)- and (*Z*)-**5** could be determined with high accuracy on the basis of the base-line separated resonances of one of the diastereotopic C-2 methylene protons in their  $^1\text{H}$ -NMR spectra (400 MHz,  $\text{CDCl}_3$ ) [7]. The stereochemistry of (*E*)- and (*Z*)-**5** was determined unequivocally on the basis of their  $^{13}\text{C}$ -NMR spectra (100 MHz) [7].

The product was accompanied with **6** in substantial quantities (Table 3). In the light of the results shown in runs 4–6 (Table 1), the origin of **6** is not clear at present. However, this is, to the best of our knowledge, the first example that indicates that  $\text{Ph}_3\text{B}$  is capable of

undergoing the Grignard-like addition to aldehydes [8–10] under certain conditions in the presence of a nickel complex.


Encouraged by the favourable reaction features of  $\text{Ph}_3\text{B}$  as the phenylating agent, we next examined the reaction with the ate complexes of  $\text{Ph}_3\text{B}$ . The organoborate complexes were generated in situ by the reaction of  $\text{Ph}_3\text{B}$  with an equal amount of an organometallic reagent.









The results obtained for the reaction with benzaldehyde and a variety of organoborates are summarised in runs 1–6 in Table 4. As is apparent from these results, the reactivity markedly depended on the kinds of the organic substituents as well as the metal counter ions of these borate complexes. The most striking feature regarding the dependence of reactivity on the metal coun-

The reaction was performed typically as follows (run 6, Table 4): Into a solution of  $\text{Ph}_3\text{B}$  (2.4 mmol, 0.25 M

In conclusion, we have developed the nickel-catalysed three-component connection reaction of a phenyl group, 1,3-dienes, and aldehydes. For the reaction with 1,3-butadiene,  $\text{Ph}_2\text{Zn}$  serves as an efficient phenylating reagent and 1-alkyl and 1-aryl substituted 5-phenyl-3-penten-1-ols (**3**) are obtained in good yields. For the reaction with isoprene, 1-alkyl and 1-aryl substituted 3-methyl-5-phenyl-3-penten-1-ols (**5**) are obtained in


Table 3  
Synthesis of 3-methyl-5-phenyl-3-penten-1-ols (**5**) via the three component reaction of  $\text{Ph}_3\text{B}$ , isoprene, and aldehydes in the presence of  $\text{Ni}(\text{cod})_2^a$










| Run | Aldehyde                                                                            | Reaction time (h) | Product <b>5</b> (% yield) <sup>b</sup> [ <i>E:Z</i> ] <sup>c</sup>                                                             | Product <b>6</b> (% yield) <sup>b</sup>                                                                                    |
|-----|-------------------------------------------------------------------------------------|-------------------|---------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|
| 1   |  | 10                | <br><b>5a</b> ( <b>62</b> ) [10:1]           | <br><b>6a</b> ( <b>8</b> )              |
| 2   |  | 27                | <br><b>5b</b> ( <b>38</b> ) [only <i>E</i> ] | <br><b>6b</b> ( <b>22</b> )             |
| 3   |  | 24                | <br><b>5c</b> ( <b>63</b> ) [7.3:1]          | <br><b>6c</b> ( <b>20</b> )             |
| 4   |  | 28                | <br><b>5d</b> ( <b>71</b> ) [10:1]           | <br><b>6d</b> ( <b>0</b> ) <sup>d</sup> |

<sup>a</sup> A mixture of an aldehyde (1 mmol), isoprene (4 mmol), Ph<sub>3</sub>B (2.4 mmol), and Ni(cod)<sub>2</sub> (0.1 mmol) in dry THF was stirred at ambient temperature under N<sub>2</sub>. <sup>b</sup> Yields refer to the isolated spectroscopically homonegeous materials. <sup>c</sup> The *E,Z*-ratios were determined by means of <sup>1</sup>H NMR (400 MHz) and <sup>13</sup>C NMR (100 MHz). <sup>d</sup> The alcohol could not be detected.

Synthesis of 3-methyl-5-phenyl-3-penten-1-ols (**5**) via the three component reaction of  $\text{Ph}_3\text{B}$ , isoprene, and aldehydes in the presence of  $\text{Ni}(\text{acac})_2$  and organometallic reagents (RM)<sup>a</sup>



| Run | Aldehyde                                                                           | RM                               | Time (h)/Temp (°C) | Product <b>5</b> (% yield) <sup>b</sup> [ <i>E</i> : <i>Z</i> ] <sup>c</sup>                                         | Product <b>6</b> (% yield) <sup>b</sup>                                                                          |
|-----|------------------------------------------------------------------------------------|----------------------------------|--------------------|----------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|
| 1   |   | MeLi                             | 24/25              | <br><b>5a</b> (86) [only <i>E</i> ] | <br><b>6a</b> (8)               |
| 2   |                                                                                    | MeMgBr                           | 20/25 then 12/50   | <b>5a</b> (91) [10:1]                                                                                                | <b>6a</b> (0)                                                                                                    |
| 3   |                                                                                    | PhMgBr                           | 24/25 then 10/50   | <b>5a</b> (49) [10:1]                                                                                                | <b>6a</b> (0)                                                                                                    |
| 4   |                                                                                    | Ph <sub>4</sub> BNa <sup>d</sup> | 23/35              | no reaction                                                                                                          |                                                                                                                  |
| 5   |                                                                                    | Me <sub>2</sub> Zn               | 15/25              | <b>5a</b> (80) [only <i>E</i> ]                                                                                      | <b>6a</b> (17)                                                                                                   |
| 6   |                                                                                    | Et <sub>2</sub> Zn               | 1.5/25             | <b>5a</b> (80) [only <i>E</i> ]                                                                                      | <b>6a</b> (0)                                                                                                    |
| 7   |   | Et <sub>2</sub> Zn               | 2/25               | <br><b>5b</b> (60) [only <i>E</i> ] | <br><b>6b</b> (20)              |
| 8   |   | Et <sub>2</sub> Zn               | 2/25               | <br><b>5c</b> (60) [only <i>E</i> ] | <br><b>6c</b> (10)              |
| 9   |  | Et <sub>2</sub> Zn               | 10/25              | <br><b>5d</b> (63) [20:1]          | <br><b>6d</b> (0) <sup>e</sup> |

<sup>a</sup> A mixture of an aldehyde (1 mmol), isoprene (4 mmol), Ph<sub>3</sub>B (2.4 mmol), RM (2.4 mmol), and Ni(acac)<sub>2</sub> (0.1 mmol) in dry THF was stirred under N<sub>2</sub> at the temperature indicated. <sup>b</sup> Yields refer to the isolated spectroscopically homogeneous materials. <sup>c</sup> The *E,Z*-ratios were determined by means of <sup>1</sup>H NMR (400 MHz) and <sup>13</sup>C NMR (100 MHz). <sup>d</sup> Ph<sub>4</sub>B Na (2.4 mmol) was used as received (Aldrich). <sup>e</sup> The alcohol could not be detected.

good yields when isoprene is subjected to the reaction with  $\text{Ph}_3\text{BZnEt}_2$  and aldehydes at room temperature in the presence of 10 mol% of  $\text{Ni}(\text{acac})_2$ . In both reactions, the (*E*)-isomers, (*E*)-**3** and (*E*)-**5**, are obtained with excellent stereoselectivity.

## Acknowledgements

The authors thank the Ministry of Education, Science, Sports and Culture, Japanese Government for financial support (Grant-in-Aid for Scientific Research B).

## References

- good yields when isoprene is subjected to the reaction with  $\text{Ph}_3\text{BZnEt}_2$  and aldehydes at room temperature in the presence of 10 mol% of  $\text{Ni}(\text{acac})_2$ . In both reactions, the (*E*)-isomers, (*E*)-**3** and (*E*)-**5**, are obtained with excellent stereoselectivity.
- ### Acknowledgements
- The authors thank the Ministry of Education, Science, Sports and Culture, Japanese Government for financial support (Grant-in-Aid for Scientific Research B).
- ### References
- [1] J. Tsuji, *Transition Metal Reagents and Catalysts*, Wiley, Chichester, 2000 Chapter 5.
  - [2] M. Kimura, A. Ezoe, K. Shibata, Y. Tamaru, *J. Am. Chem. Soc.* 120 (1998) 4033.
  - [3] M. Kimura, H. Fujimatsu, A. Ezoe, K. Shibata, M. Shimizu, S. Matsumoto, Y. Tamaru, *Angew. Chem. Int. Ed. Engl.* 38 (1999) 397.
  - [4] M. Kimura, S. Matsuo, K. Shibata, Y. Tamaru, *Angew. Chem. Int. Ed. Engl.* 38 (1999) 3386.
  - [5] M. Kimura, K. Shibata, Y. Koudahashi, Y. Tamaru, *Tetrahedron Lett.* 41 (2000) 6789.
  - [6] P. Knochel, P. Jones, *Organozinc Reagents*, Oxford University, Oxford, 1999.
  - [7] (*E*)-3-Methyl-1,5-diphenyl-3-penten-1-ol ((*E*)-**5a**): IR (neat) 3400 (s), 1495 (s), 1450 (s), 1030 (s)  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.80 (brs, 3 H), 2.07 (d,  $J = 2.4$  Hz, 1 H), 2.41 (dd,  $J = 13.5$ , 9.0 Hz, 1 H), 2.48 (dd,  $J = 13.5$ , 4.7 Hz, 1 H), 3.40 (d,  $J = 7.3$  Hz, 2 H), 4.81 (ddd,  $J = 9.0$ , 4.7, 2.4 Hz, 1 H), 5.50 (brt,  $J = 7.3$  Hz, 1 H), 7.11–7.38 (m, 10 H);  $^{13}\text{C-NMR}$  (100 MHz,  $\text{CDCl}_3$ )  $\delta$  16.3, 34.4, 50.2, 71.8, 125.9, 127.4, 127.7, 128.4, 128.5, 132.8, 141.1, 144.2. Anal. Calc. for  $\text{C}_{18}\text{H}_{20}\text{O}$ : C, 85.67; H, 7.99. Found: C, 85.32; H, 8.18. (*Z*)-3-Methyl-1,5-diphenyl-3-penten-1-ol ((*Z*)-**5a**):  $^1\text{H-NMR}$  (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.81 (brs, 3 H), 1.98 (d,  $J = 2.2$  Hz, 1 H), 2.42 (dd,  $J = 13.5$ , 5.1 Hz, 1 H), 2.76 (dd,

- $J = 13.5$ ,  $9.0$  Hz, 1 H), 3.33 (d,  $J = 7.3$  Hz, 2 H), 4.86 (ddd,  $J = 9.0$ ,  $5.1$ ,  $2.2$  Hz, 1 H), 5.54 (brt,  $J = 7.3$  Hz, 1 H), 7.20–7.41 (m, 10 H);  $^{13}\text{C}$ -NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  23.8, 34.2, 42.4, 72.4, 125.8, 125.9, 127.6, 127.9, 128.5, 128.9, 132.3, 141.3, 144.4.
- [8] Alkylation of aldehydes with alkylboranes: (a) B.M. Mikhailov, T.K. Baryshnikova, A.S. Shashkov, J. Organomet. Chem. 219 (1981) 301. (b) G.W. Kabalka, Z. Wu, S.E. Trotman, X. Gao, Org. Lett. 2 (2000) 255.
- [9] Allylation of aldehydes with allylboranes: (a) P.V. Ramachandran, G.-M. Chen, H.C. Brown, Tetrahedron Lett. 38 (1997) 2417. (b) M. Suginome, H. Nakamura, T. Matsuda, Y. Ito, J. Am. Chem. Soc. 120 (1998) 4248. (c) J.D. White, J. Hong, L.A. Robarge, Tetrahedron Lett. 40 (1999) 1463.
- [10] Arylation and vinylation of aldehydes with aryl- and vinylboranes: (a) P. Jacob, III, H.C. Brown, J. Org. Chem. 42 (1977) 579. (b) M. Sakai, M. Ueda, N. Miyaura, Angew. Chem. Int. Ed. Engl. 37 (1998) 3279. (c) R.A. Batey, D.B. MacKay, V. Santhakumar, J. Am. Chem. Soc. 121 (1999) 5075. (d) R.A. Batey, A.N. Thadani, D.M. Smil, Org. Lett. 1 (1999) 1683.