

*Transformation of Swertiamarin to Gentiopicrin.
The Structure of Gentiopicrin*

By Takashi KUBOTA and Yutaka TOMITA

(Received July 17, 1961)

In a preceding communication, it was reported that swertiamarin, the bitter component of *Swertia japonica* Makino, is represented by structure I¹⁾, and that dehydration of the tertiary hydroxyl group of swertiamarin acetate (II) with potassium bisulfate and acetic anhydride affords monoanhydroswertiamarin acetate, C₂₄H₂₈O₁₃ (III), m. p. 143~144°C which shows bands in its ultraviolet absorption spectrum at 246 m μ log ϵ 3.44 and 269 m μ log ϵ 3.37¹⁾.

However, on further purification by chromatography and by recrystallization the melting point of II was raised to 151~152°C (IIIA), and the ultraviolet absorption spectrum showed a band at λ_{max} 245 m μ log ϵ 3.65 and no band at 269 m μ . On the other hand, II treated with BF₃ in AcOH at low temperature for 3 hr. afforded a dehydrated product (IIIB) as colorless plates, m. p. 137~138°C. The ultraviolet absorption spectrum showed a band only at 270 m μ log ϵ 3.8 and none at 246 m μ . Therefore, the monoanhydroswertiamarin acetate (III) described in the previous paper¹⁾ is a mixture of IIIA and IIIB and from the ultraviolet absorption spectra the following formulae are given for IIIA and IIIB respectively. (Fig. 1). The fact that a greater amount of IIIA is produced by the dehydration reaction with potassium bisulfate than the more stable triene

type isomer IIIB, indicates the relationship between the hydroxyl group at C₄ and the hydrogen atom at C₅ to be in cis to each other. In the case of the dehydration reaction with BF₃, double bond migration of IIIA occurs by the action of BF₃ and consequently II gives the stable IIIB.

The infrared absorption spectrum of IIIA shows a band at 12.2 μ which is due to the tri-substituted double bond. Therefore the tertiary hydroxyl group of I must exist at C₄ and not at C₅. From these results, an alternative structure of swertiamarin IA¹⁾ can be excluded.

In order to establish the structure of gentiopicrin a detailed study on its degradation reactions was carried out by Asahina, Asano, Tanase and Ueno²⁾. On the basis of their data the several possible structures were suggested by Sakurai³⁾, Korte³⁾ and Canonica and Pelizzoni⁴⁾, respectively. Although, Canonica and Pelizzoni⁵⁾ have recently proposed the formula IV as the most probable structure for gentiopicrin, the definite evidence for its carbon skeleton is lacking in their paper. If the structure IV is correct, IIIB, a monoanhydro derivative of II should be identical with gentiopicrin acetate.

Now it has been confirmed that IIIB and the acetate, m. p. 139°C, of gentiopicrin, extracted from the root of *Gentiana scabra*, are identical in their infrared absorption spectra.

Concerning the structure of swertiamarin, the correctness of the carbon skeleton of I has been definitely demonstrated by synthesis of erythrocentaurin⁶⁾, the hydrolysis product of swertiamarin with emulsin, and by transformation of swertiamarin to gentianin⁷⁾ as reported in the previous papers. Although several structures were suggested for gentiopicrin, the structure IV proposed by Canonica and Pelizzoni has been established to be correct from the above-mentioned results.

Faculty of Science
Osaka City University
Kita-ku, Osaka




Fig. 1

1) T. Kubota and Y. Tomita, *Tetrahedron Letters*, **5**, 176 (1961).

2) Y. Asahina, J. Asano, Y. Tanase and Y. Ueno, *Ber.*, **69**, 771 (1936); Y. Asahina and Y. Sakurai, *ibid.*, **72**, 1534 (1939).

3) F. Korte, *Chem. Ber.*, **87**, 512, 780 (1954).

4) L. Canonica and F. Pelizzoni, *Gazz. chim. ital.*, **87**, 1251 (1957).

5) L. Canonica, F. Pelizzoni, P. Manitto and G. Jomi, *Tetrahedron Letters*, **24**, 7 (1960).

6) T. Kubota, Y. Tomita and K. Suzuki, *ibid.*, **6**, 223 (1961).

7) T. Kubota and Y. Tomita, *ibid.*, in press.