Absolute Stereochemical Determination of Chiral Carboxylates Using an Achiral Molecular Tweezer

Hongsik Yoon, Chi-Hwa Lee, and Woo-Dong Jang*^[a]

Abstract: A new type of molecular tweezer (1) has been synthesized for the direct determination of the absolute configuration of chiral carboxylates without analyte derivatization. Upon the addition of diamine and anionic guests, 1 exhibited shifts in its absorption spectrum with clear isosbestic points. The continuous variation method indicated that both the diamine and anionic guests form 1:1 host-guest complexes with **1** with very high binding affinity. When Boc-L-Ala (BLA) as a form of tetrabutylammonium salt was added to **1**, a weak negative CD signal was observed. This weak CD signal was dramatically

Keywords: chirality • circular dichroism • host-guest systems • molecular tweezer • porphyrinoids

Introduction

Determination of absolute configuration remains a very important topic in chemistry because many bioactive molecules are only effective if they have a specific stereo-configuration.^[1] To date, various techniques have been developed for the determination of absolute configuration in chiral compounds, including optical rotation, circular dichroism (CD), nuclear magnetic resonance, chiral chromatography, and X-ray crystallography.^[2] Among the various methods, one of the most powerful tools for determining absolute configuration is CD spectroscopy, in which enantiomers give spectroscopic data with differential absorption of either leftor right-circularly polarized light.^[3] To obtain reliable CD signals, molecules should have an appropriate chromophore with a strong absorption. Recently, several porphyrin tweezer systems as probes for absolute stereochemical determination have been developed based on exciton-coupled circular dichroism (ECCD).^[4] The binding of a chiral guest to the porphyrin tweezer induces a chiral twist of the two porphyrin units to minimize steric repulsion. The ECCD method has been successfully applied to a wide variety of compounds, including chiral alcohols, amines, amino alcohols, amino acid derivatives, and bifunctional amide conjugates.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/chem.201200371.

Chem. Eur. J. 2012, 00, 0-0

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

WILEY CONLINE LIBRARY

These are not the final page numbers! **77**

changed to a strong positive CD couplet upon addition of achiral 1,12-diaminododecane. Such a positive CD couplet was observed for all of the tested L-amino acid derivatives, while the D-amino acid derivatives gave the opposite signals. As a result of these unique characteristics of **1**, it can be utilized as a highly sensitive probe for the absolute stereochemical determination of chiral carboxylates.

However, this potentially powerful approach suffers from a severe limitation that precludes its current use in highthroughput screening or other real-time analyses. Specifically, derivatization is required in the case of substrates bearing only a single ligation site.^[5] Herein, we present a solution to this problem that is based on the combination of a porphyrin tweezer system with remote guest-binding receptor modules. In general, the chirality of the guest molecule is directly transferred to the achiral porphyrin tweezer in the form of a twisted conformation through bidentate metal coordination. Owing to the significantly large extinction coefficient of porphyrin units in the visible range, the conformational twist of the porphyrin tweezer system results in large signal amplification.^[6] However, tedious derivatizations of monoamines or monocarboxylic acids tend to decrease their applicability in chiral sensing. Furthermore, a large excess of these guest molecules is often needed to elicit a reliable CD signal due to their weak binding affinities. Because most naturally occurring chiral compounds can only be isolated in very small quantities, stereochemical determination without derivatization as well as probes with high binding affinity would be highly desirable. To this end, we have introduced two urea groups in a porphyrin tweezer system as remote guest-binding modules, which provided significantly high binding affinity for chiral carboxylates without derivatization. Although the ECCD signal of the tweezer system manifested upon chiral carboxylate binding was relatively weak, great signal amplification was achieved upon additional binding of an achiral diamine guest to the porphyrin units.

[[]a] H. Yoon, C.-H. Lee, Prof. W.-D. Jang Department of Chemistry College of Science Yonsei University 50 Yonsei-ro Seodaemun-gu, Seoul, 120-749 (Korea) Fax: (+82)2-364-7050 E-mail: wdjang@yonsei.ac.kr

quent removal of the trimethylsilyl group gave 8. Finally, dimerization of 8 yielded 1, which was unambiguously characterized by ¹H and ¹³C NMR and MALDI-TOF-MS analyses.

Compound 1 exhibited Soret absorption at 419 nm with a

remarkably strong shoulder at 409 nm (Figure 1), at which

the extinction coefficient in CH_2Cl_2 for the Soret band was determined to be $5.42 \times 10^5 \,\text{m}^{-1} \,\text{cm}^{-1}$. The absorption spec-

trum of the monomeric porphyrin (8) was also measured

and featured a single sharp Soret absorption at 409 nm that

was coincident with the position of the shoulder. Since the

monomeric porphyrin absorbs at the same position as the

shoulder of 1, the major absorption of 1 at 419 nm is proba-

bly caused by exciton coupling of the two porphyrin units.^[7]

Results and Discussion

The synthesis of the molecular tweezer (1) is outlined in Scheme 1. Briefly, iodination of *tert*-butyl aniline (2) afforded 3, which was reacted with trimethylsilylacetylene to give 4. The amine group in 4 was converted into an isocyanate group using triphosgene in CH_2Cl_2 , and then the isocyanate was further reacted with the amino-porphyrin (7) to generate the urea-bearing porphyrin derivative (8). The aminoporphyrin was synthesized from 5, 6, and dipyrrolomethane by an acid-catalyzed cross-condensation reaction and subsequent deprotection of the acetyl group. Metallation of the porphyrin was accomplished with $Zn(OAc)_2$, and subse-



bind to the Lewis acidic zinc porphyrin as an axial ligand,^[8] we expected that 1 could accommodate diamine guests. As shown in Figure 2a, the addition of 1,12-diaminododecane (DAD) to 1 resulted in a hypsochromic shift of the absorption maximum from 419 to 418.5 nm with clear isosbestic points at 379, 414, 511, 545, and 575 nm. The bandwidth of the Soret absorption band was also reduced along with disappearance of the shoulder after formation of the host-guest complex. The continuous variation method indicated that 1 and DAD formed a 1:1 host-guest complex with a very high binding affinity, the association constant for which was determined to be $1.2 \times$ $10^7 \,\mathrm{M}^{-1}$ through a nonlinear curve-fitting method using hypspec software. Because axial ligation by the amine increases the electron density of the zinc porphyrin, the absorption bands should exhibit a bathochromic shift. However, after complexation of DAD, 1 exhibited a hypsochromic shift of the absorption maximum but a bathochromic shift in the position of the shoulder. This spectroscopic change indicated that the two porphyrin units in 1 partially engage in the formation of а slipped-cofacial structure through $\pi - \pi$ interaction, which was diminished upon insertion of DAD between them.^[8]

Because an amine group can

Scheme 1. Synthesis of **1**. Reagents and conditions: a) I_2 , Ag_2SO_4 , EtOH, 25 °C, 2 h; b) trimethylsilylacetylene, CuI, $[Pd(PPh_3)_2Cl_2]$, Et₃N, 25 °C, 12 h; c) triphosgene, Et₃N, CH₂Cl₂, 0 °C, 2 h; d) dipyrromethane, TFA, CH₂Cl₂/MeOH, 25 °C, 12 h; e) *p*-chloranil, 25 °C, 4 h; f) HCl, EtOH, reflux, 2 h; g) **7**, reflux, 12 h; h) Zn(OAc)₂, CH₂Cl₂/MeOH, 25 °C, 12 h; i) TBAF, THF, 25 °C, 30 min; j) Cu(OAc)₂:H₂O, pyridine, 25 °C, 12 h.



Figure 1. UV/Vis absorption spectra of a) 1 (solid line) and b) 8 (broken line) in CD₂Cl₂ at 25 °C.

Meanwhile, the two urea moieties in **1** provide strong hydrogen-bond donor groups for binding anionic guest molecules.^[9] In fact, upon the addition of several different anionic guests, **1** exhibited a remarkably strong spectral shift with peak broadening, indicating a strong interaction between its two porphyrin units after anion binding (Figure 2b). The existence of clear isosbestic points and the result of the continuous variation method (Figure 2; Job's plot) indicated the formation of a very stable host–guest complex with a 1:1 binding stoichiometry, with the association constant for BLA exceeding 10^8 M^{-1} .

As mentioned above, the UV/Vis absorption spectrum of 1 is indicative of an interaction between the two porphyrin units of the tweezer. Variable-temperature (VT)-NMR and temperature-dependent UV/Vis absorption studies of 1 in $C_2D_2Cl_4$ were carried out to obtain further information on the initial conformation (Figure S1 in the Supporting Information). In the VT-NMR study, all peaks became sharp upon increasing the temperature, indicating rapid conformational interchange at high temperature. In contrast, all peaks disappeared on lowering the temperature due to poor solubility, with 1 precipitating from C₂D₂Cl₄ solution at low temperature. Although the VT-NMR experiments indicated slight downfield shifts of all peaks, the information provided was insufficient to confirm the initial conformation of 1. Therefore, the ¹H NMR spectrum of **8** was compared with that of 1 (Figure 3). The signals of the pyrrolic and meso protons of 1 were seen to be located further upfield than those of 8, indicating a partial overlapping of the porphyrin rings such that they were mutually affected by their respective ring currents. A UV/Vis absorption study in $C_2D_2Cl_4$ also indicated partial overlapping of the porphyrin rings (Figure S2 in the Supporting Information). The absorption spectrum of **1** in $C_2D_2Cl_4$ showed similar spectral shape to that in CH₂Cl₂, with the Soret absorption at 421 nm and a remarkably strong shoulder at 411 nm. The absorption intensity at 421 nm was gradually decreased on increasing the temperature, indicating a higher proportion of the trans-conformation at high temperature.

On the other hand, ¹H NMR study clearly showed spectral changes upon the addition of anionic guests. Upon addi-



Figure 2. UV/Vis spectroscopic titration of **1** with DAD and BLA in CH₂Cl₂ at 25 °C. Lower left and right graphs are binding isotherm and Job's plot, respectively. a) [DAD]/[**1**]=0-3.0, [**1**]= 2.15×10^{-6} M, Δ Abs was monitored at 408 nm, b) [BLA]/[**1**]=0-3.0, [**1**]= 2.15×10^{-6} M, Δ Abs was monitored at 420 nm.

tion of Boc-L-alanine (BLA) as the tetrabutylammonium salt, the urea NH signals exhibited a significant downfield shift, indicating the formation of strong hydrogen bonds between the carboxylate anion and the NH protons (Figure 3). At the same time, the signals of the pyrrole β and meso protons of the porphyrin were remarkably upfield shifted upon the addition of BLA. This strong upfield shift can be explained in terms of π - π interaction between the two porphyrin units. Due to the partial overlapping of the porphyrin

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

 GaA, Weinheim
 www.chemeurj.org

 Image: These are not the final page numbers!

FULL PAPER



a)

4

1.5

1.0

1.5

1.0

0.5

Figure 3. Partial ¹H NMR spectra of a) 8, b) 1, c) $1 \supset BLA$, and \exists d) $1 \supset BLA \cdot DAD$ in CD_2Cl_2 at 25 °C.



units by π - π interaction, the pyrrole β and meso protons were greatly influenced by the porphyrin ring-current effect. When both BLA and DAD were added to **1**, the degree of upfield shift of the pyrrole β and meso proton signals was greatly decreased. The distance between the two porphyrin units was increased by the insertion of DAD, and hence the influence of their ring-current effects was greatly decreased. In sharp contrast to the proton signals corresponding to the porphyrin units, the chemical shift of the urea NH protons was not changed by the addition of DAD, indicating that the urea groups maintained their hydrogen bonds.

In the presence of one equivalent of BLA, **1** was again titrated with DAD. In this experiment, the absorption spectrum exhibited a bathochromic shift with clear isosbestic points at 441, 444, 516, and 540 nm (Figure 4a). The bandwidth of the Soret absorption also narrowed. When **1** was ti-

Figure 4. UV/Vis spectroscopic titration of **1** with DAD and BLA in CH₂Cl₂ at 25 °C in the presence of 1 equivalent BLA and DAD, respectively. Lower left and right graphs are binding isotherm and Job's plot, respectively. a) [BLA]/[1 \supset DAD]=0-3.0, [1 \supset DAD]=2.13×10⁻⁶ M, Δ Abs was monitored at 420 nm, b) [DAD]/[1 \supset BLA]=0-3.0, [1 \supset BLA]=2.13×10⁻⁶ M, Δ Abs was monitored at 402 nm.

trated with BLA in the presence of one equivalent of DAD, the Soret absorption was slightly broadened with clear isosbestic points at 412, 429, and 559 nm, indicating simultaneous accommodation of the anion and diamine guests within **1** (Figure 4b). The results of the continuous variation method indicated the formation of 1:1 host-guest complexes for both the diamine and anionic guests.

Because 1 exhibited strong binding affinities towards both the diamine and anionic guests, the binding of chiral guests induced a conformational change of the host compound, which in turn generated ECCD signals. Therefore, CD spectra of 1 were measured at 20 °C upon the addition of several chiral amino acid derivatives. While 1 showed essentially no observable CD signals, a very weak negative CD signal appeared around the Soret region after the addition of three equivalents of BLA (Figure 5). Conversely, 1 exhibited a



Figure 5. CD spectra of 1 in CH_2Cl_2 in the presence of three equivalents of BLA or BDA, with or without DAD (scan number 8, scan speed 200 nm min⁻¹, 293 K, bandwidth = 1 nm, [1] = 1.69×10^{-6} M).

weak positive CD signal upon the addition of Boc-D-alanine (BDA) as the tetrabutylammonium salt. For many other amino acid derivatives, very weak induced CD signals appeared. Although induced CD signals appeared after the addition of chiral amino acid derivatives, the intensity of these signals was very weak and their signs were not perfectly correlated with the chirality of guest molecules. The low intensity of the induced CD signal upon carboxylate binding to **1** might be explained in terms of the formation of a slipped-cofacial structure due to the strong π - π interaction between the two porphyrin units. In addition, the binding of anionic guests at the remote site away from the porphyrin units will not cause strong ECCD signals

at the Soret band.

Because the strong interaction between the two porphyrin units in **1** was diminished by the insertion of DAD, we speculated that a conformational change of the porphyrin units might be enhanced by the addition of this diamine. Therefore, CD spectra were again meas-



Figure 6. Energy-minimized structures of a) 1, b) 1>BLA, and c) 1>BLA•DAD with omission of alkyl chains.

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim www.chemeurj.org _____5 These are not the final page numbers!

ured after the addition of three equivalents of DAD. Very interestingly, the weak induced CD signals observed with BLA were converted into a strong positive couplet after the addition of DAD (Figure 5). Based on this information, we re-evaluated the CD signal amplification of **1** upon the addition of several other amino acid derivatives. All of the tested L-amino acid derivatives gave rise to a positive CD couplet, while D-amino acid derivatives gave the opposite signals. Furthermore, the intensities of the CD couplet were sufficiently strong to identify the absolute stereo-configurations (Table 1). Because **1** exhibited a significant and useful

Table 1. CD amplitudes of **1** induced by the binding of carboxylates in the presence of 3 equiv DAD at 20 $^{\circ}$ C in a 1 cm optical cell.^[a]

Guests	Asymmetry	1st/nm [$\Delta \varepsilon$]	2nd/nm [$\Delta \varepsilon$]	$A_{\rm CD}^{[b]}$
BDA	R	427.2	419.2	-58.2
		[-32.5]	[+25.7]	
BLA	S	428.2	418.4	+61.4
		[+28.0]	[-33.4]	
Boc-D-Leu	R	426.9	418.9	-98.5
		[-56.0]	[+42.5]	
Boc-L-Leu	S	426.7	418.5	+102.3
		[+49.2]	[-53.1]	
Boc-D-Phe	R	427.6	418.4	-135.2
		[-67.9]	[+67.3]	
Boc-L-Phe	S	427.4	418.5	+136.1
		[+65.2]	[-70.9]	
Boc-L-Pro	S	428.3	418.5	+138.2
		[+70.4]	[-67.9]	
Boc-L-Ser	S	429.1	415.5	+14.7
		[+6.8]	[-7.9]	
Boc-L-Thr	S	426.3	418.7	+61.8
		[+32.5]	[-29.3]	
Boc-L-Val	S	427.7	418.6	+99.7
		[+48.5]	[-51.2]	
Mandelate	S	427.4	417.6	+72.3
		[+32.1]	[-40.2]	

[a] Solvent: CH₂Cl₂, scan speed: 200 nmmin⁻¹, scan number: 8, [1]= 1.69×10^{-6} M, [carboxylates]= 5.07×10^{-6} M, [b] L mol⁻¹ cm⁻¹.

CD signal response upon the addition of amino acid derivatives together with achiral DAD, computer-aided molecular modeling studies were carried out using the MM3 forcefield to study the possible underlying conformational changes.^[10] Depending on the initial conformation, the energy-minimized structure of **1** exhibited both *cis*- and *trans*-conformers (Figure 6a and Scheme 2). A slipped-cofacial structure of the two porphyrin units could be observed A EUROPEAN JOURNAL



Scheme 2. Proposed mechanism of CD enhancement of **1** by the binding of guests.

in the cis-conformation, which is consistent with the observations based on the electronic absorption spectra. Accordingly, upon formation of the BLA adduct with $1 (1 \supset BLA)$, 1 adopted the cis-conformation through the formation of multiple hydrogen bonds (Figure 6b). In this state, the two porphyrin units are spatially very close to each other due to their π - π interaction. More importantly, the directions of the transition dipole moments for the two porphyrin rings, which run from the 5- to the 15-position of the porphyrin units, become almost parallel in the $1 \supset BLA$ complex. This could be the reason for the generation of weak ECCD signals upon the addition of BLA to 1. Considering the addition of DAD to $1 \supset BLA$, the angle between the transition dipole moments of the two porphyrin units becomes much greater following insertion of the diamine. Therefore, the molecular chirality of the carboxylate may strongly induce either a right-handed or left-handed twist to the structure of 1. In fact, the angle between the transition dipole moments of the two porphyrin units in the energy-minimized structure of $1 \supset BLA \cdot DAD$ was much wider than that in $1 \supset BLA$, which presumably gave rise to the strong ECCD signals (Figure 6c). The process of chiral recognition can be summarized as shown in Scheme 2. The porphyrin tweezer adopts a *cis*-conformation by carboxylate inclusion due to the formation of multiple hydrogen bonds. In this process, the two urea groups are twisted in different directions depending on the chirality of the amino acid derivatives in order to minimize steric repulsion (Figure 7). Because of the chiral twist of the urea groups, 1 should exhibit ECCD signals. However, the electronic absorption of the urea groups lies in the invisible UV range. Moreover, the two porphyrin units in 1 form a slipped-cofacial structure due to the strong π - π interaction. Therefore, **1** exhibits only weak CD signals due to the parallel orientation of transition dipoles. On the other hand, the chiral distortion of the urea groups is transferred to the two porphyrin units by the insertion of DAD, because the strong π - π interaction is diminished. Therefore,



Figure 7. Twist of urea groups by the formation of multiple hydrogen bonding with carboxylate.

strong ECCD signals become observable upon the addition of DAD to a chiral carboxylate complex of **1**. In this process, the size and rigidity of the side groups of chiral amino acids greatly influence the intensity of the CD signals, with the order of signal intensity being as follows: $Pro \approx Phe >$ $Leu \approx Val > Thr \approx Ala > Ser$. The ECCD is induced by a structural twist of the two porphyrin units to minimize steric interaction, and bulky and rigid side groups will induce more twisted structures than smaller ones. The relative intensities of ECCD upon additions of Phe, Leu, Val, and Ala clearly reflect the influences of the side groups.

Conclusion

We have synthesized a new type of porphyrin-based molecular tweezer with two urea groups, which exhibits strong ECCD signals upon the simultaneous complexation of chiral carboxylate and achiral diamine guests. Because the two urea groups and porphyrin units exhibit significantly strong binding affinities towards chiral carboxylate and diamine guests, respectively, the molecular tweezer can be successfully utilized as a highly sensitive probe for the absolute stereochemical determination of chiral carboxylates. Unlike other porphyrin-based molecular tweezers, monodentate carboxylates can be directly utilized without derivatization to determine their molecular chirality. Therefore, the concept demonstrated herein provides a new strategy for the design of efficient chiral probes.

Experimental Section

Measurements: Electronic absorption spectra were recorded on a JASCO model V-660 spectrometer. CD spectra were recorded on a JASCO model J-815 spectrometer at 20°C in CH₂Cl₂ (scan speed: 200 nm min⁻¹, scan number: 8, bandwidth: 5 nm). ¹H and ¹³C NMR spectra were recorded on Bruker Advance DPX 250 and DPX 400 spectrometers at 25°C in CD₂Cl₂ or CDCl₃. MALDI-TOF-MS was performed on an Applied Biosystems 4700 proteomics analyzer with α -cyano-4-hydroxycinnamic acid as the matrix.

Synthesis of $3^{[111]}$ I₂ (3.57 g, 14.07 mmol) and Ag₂SO₄ (4.39 g, 14.07 mmol) were added to a solution of 4-*tert*-butylaniline (2.00 g, 13.4 mmol) in

EtOH (50 mL) and the mixture was stirred for 2 h at 25 °C and then filtered through Celite. The filtrate was concentrated and the residue was purified by column chromatography on silica eluting with CH₂Cl₂/hexane (1:3) to give **3** (1.1 g, 30%). ¹H NMR (250 MHz, CDCl₃, 25 °C): δ =7.62 (s, 1H), 7.19–7.16 (d, *J*=8 Hz, 1H), 6.72–6.69 (d, *J*=8 Hz, 1H), 3.96 (s, 2H), 1.26 ppm (s, 9H).

Synthesis of 4:[11] 3 (1.05 g, 3.81 mmol), CuI (36 mg, 0.19 mmol), and [Pd-(PPh₃)₂Cl₂] (133 mg, 0.19 mmol) were placed in a Schlenk flask. The flask was degassed under high vacuum and back-filled with N2; this process was repeated three times. Dried THF (15 mL), Et₃N (3 mL), and trimethvlsilvlacetvlene (0.94 mL, 15.24 mmol) were added. The reaction mixture was stirred for 16 h at 50 °C and then filtered through Celite. The filtrate was concentrated and then purified by column chromatography on silica eluting with CH_2Cl_2 /hexane (1:5) to give 4 (0.80 g, 86%). ¹H NMR (250 MHz, CDCl₃, 25 °C): $\delta = 7.30$ (s, 1H), 7.18–7.15 (d, J = 8 Hz, 1H), 6.66-6.63 (d, J=8 Hz, 1 H), 4.11 (s, 2 H), 1.26 (s, 9 H), 0.27 ppm (s, 9 H). Synthesis of 7:^[6b] TFA (2 mL, 26.12 mmol) was added to a solution of 5 (1.23 g, 7.51 mmol), 6 (2.72 g, 7.51 mmol), and dipyrromethane (2.20 g, 15.03 mmol) in CH₂Cl₂ (830 mL) and MeOH (170 mL), and the mixture was stirred for 12 h at 25 °C. Thereafter, p-chloranil (8.49 g, 34.2 mmol) was added and the reaction mixture was stirred for a further 4 h. It was then concentrated to a volume of 200 mL and chromatographed on silica gel eluting with 20% EtOAc/CH2Cl2. Without further purification, the product was dissolved in a mixture of EtOH (40 mL) and aqueous HCl (60 mL) and the solution was refluxed for 12 h. The reaction mixture was purified by column chromatography on silica, eluting with CH2Cl2, and the eluate was concentrated to dryness. The residue was recrystallized from CH₂Cl₂/hexane to give 7 as a reddish-purple powder (0.44 g, 8%). ¹H NMR (250 MHz, CDCl₃, 25 °C): $\delta = 10.29$ (s, 2H), 9.40–9.37 (m, 4H), 9.20-9.16 (m, 4 H), 8.08-8.05 (d, J=8 Hz, 2 H), 7.43 (s, 2 H), 7.14-7.11 (d, J=8 Hz, 2H), 6.92 (s, 1H), 4.18–4.13 (t, J=6.5 Hz, 4H), 4.06 (s, 2H), 1.92-1.86 (m, 4H), 1.53-1.28 (m, 10H), 0.90-0.85 (t, J=6 Hz, 6H), -3.08 ppm (brs, 2H); MALDI-TOF-MS: m/z: calcd for C₄₈H₅₅N₅O₂: 733.98 [*M*⁺]; found: 736.61.

Synthesis of $8{:}^{[12]}$ Triphosgene (0.14 g, 0.40 mmol) in CH_2Cl_2/Et_3N (3mL:0.2mL) was added to a solution of 4 (0.11 g, 0.40 mmol) in CH₂Cl₂ (7mL) and the mixture was stirred for 2 h at 0°C. Thereafter, a solution of 7 (0.2 g, 0.27 mmol) in CH₂Cl₂ (10 mL) was added, the reaction mixture was refluxed for 12 h, and then the solvents were evaporated. The residue was filtered through silica gel with EtOAc. Without further purification, the product was dissolved in 10% MeOH/CH2Cl2 (20mL) containing Zn(OAc)₂ (0.434 g, 1.98 mmol) and the solution was stirred for 12 h at 25 °C. The reaction mixture was purified by column chromatography on silica, eluting with 20% EtOAc/hexane to give a reddish-purple solid. The residue was redissolved in THF containing TBAF (78 µL, 0.27 mmol) and the solution was stirred for 30 min at 25 °C. The reaction mixture was purified by column chromatography on silica eluting with EtOAc/hexane (1:4) to give 8 (0.1 g, 37%). ¹H NMR (250 MHz, CDCl₃, 25°C): $\delta = 10.26$ (s, 2 H), 9.42–9.36 (m, 4 H), 9.28–9.26 (d, J = 4.5 Hz, 2 H), 9.06-9.04 (d, J=4.5 Hz, 2H), 8.16-8.13 (d, J=8 Hz, 2H), 7.45-7.40 (m, 4H), 7.33-7.26 (m, 2H), 7.03 (s, 1H), 6.92 (s, 1H), 5.73 (s, 2H), 4.17-4.10 (m, 4H), 3.36 (s, 1H), 1.92-1.81 (m, 4H), 1.48-1.23 (m, 10H), 0.88-0.82 ppm (m, 6H); MALDI-TOF-MS: m/z: calcd for C₆₁H₆₆N₆O₃Zn: 996.63 [*M*⁺]; found: 994.13.

Synthesis of 1:^[13] **8** (0.1 g, 0.10 mmol) and Cu(OAc)₂·H₂O (40 mg, 0.20 mmol) were placed in a Schlenk flask. The flask was degassed under high vacuum and back-filled with N₂; this process was repeated three times. Dry pyridine (10 mL) was added. The reaction mixture was stirred for 12 h at 25 °C, and then the solvent was evaporated. The residue was purified by column chromatography on silica eluting with EtOAc/hexane (1:3) to give **1** (80 mg, 40%). ¹H NMR (400 MHz, CD₂Cl₂, 25 °C): δ = 10.08 (s, 4H), 9.46 (brs, 4H), 9.30 (brs, 4H), 8.98 (brs, 4H), 8.89 (brs, 4H), 8.64 (brd, J = 5.6 Hz, 4H), 7.50 (s, 4H), 7.08 (s, 2H), 6.94 (s, 2H), 6.75 (brs, 6H), 6.70 (brs, 2H), 6.28 (brs, 2H), 4.78 (brs, 2H), 4.19–4.15 (t, J = 6.4 Hz, 8H), 1.59–1.05 (m, 20H), 0.84 ppm (s, 12H); ¹³C NMR (100 MHz, CD₂Cl₂, 25 °C): δ = 158.59, 150.41, 149.93, 149.89, 149.46, 145.04, 136.11, 133.88, 132.01, 131.94, 130.15, 128.09, 120.73, 120.29, 118.47, 115.02, 106.55, 101.17, 79.58, 79.07, 72.44, 68.85, 61.94, 34.36,

32.18, 31.28, 30.08, 29.82, 29.78, 29.61, 26.50, 23.00, 14.21 ppm; MALDI-TOF-MS: m/z: calcd for $C_{132}H_{96}N_{12}O_8Zn_2$: 1991.23 [M^+]; found: 1990.16.

Acknowledgements

This work was supported by a grant from the National Research Foundation (NRF) of Korea funded by the Korean government (MEST) (no. 2011-0001126). H. Yoon and C.-H. Lee acknowledge fellowships from the BK21 program of the Ministry of Education, Science and Technology, Korea.

- a) P. Metola, E. V. Anslyn, T. D. James, S. D. Bull, *Chem. Sci.* 2012, 3, 156–161; b) L. A. Joyce, M. S. Maynor, J. M. Dragna, G. M. da Cruz, V. M. Lynch, J. W. Canary, E. V. Anslyn, *J. Am. Chem. Soc.* 2011, 133, 13746–13752; c) D. Leung, S. O. Kang, E. V. Anslyn, *Chem. Soc. Rev.* 2012, 41, 448–479; d) X.-C. Li, D. Ferreira, Y. Ding, *Curr. Org. Chem.* 2010, 14, 1678–1697.
- [2] a) K. Claborn, C. Isborn, W. Kaminsky, B. Kahr, Angew. Chem.
 2008, 120, 5790-5801; Angew. Chem. Int. Ed. 2008, 47, 5706-5717;
 b) S. Allenmark, Chirality 2003, 15, 409-422; c) D. Parker, Chem.
 Rev. 1991, 91, 1441-1457; d) T. J. Wenzel, J. D. Wilcox, Chirality
 2003, 15, 256-270; e) Y. Okamoto, E. Yashima, Angew. Chem. 1998, 110, 1072-1095; Angew. Chem. Int. Ed. 1998, 37, 1020-1043; f) T.
 Ikai, Y. Okamoto, Chem. Rev. 2009, 109, 6077-6101; g) H. D. Flack,
 G. Bernardinelli, Chirality 2008, 20, 681-690.
- [3] a) G. Tarczay, G. Magyarfalvi, E. Vass, Angew. Chem. 2006, 118, 1807–1809; Angew. Chem. Int. Ed. 2006, 45, 1775–1777; b) S. Nieto, V. M. Lynch, E. V. Anslyn, H. Kim, J. Chin, J. Am. Chem. Soc. 2008, 130, 9232–9233; c) X. Peng, N. Komatsu, T. Kimura, A. Osuka, J. Am. Chem. Soc. 2007, 129, 15947–15953.
- [4] a) Q. Yang, C. Olmsted, B. Borhan, Org. Lett. 2002, 4, 3423-3426; b) X. Li, B. Borhan, J. Am. Chem. Soc. 2008, 130, 16126-16127; c) M. Anderson, K. Wilcox, M. Guericke, H. Chu, M. V. Wilson, E. Wilson, K. Lucas, A. E. Holmes, Chirality 2010, 22, 398-402; d) J.-K. Choi, G. Sargsyan, M. Shabbir-Hussain, A. E. Holmes, M. Balaz, J. Phys. Chem. B 2011, 115, 10182-10188; e) M. Tanasova, C. Vasileiou, O. O. Olumolade, B. Borhan, Chirality 2009, 21, 374-382; f) B. I. Morinaka, T. F. Molinski, Chirality 2008, 20, 1066-1070; g) D. Das, Z. Dai, A. E. Holmes, J. W. Canary, Chirality 2008, 20, 585-591; h) V. V. Borovkov, J. M. Lintuluoto, Y. Inoue, Org. Lett. 2000, 2, 1565-1568; i) G. Proni, G. Pescitelli, X. Huang, K. Nakanishi, N. Berova, J. Am. Chem. Soc. 2003, 125, 12914-12927; j) X. Huang, N. Fujioka, G. Pescitelli, F. E. Koehn, R. T. Williamson, K. Nakanishi, N. Berova, J. Am. Chem. Soc. 2002, 124, 10320-10335; k) N. Berova, G. Pescitelli, A. G. Petrovica, G. Proni, Chem. Commun. 2009, 5958-5980; 1) L. Xi, H. Fu, W. Yang, J. Yao, Chem. Commun. 2005, 492-494; m) D. S. Dalisay, S. Tsukamoto, T. F. Molinski, J. Nat. Prod. 2009, 72, 353-359; n) A. Sugasaki, M. Ikeda, M. Takeuchi, K. Koumoto, S. Shinkai, Tetrahedron 2000, 56, 4717-4723; o) O. Hirata, Y. Kubo, M. Takeuchi, S. Shinkai, Tetrahedron 2004, 60, 11211-11218.
- [5] a) G. Proni, G. Pescitelli, X. Huang, N. Q. Quraishi, K. Nakanishi, N. Berova, *Chem. Commun.* **2002**, 1590–1591; b) V. V. Borovkov, J. M. Lintuluoto, G. A. Hembury, M. Sugiura, R. Arakawa, Y. Inoue, *J. Org. Chem.* **2003**, 68, 7176–7192; c) X. Li, M. Tanasova, C. Vasileiou, B. Borhan, *J. Am. Chem. Soc.* **2008**, *130*, 1885–1893; d) Q. F. Liang, J. J. Liu, J. Chen, *Tetrahedron Lett.* **2011**, *52*, 3987–3991.
- [6] a) J. F. Lovell, C. S. Jin, E. Huynh, H. Jin, C. Kim, J. L. Rubinstein, W. C. W. Chan, W. Cao, L. V. Wang, G. Zheng, *Nat. Mater.* 2011, *10*, 324–332; b) C.-H. Lee, S. Lee, H. Yoon, W.-D. Jang, *Chem. Eur. J.* 2011, *17*, 13898–13903; c) W.-D. Jang, C.-H. Lee, M. Osada, M.-S. Choi, *J. Porphyrins Phthalocyanines* 2009, *13*, 787–793.
- [7] a) T. Yamaguchi, T. Kimura, H. Matsuda, T. Aida, Angew. Chem.
 2004, 116, 6510-6515; Angew. Chem. Int. Ed. 2004, 43, 6350-6355;
 b) F. Hajjaj, Z. S. Yoon, M.-C. Yoon, J. Park, A. Satake, D. Kim, Y. Kobuke, J. Am. Chem. Soc. 2006, 128, 4612-4623; c) N. C. Maiti, S.

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

www.chemeurj.org



These are not the final page numbers! **77**

Mazumdar, N. Periasamy, *J. Phys. Chem. B* **1998**, *102*, 1528–1538; d) Y. Egawa, R. Hayashida, J. Anzai, *Langmuir* **2007**, *23*, 13146–13150.

- [8] a) F. A. Walker, M. Benson, J. Am. Chem. Soc. 1980, 102, 5530–5538; b) M. Schmittel, S. K. Samanta, J. Org. Chem. 2010, 75, 5911–5919; c) C.-H. Lee, H. Yoon, W.-D. Jang, Chem. Eur. J. 2009, 15, 9972–9976; d) C.-H. Lee, H. Yoon, P.-S. Kim, S. Cho, D. Kim, W.-D. Jang, Chem. Commun. 2011, 47, 4246–4248.
- [9] a) B. P. Hay, T. K. Firman, B. A. Moyer, J. Am. Chem. Soc. 2005, 127, 1810–1819; b) M. Pittelkow, C. B. Nielsen, M. A. C. Broeren, J. L. J. van Dongen, M. H. P. van Genderen, E. W. Meijer, J. B. Christensen, Chem. Eur. J. 2005, 11, 5126–5135; c) M. A. C. Broeren, B. F. M. de Waal, M. H. P. van Gendern, H. M. H. F. Sanders, G. Fytas, E. W. Meijer, J. Am. Chem. Soc. 2005, 127, 10334–10343; d) J.-D. Lee, D. Jang, J.-I. Hong, B. Kor. Chem. Soc. 2010, 31, 2685–2688; e) J.-D. Lee, Y.-H. Kim, J.-I. Hong, J. Org. Chem. 2010, 75, 7588–7595; f) K. Calderon-Kawasaki, S. Kularatne, Y. H. Li, B. C.

Noll, W. R. Scheidt, D. H. Burns, *J. Org. Chem.* **2007**, *72*, 9081–9087; g) X. P. Bao, H. Zhang, Z. Zhang, L. Wu, Z. Y. Li, *Inorg. Chem. Commun.* **2007**, *10*, 728–730; h) G. Baggi, M. Boiocchi, L. Fabbrizzi, L. Mosca, *Chem. Eur. J.* **2011**, *17*, 9423–9439.

- [10] a) L. R. Schmitz, N. L. Allinger, J. Am. Chem. Soc. 1990, 112, 8307– 8315; b) J. B. Sorensen, A. H. Lewin, J. P. Bowen, J. Mol. Struct. (Theochem) 2003, 623, 145–158.
- [11] K.-J. Chang, D. Moon, M. S. Lah, K.-S. Jeong, Angew. Chem. 2005, 117, 8140–8143; Angew. Chem. Int. Ed. 2005, 44, 7926–7929.
- [12] M. Dudic?, P. Lhoták, I. Stibor, K. Lang, P. Proškova, Org. Lett. 2003, 5, 149–152.
- [13] M. Ikeda, A. Sugasaki, Y. Kubo, K. Sugiyasu, M. Takeuchi, S. Shinkai, *Chem. Lett.* 2001, 1266–1267.

Received: February 3, 2012 Revised: June 4, 2012

Published online: ■ ■ , 0000

Molecular tweezer for chirality deter-

mination: A new type of porphyrinbased molecular tweezer has been found to exhibit strong CD signals upon the simultaneous complexation of chiral carboxylate and achiral diamine guests (see scheme). By virtue of this unique feature of the tweezer, it can be utilized as a highly sensitive probe for the absolute stereochemical determination of chiral carboxylates.



Chirality Determination -

H. Yoon, C.-H. Lee, W.-D. Jang*.....

Absolute Stereochemical Determination of Chiral Carboxylates Using an Achiral Molecular Tweezer

Chem. Eur. J. 2012, 00, 0-0

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

www.chemeurj.org

9

These are not the final page numbers! 77