

PII: S0040-4039(96)00827-1

## Aziridine-2-Carboxylic Acid Mediated Asymmetric Synthesis of D-*erythro*and L-*threo*-Sphingosine from a Common Precursor

Franklin A. Davis\*, and G. Venkat Reddy

Department of Chemistry, Temple University, Philadelphia, PA 19122-2585

**Summary:** Short, two and three step asymmetric syntheses of L-*threo*- and D-*erythro*sphingosine (-)-**6a** and (-)-**9a**, respectively, in 58 and 31 percent overall yield were accomplished via the stereo and regioselective ring opening of N-sulfinylaziridine 2carboxylic acid **4**. Copyright © 1996 Elsevier Science Ltd

Enantiomerically pure aziridines serve as versatile intermediates for the asymmetric synthesis of biologically active materials because they undergo highly regio- and stereocontrolled ring opening reactions with nucleophiles.<sup>1</sup> This is particularly true for aziridine 2-carboxylic acids which afford  $\beta$ -substituted  $\alpha$ -amino acids upon opening.<sup>2</sup> Using *cis-N-(p*-toluenesulfinyl)-2-carbomethoxyaziridine 1, we recently described a highly efficient asymmetric synthesis of *threo-(-)*-2-amino-1-[(4-methylthio)phenyl]-1,3-propanediol (3), the key intermediate in the manufacture of the broad spectrum antibiotics thiamphenicol and florfenicol.<sup>3</sup> This procedure involved the regio-and stereospecific S<sub>N</sub>2 ring opening of the corresponding aziridine-2-methanol 2 with TsOH/H<sub>2</sub>O affording *threo-*3 in 93% yield with none of the *erythro-*3 isomer being detected (Scheme 1). A similar 2-amino-1,3-diol unit is found in sphingosine [*erythro-*(-)-2-amino-D-4(E)-octadecene-1,3-diol (9a)], the major constituent of the lipid backbone of the sphingolipids (e.g. ceramides, cerebrosides, sphingomyelins, gangliosides and glycosphingolipids).<sup>4</sup> The sphingolipids are important membrane components that play crucial roles in cell recognition events such as growth,

Scheme 1



Ar = p-MeSPh

differentiation, the immune response<sup>5</sup> and as receptors for HIV binding in cells lacking the CD4 receptor.<sup>6</sup> Furthermore, all of the sphingosine enantiomers are reported to be potent inhibitors of protein kinase C (PKC)<sup>7</sup> as well as stimulators of DNA synthesis and cell proliferation.<sup>8</sup> In this letter we describe the first aziridine-mediated asymmetric syntheses of L-*threo* and D-*erythro* sphingosines **6a** and **9a**, via a common intermediate and we also discoverd a new Pummerer induced rearrangement of N-sulfinyl aziridines.<sup>9</sup>

Synthesis of the less common L-*threo* sphingosine **6a** by the protocol outlined in Scheme 1 requires aziridine (2R,3R)-(-)-**4**<sup>13</sup> and would appear to be straightforward (Scheme 2). However, model studies with aziridine *rac*-**4** (R = *n*-C<sub>3</sub>H<sub>7</sub>) quickly demonstrated the problematic nature of this approach. Reduction to the aziridine methanol with lithium aluminum hydride (LAH) and other reducing reagents (DIBAL, NaBH<sub>4</sub>) gave the alcohol in low yield (ca 30%) and hydrolysis (TsOH/H<sub>2</sub>O) gave an inseparable mixture of stereo- and regioisomeric aminodiols. On the other hand, treatment of (-)-4 in acetone/TFA/H<sub>2</sub>O gave β-hydroxy-α-amino acid (-)-5 as a single isomer in 72% yield following flash chromatography. This hydrolysis proved to be very sensitive to the reaction conditions with higher temperatures and other solvents (MeCN, THF) giving multiple, unidentified products, and up to 10% of the *erythro*-isomer. Reduction of (-)-5 with LiBH<sub>4</sub> gave L-(-)-*threo* sphingosine (**6a**) in 80% yield. This was quantitatively acetylated to the more stable triacetate (+)-**6b**. Both materials had spectral properties identical with literature values.<sup>12a</sup>



(a) 50% aqueous TFA/acetone, 0 °C, 24 h, 72%; (b) TFAA/CH<sub>2</sub>Cl<sub>2</sub>, 35 °C, 15 min., radial chromatography, 59%; (c) LiBH<sub>4</sub>/MeOH, rt., 30 min. 80%; (d) LiBH<sub>4</sub>/MeOH, rt., 1 h, 71%; (e) K<sub>2</sub>CO<sub>3</sub>/EtOH, 50 °C, 5 h, 75%; (f) Ac<sub>2</sub>O/pyr. 100%.

4350

Selectivity in aziridine ring-opening depends on the ring substituents, the activating group on nitrogen and the reaction conditions because competitive  $S_N1$  and  $S_N2$  process are involved. Since anti selectivity generally predominates, it would appear impossible to obtain the naturally occurring D-(-)-*erythro*-sphingosine **9a** as the major isomer from cis-aziridine (-)-**4**. Remarkably, treatment of (-)-**4** with trifluoroacetic anhydride (TFAA) in  $CH_2Cl_2$  at 35 °C resulted in an 88:12 mixture of *erythro*- and *threo*-**7** which were separated by radial chromatography (ether:*n*-pentane) to give the major isomer in 59% yield. Formation of *erythro*-**7** from cis-**4** requires the highly unusual attack of the nucleophile from the same side as the departing amino group. While details of this transfromation are at present unclear, we suggest that activated sulfoxide complex **10**, well precedented in the Pummerer rearrangement of sulfoxides,<sup>14</sup> undergoes a stereospecific [3,3]-sigmatropic rearrangement of the trifluoromethyl acetoxy group to a developing carbocation or ion pair **11**. The aziridine N-activating group which promotes this type of ring-opening is the amino-sulfonium salt.



Despite literature to the contrary,<sup>15</sup> all attempts to remove the *N*-trifluoroacetyl group in *erythro*-(-)-**7** by reaction with aqueous K<sub>2</sub>CO<sub>3</sub>/EtOH failed, affording instead the retro-aldol aldehyde (*E*)-hexadecenal in 50% yield.<sup>16</sup> This problem was solved by reduction of (-)-**7** with LiBH<sub>4</sub> to give (-)-**8** in 80% and then by hydrolysis with aqueous K<sub>2</sub>CO<sub>3</sub>/EtOH at 50 °C for 5 h. *erythro*-(-)-Sphingosine **9a** was isolated by flash chromatography in 71% yield and was identical in all respects with an authentic sample.<sup>12a</sup>

In summary, short two and three step asymmetric syntheses of *threo* and *erythro* sphingosine (-)-**6a** and (-)-**9a**, respectively, in 58 and 31 % overall yield from the N-sulfinylaziridine 2-carboxylic acid **4** are described. Significantly, the discovery of a novel Pummerer induced rearrangement of N-sulfinylaziridines makes it possible for the first time to prepare both *erythro* and *threo*  $\beta$ -hydroxy- $\alpha$ -amino acid derivatives from a single precursor.

Acknowledgments. We thank Ping Zhou for helpful discussion. Financially support of this work by the National Institutes of Health is gratefully acknowledge.

## **References and Notes**

- For a review on optically active aziridines see: Tanner, D. Angew. Chem., Int. Ed. Engl. 1994, 1. 33. 599.
- For leading references to optically active aziridine 2-carboxylic acids see: 2. Davis, F. A.; Zhou, P.; Reddy, G. V. J. Org. Chem. 1994, 59, 3243.
- Davis, F. A.; Zhou, P. Tetrahedron Lett. 1994, 35, 7525. 3.
- Kanfer, J. N.; Hakomori, S. Sphingolipid Biochemistry, Plenum Press: New York 1983. 4.
- Radin, N. S.; Shayman, J. A.; Inokuchi, J. Avd. Lipid Res., 1993, 26, 183. Springer, T. A.; Lasky, L. A. Nature 1991, 349, 196. Feizi, T. Trends Biochem. Sci. 1991, 16, 5. 84. Karlsson, K.-A. Trends Pharm. Sci. 1991, 12, 265.
- Harouse, J. M.; Bhat, S.; Spitalnik, S. L.; Laughlin, M.; Stefano, K.; Silberberg, D. H.; 6. Gonzalez-Scarano, F. Science 1991, 253, 320.
- Merrill, A. H. Jr.; Nimkar, S.; Menaldino, D.; Hannun, Y. A.; Loomis, C.; Bell, R. M.; Tyahi, S. R.; 7. Lambeth, J. D.; Stevens, V. L.; Hunter, R.; Liotta, D. C. *Biochemistry*, **1989**, *28*, 3138. Zhang, H.; Buckley, N. E.; Gibson, K.; Spiegel, S.; *J. Biol. Chem.* **1990**, *265*, 76.
- 8. Olivera, A.; Spiegel, S. Glyococonjugate J. 1992, 9, 110.
- (±)-Sphingosine triacetate<sup>10</sup> and (±)-sphinganine triacetate<sup>11</sup> its saturated analog, have 9. been prepared from trans-aziridine 2-methanols with varying degrees of success.
- 10.
- Cardillo, G.; Orena, M.; Sandri, S.; Tomasini, C. *Tetrahedron* **1986**, *42*, 917. Bongini, A.; Cardillo, G.; Orena, M.; Sandri, S.; Tomasini, C. J. C. S. Perkin Trans I, **1986**, 11. 1339. Bongini, A.; Cardillo, G.; Orena, M.; Sandri, S.; Tomasini, C. J. C. S. Perkin Trans I, **1986**, 1345.
- For leading references to the asymmetric synthesis of sphingosine see: (a) Enders, D.; 12. Whitehouse, D. L.; Runsink, J. Chem. Eur. J. 1995, 1, 382. (b) Hudlicky, T.; Nugent, T.; Griffith, W. J. Org. Chem. 1994, 59, 7944. (c) Nimkar, S.; Menaldino, D.; Merrill, A. H.; Liotta, D. Tetrahedron Lett. 1988, 29, 3037.
- Davis, F. A.; Reddy, G. V.; Liu, H. J. Am. Chem. Soc. 1995, 117, 3651. 13.
- For recent examples of the Pummerer rearrangement using TFAA see: Jung, M. E.; Kim, C.; 14. von dem Bussche, L. J. Org. Chem. 1994, 59, 3248. Page, P. C. B.; Shuttleworth, S. J.; McKenzie, M. J.; Schilling, M. B.; Tapolczay, D. J. Synthesis, 1995, 73. Bueno, A. B.; Carreno, M. C.; Garcia Ruano, J. L.; Rubio, A. Tetrahedron: Asymmetry 1992, 3, 251.
- Newman, H. J. Org. Chem. 1965, 30, 1287. Franck, B.; Lubs, H. J. Angew. Chem., Int. 15.
- Ed. Engl. 1968, 7, 223. Imazawa, M.; Eckstein, F. J. Org. Chem. 1979, 44, 2039. See Solladie-Cavallo, A.; Koessler, J. L. J. Org. Chem. 1994, 59, 3240. 16.

(Received in USA 11 April 1996; revised 25 April 1996; accepted 26 April 1996)