Contents lists available at ScienceDirect

ELSEVIER



journal homepage: www.elsevier.com/locate/bmcl

Synthesis and evaluation of a cyclic imine derivative conjugated to a fluorescent molecule for labeling of proteins

Hai-Ming Guo^a, Maki Minakawa^a, Lynn Ueno^b, Fujie Tanaka^{a,*}

^a Department of Molecular Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA ^b Department of Molecular and Experimental Medicine, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

ARTICLE INFO

ABSTRACT

Article history: Received 12 November 2008 Revised 15 December 2008 Accepted 17 December 2008 Available online 25 December 2008

Keywords: Imine Tyrosine Protein modification

Covalent labeling reactions of proteins are required for preparation of protein conjugates that are useful in research and medicine. For example, antibody–drug conjugates are more efficient therapeutics than antibody alone or drug alone.¹ When conjugated to protein, small molecule drugs can be selectively delivered to targets, reducing side effects and extending the half-life of the drugs.^{1,2} Fluorescently labeled proteins can be traced by detection of the fluorescence.³ Currently available labeling methods include labeling at lysine with succinimidyl ester derivatives, labeling at cysteine with maleimide derivatives, and labeling of an unnatural amino acid residue incorporated in protein sequence during its expression.^{1c,d,4,5} Small peptide tag-based strategies for labeling of proteins have also been developed.⁶ Most of the methods have advantages and disadvantages.

To provide labeling reactions complementary to the existing reactions, we have recently developed cyclic imines that covalently react with phenols, including tyrosine residues, in aqueous solutions⁷ (Scheme 1). The reactions of the imines with phenols proceeded over a wide pH range without the need of additional catalysts.⁷ The imines were relatively stable and were resistant to hydrolysis in aqueous solutions but efficiently reacted with phenols.⁷

Tyrosine residues are observed less often on the surface of folded proteins than lysines or carboxylic acid-containing residues,⁸ which are commonly used as labeling sites. Thus, accessible tyrosine is an attractive covalent labeling site.^{7–9} For proteins and peptides that do not have an accessible tyrosine, a tyrosine residue

* Corresponding author. E-mail address: ftanaka@scripps.edu (F. Tanaka). can be added by usual site-directed mutagenesis or by usual peptide synthesis. Although imines prepared from formaldehyde and aniline derivatives in situ have been used for labeling of proteins at tyrosine,⁸ the optimal pH range for this reaction is 5.5–6.5 and the reaction does not proceed above pH 8.^{8a} In addition, formaldehyde cannot be used in many cases; for example, use of formaldehyde is not suitable for labeling of proteins at tyrosine inside living cells or in the presence of living cells. Reactions at tyrosine using reagents other than imines typically require additional catalysts or multistep conversions after the first reaction step on the phenol moiety of tyrosine.^{9a–c} Use of the imine shown in Scheme 1 may bypass the limitations and disadvantages of the existing labeling methods.

A cyclic imine conjugated to a fluorescent dansyl group was synthesized and used for covalent labeling of

proteins. The covalent attachment to proteins was confirmed by gel electrophoresis and mass analysis.

To demonstrate the use of the cyclic imines shown in Scheme 1 for covalent attachment of non-native molecules to proteins and peptides, here, we have synthesized a conjugate of the imine with a fluorescent dansyl group and examined the reactions of the dansyl-conjugated imine with proteins and peptides.







© 2008 Elsevier Ltd. All rights reserved.

⁰⁹⁶⁰⁻⁸⁹⁴X/ $\$ - see front matter \odot 2008 Elsevier Ltd. All rights reserved. doi:10.1016/j.bmcl.2008.12.071





The imine derivative linked to a fluorescent dansyl group was synthesized as shown in Scheme 2.¹⁰ Diazidation of **1** afforded a mixture of stereoisomers of 2; the major isomer was expected to have trans-diazide groups due to the procedures used.¹¹ Hydrolysis of the ester with LiOH afforded 3, which was converted to succinimidyl ester 4. Reaction of 4 with amine 5, prepared from the corresponding diamine and dansyl chloride, afforded compound 6. Diamine 7 was obtained by reduction of the azide groups by hydrogenation with Pd-C under H₂ in EtOAc-MeOH (3:1). The solvents for this reduction were critical; reduction of 6 with Pd-C under H₂ but in MeOH mainly afforded monoamine, which might have been generated via elimination of HN₃. Diamine 7 was then transformed to imine **8** as a mixture of regioisomers (**8a** + **8b**). To confirm the imine functionality of 8, reaction of 8 with p-cresol was performed in the presence of TFA (Scheme 3).¹² Formation of addition product 9 confirmed the presence of the imine functionality in compound 8.

Imine **8** was used for modification of proteins and peptides. Reaction of peptide was performed using GlyTyr or AlaTyrAla (90 mM) with **8** (20 mM) in 10% DMSO/100 mM sodium phosphate, pH 6.0 for 48 h at 37 °C. Imine **8** was completely consumed as judged by TLC analysis of the reaction mixture. Formation of the addition products were confirmed by mass analysis.¹³

To test reaction of **8** with proteins, lysozyme from chicken egg white, α -chymotrypsinogen A type II from bovine pancreas, myoglobin from equine skeletal muscle, carbonic anhydrase isozyme II from bovine erythrocytes, and cytochrome *C* from horse heart were used. Reactions of proteins (475 μ M) were performed in the presence of **8** (5 mM) in 5% DMSO/100 mM sodium phosphate, pH 6.0 for 4 days at 37 °C and were analyzed by gel electrophoresis (Fig. 1). Gel bands corresponding to modified proteins were fluo-



Scheme 3.



Figure 1. Analysis of proteins modified with **8** after separation by gel electrophoresis. Lane a, protein standard marker; b, lysozyme; c, chymotrypsinogen (see text); d, myoglobin; e, carbonic anhydrase; f, cytochrome C. (A) Analysis of fluorescence of dansyl group attached to proteins under UV 312 nm. (B) Analysis of proteins by Coomassie staining.

rescent under a UV lamp (312 nm). Unmodified proteins were not fluorescent on a gel under the UV lamp (data not shown). Gel analysis showed that lysozyme was most easily modified among proteins tested (lane b). Mass analysis of the reaction mixture of lysozyme with 8 showed the formation of the mono-addition product.¹⁴ We previously demonstrated that the imine derivative shown in Scheme 1, prepared from trans-diaminocyclohexane, selectively reacted at tyrosine and did not react with other natural amino acid residues except cysteine thiol.⁷ Thus, reaction of **8** with lysozyme probably occurred at tyrosine. Cytochrome C was not labeled with 8 under the conditions used (lane f). Myoglobin and carbonic anhydrase were slightly modified (lanes d and e).¹⁵ Chymotrypsinogen was hydrolyzed into small fragments by autolysis and by catalysis of generated chymotrypsin under the conditions used (lane c). Due to this hydrolysis, we were unable to evaluate labeling of chymotrypsinogen with 8. These results indicated that 8 did not inhibit the serine protease activity under the conditions used; this implies that the nucleophilic serine in the active site of chymotrypsin did not efficiently react with imine 8.

Mass analysis of the reaction of **8** with lysozyme at 48 h suggested that the reaction of **8** was slower than that of the imine synthesized from *trans*-cyclohexanediamine shown in Scheme 1. We previously observed that reaction of the *trans*-isomer of the imine shown in Scheme 1 was faster than the corresponding *cis*-isomer.⁷ Relative stereochemistries of the two nitrogen functionalities and the amide carbonyl group on the cyclohexane ring of **8** may affect the reactivity of the imine moiety. In order to increase the reaction rate, stereoselective synthesis of **8** may be required. Alternatively, introduction of moieties that provide noncovalent interactions with the target protein may be used to enhance the reaction rate of the imine with the target.^{9d}

In these reactions, excess of compound **8** was used compared to protein (5 mM vs 475 μ M). After protein was separated by gel filtration or C18 column, compound **8** was recovered by extraction with organic solvent (such as CH₂Cl₂). This recoverable feature of the imine should be beneficial when a toxic drug or an expensive molecule is covalently attached to protein.

In summary, we have synthesized a fluorescent conjugate of cyclic imines that we have recently developed for covalent reactions with phenols, including accessible tyrosine. We have demonstrated that the imine derivative can be used to fluorescently label proteins, although the reaction of the imine was relatively slow. The imine derivative was stable in aqueous solutions and recoverable after reactions. Reactions of the imine derivative proceeded without the need of additional catalysts. This system does not require formaldehyde or multistep conversions after the first reaction step on tyrosine and thus is an improvement over previously reported labeling systems. Further studies and improvements on the imine derivatives described here are under investigation.

Acknowledgments

This study was supported by NIH Grant R21 GM078447. We thank Prof. Peter K. Vogt for helpful technical discussions.

References and notes

- (a) Schrama, D.; Reisfeld, R. A.; Becker, J. C. Nat. Rev. Drug Discov. 2006, 5, 147;
 (b) Wu, A. M.; Senter, P. D. Nat. Biotechnol. 2005, 23, 1137;
 (c) Hamann, P. R.; Hinman, L. M.; Hollander, I.; Beyer, C. F.; Lindh, D.; Holcomb, R.; Hallett, W.; Tsou, H.-R.; Upeslacis, J.; Shochat, D.; Mountain, A.; Flowers, D. A.; Bernstein, I. Bioconj. Chem. 2002, 13, 47;
 (d) Lillo, A. M.; Sun, C.; Gao, C.; Ditzel, H.; Parrish, J.; Gauss, G.-M.; Moss, J.; Felding-Habermann, B.; Wirsching, P.; Boger, D. L.; Janda, K. D. Chem. Biol. 2004, 11, 897.
- Vincent, S.; Thomas, A.; Brasher, B.; Benson, J. D. Nat. Biotechnol. 2003, 21, 936.
 Keppler, A.; Pick, H.; Arivoli, C.; Vogel, H.; Johnsson, K. Proc. Natl. Acad. Sci. U.S.A.
- **2004**, *101*, 9955.

- (a) Prescher, J. A.; Bertozzi, C. R. Nat. Chem. Biol. 2005, 1, 13; (b) Antos, J. M.; Francis, M. B. Curr. Opin. Chem. Biol. 2006, 10, 253.
- (a) Link, A. J.; Tirrell, D. A. J. Am. Chem. Soc. 2003, 125, 11164; (b) Agard, N. J.; Prescher, J. A.; Bertozzi, C. R. J. Am. Chem. Soc. 2004, 126, 15046; (c) Saxon, E.; Bertozzi, C. R. Science 2000, 287, 2007; (d) Dirksen, A.; Hackeng, T. M.; Dawson, P. E. Angew. Chem., Int. Ed. 2006, 45, 7581; (e) Crich, D.; Banerjee, A. J. Am. Chem. Soc. 2007, 129, 10064; (f) Tanaka, K.; Masuyama, T.; Hasegawa, K.; Tahara, T.; Mizuma, H.; Wada, Y.; Watanabe, Y.; Fukase, K. Angew. Chem., Int. Ed 2008, 47, 102.
- (a) Tanaka, F.; Fuller, R.; Asawapornmongkol, L.; Warsinke, A.; Gobuty, S.; Barbas, C. F., III *Bioconj. Chem.* 2007, *18*, 1318; (b) Tanaka, F.; Fuller, R. *Bioorg. Med. Chem. Lett.* 2006, *16*, 4059.
- 7. Minakawa, M.; Guo, H.-M.; Tanaka, F. J. Org. Chem. **2008**, 73, 8669.
- 8. (a) Joshi, N. S.; Whitaker, L. R.; Francis, M. B. J. Am. Chem. Soc. 2004, 126, 15942;
- (b) McFarland, J. M.; Joshi, N. S.; Francis, M. B. J. Am. Chem. Soc. 2008, 130, 7639.
 (a) Hooker, J. M.; Kovacs, E. W.; Francis, M. B. J. Am. Chem. Soc. 2004, 126, 3718; (b) Tilley, S. D.; Francis, M. B. J. Am. Chem. Soc. 2006, 128, 1080; (c) Hass, J. A.; Frederick, M. A.; Fox, B. G. Protein Expr. Purif. 2000, 20, 274; (d) Koshi, Y.; Nakata, E.; Miyagawa, M.; Tsukiji, S.; Ogawa, T.; Hamachi, I. J. Am. Chem. Soc. 2008, 130, 245.
- Compound 2. To a mixture of Mn(OAc)₃·2H₂O (16.1 g, 60 mmol) and NaN₃ (6.50 g, 100 mmol) in CH₃CN (180 mL), compound 1 (2.75 mL, 20 mmol) was added followed by trifluoroacetic acid (TFA, 20 mL) at -20 °C under Ar.¹¹ After 5 min, the mixture was added to saturated NaHSO₃ and extracted with CH₂C1₂. Usual workup and purification by silica gel flash column chromatography afforded 2⁷ (3.83 g, 85%).*Compound* 3. To a solution of 2 (3.38 g, 17 mmol) in THF (20 mL) was added to solution of LiOH (1.43 g, 34 mmol) in water (20 mL) at room temperature and the mixture was stirred for 15 h. After THF was evaporated in vacuo, 3 N HCl was added to adjust the pH to 3 and the mixture was extracted with CH₂C1₂. Usual workup and purification by silica gel flash column chromatography (EtOAc/hexane) afforded 3 (3.27 g, 98%).¹ H NMR (400 MHz, CDCl₃): δ 11.1 (br s, 1H), 3.94 to 3.49 (m, 1H), 3.42 to 3.21 (m, 1H), 2.82 to 2.68 (m, 1H), 2.46 to 2.27 (m, 1H), 2.16 to 1.85 (m, 2H), 1.83 to 1.40 (m, 3H).

Compound 4. A mixture of 3 (2.64 g, 12.6 mmol), N-hydroxysuccinimide (2.17 g, 18.9 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (3.62 g, 18.9 mmol), and DMAP (5.0 mg, 0.04 mmol) in CH₂Cl₂ (40 mL) was stirred at room temperature for 24 h. The reaction mixture was added to H₂O and extracted with CH₂Cl₂. Usual workup and purification by silica gel flash column chromatography (EtOAc/hexane) afforded 4 (3.28 g, 85%). ¹H NMR (400 MHz, CDCl₃): δ 3.73 to 3.69 (m, 1H), 3.44 to 3.28 (m, 1H), 3.12 to 3.08 (m, 1H), 2.85 (s, 4H), 2.41 to 2.35 (m, 1H), 2.24 to 2.14 (m, 1H), 2.04 to 1.92 (m, 1H), 1.83 to 1.66 (m, 3H). Compound 5. To a solution of 1,2-bis(2-aminoethoxy)ethane (2.5 mL, 17.0 mmol) in CH₂Cl₂ (20 mL), a solution of dansyl chloride (467 mg, 1.73 mmol) in CH₂Cl₂ (20 mL) was added dropwise at 0 °C. After stirring at room temperature for 1 h, the mixture was acidified with 1 N HCl and washed with CH₂Cl₂. The aqueous layer was basified (pH 9) with 3 N NaOH and extracted with CH₂Cl₂. Organic layers were combined, washed with brine, dried over Na₂SO₄, filtered, and concentrated in vacuo to afford 5 (630 mg, 95%). ¹H NMR (400 MHz, CDCl₃): δ 8.52 (d, J = 8.8 Hz, 1H), 8.34 (d, J = 8.8 Hz, 1H), 8.23 (dd, J = 7.2 Hz, 1.2 Hz, 1H), 7.56 to 7.48 (m, 2H), 7.17 (d, J = 7.6 Hz, 1H), 3.50 to 3.46 (m, 4H), 3.41 to 3.39 (m, 4H), 3.11 to 3.09 (m, 4H), 2.87 (s, 6H).

Compound **6**. A mixture of **5** (612 mg, 1.60 mmol) and **4** (493 mg, 1.60 mmol) in CH₃CN (0.5 mL)–MeOH (1.5 mL) was stirred for 15 h at room temperature. Solvents were evaporated in vacuo and the residue was purified by silica gel flash column chromatography (CH₂Cl₂/EtOAc, 4:1) to afford **6** (322 mg, 47%). ¹H NMR (400 MHz, CDCl₃): δ 8.55 (d, J = 8.8 Hz, 1H), 8.30 (d, J = 8.8 Hz, 1H), 8.24 (d, J = 7.2 Hz, 1H), 7.57 to 7.50 (m, 2 H), 7.19 (d, J = 7.6 Hz, 1H), 6.58 (br s, 1H), 5.90 (t, J = 5.6 Hz, 1H), 3.85 to 3.80 (m, 1H), 3.59 to 3.57 (m, 2H), 3.54 to 3.45 (m, 8H), 3.37 to 3.32 (m, 1H), 3.12 to 3.08 (m, 2H), 2.89 (s, 6H), 2.56 to 2.54 (m, 1H), 2.22 to 2.16 (m, 1H), 1.90 to 1.81 (m, 2H), 1.76 to 1.71 (m, 1H), 1.61 to 1.53 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 174.7, 152.1, 134.0, 130.8, 129.8, 129.6, 129.3, 128.6, 123.1, 118.2, 115.2, 115.2, 63.7, 63.6, 60.1, 45.3, 43.0, 39.1, 38.5, 29.8, 26.9, 25.3, 24.6. HRMS: calcd for C₂₅H₃₆N₉O₅S (MH⁺) 574.2555, found 574.2561.

Compound **7**. A mixture of **6** (322 mg, 0.56 mmol) and 10% Pd/C (32 mg) in MeOH (5 mL)–EtOAc (15 mL) was stirred under H_2 for 24 h at room temperature. The reaction mixture was filtered through Celite and concentrated in vacuo to afford **7** (290 mg, quant). ¹H NMR (400 MHz, CD₃OD): δ 8.52 (d, J = 8.8 Hz, 1H), 8.17 (dd, J = 7.2 Hz, 1.2 Hz, 1H), 7.56 to 7.51 (m, 2H), 7.23 (d, J = 7.6 Hz, 1H), 3.78 to 3.74 (m, 1H), 3.46 to 3.36 (m, 2H), 3.36 to 3.33 (m, 2H), 3.30 to 3.25 (m, 7H), 3.02 to 2.98 (m, 2H), 2.83 (s, 6H), 2.55 to 2.52 (m, 1H), 2.09 to 2.03 (m, 1H), 1.83 to 1.76 (m, 2H), 1.64 to 1.51 (m, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 17.46, 151.8, 135.1, 130.3, 130.2, 129.8, 129.6, 129.1, 128.18, 128.15, 128.12, 128.0, 123.1, 118.9, 115.16, 115.14, 70.23, 70.20, 70.2, 70.1, 70.0, 69.4, 56.1, 53.4, 45.3, 42.8, 39.5, 39.2, 26.1. HRMS: calcd for C₂₅H₄₀N₅O₅S (MH⁺) 522.2745, found 522.2745.

Compound **8** (*8a* + *8b*). To a solution of **7** (406 mg, 0.78 mmol) in CF₃CH₂OH (15 mL), a solution of ethyl glyoxylate polymer form (45–50% in toluene, 218 μ L, 0.78 mmol) in CF₃CH₂OH (15 mL) was added dropwise over 24 h at room temperature. The mixture was further stirred for 6 days. After solvents were removed in vacuo, the residue was purified by silica gel flash column chromatography (CH₂Cl₂/EtOH, 20:1) to afford **8** (259 mg, 60%). ¹H NMR (400 MHz, CDCl₃): δ 8.56 to 8.53 (m, 1H), 8.31 to 8.29 (m, 1H), 8.25 to 8.82 (m, 1H), 7.72 to 7.68 (m, 1H), 7.56 to 7.50 (m, 21H), 7.19 to 7.17 (m, 1H), 6.80 to 6.39 (m, 1H), 6.30 (m, 1H), 6.28 to 6.25 (m, 1H), 3.72 to 3.40 (m, 12H), 3.13 to 3.07 (m, 2H), 2.89 (s, 6H), 2.75 to 1.30 (m, 7H). ¹³C NMR (100 MHz, CDCl₃): δ 174.5, 173.8, 157.8, 129.55, 129.4, 129.3, 128.16, 128.13, 123.1, 118.8, 118.7, 115.2, 115.1, 70.4, 70.3, 70.2, 70.1, 69.3, 69.2, 62.8, 59.2, 54.0, 50.4, 45.3, 42.9, 42.8, 39.4, 39.2, 39.1, 38.1

33.0, 32.2, 28.0, 27.5, 27.4, 25.5. HRMS: calcd for C₂₇H₃₈N₅O₆S (MH⁺) 560.2537, found 560.2543.

- 11. Snider, B. B.; Lin, H. Synth. Commun. 1998, 28, 1913.
- 12. Compound 9. A mixture of compound 8 (136 mg, 0.24 mmol) and 4-methylphenol (39.4 mg, 0.36 mmol) in CF₃COOH (1 mL)-CH₂Cl₂ (1 mL) was stirred for 24 h at room temperature. After solvents were removed in vacuo, residue was dissolved in CH₂Cl₂ and was washed with saturated NaHCO₃. The aqueous layer was extracted with CH₂Cl₂. Usual workup and purification by silica gel flash column chromatography (CH₂Cl₂/MeOH, 30:1) afforded **9** (88 mg, 54%). ¹H NMR (400 MHz, CDCl₃): δ 8.54 (d, *J* = 8.8 Hz, 1H), 8.30 to 8.27 (m, 1H), 8.24 to 8.20 (m, 1H), 7.55 to 7.49 (m, 2H), 7.17 (d, J = 7.2 Hz, 1H), 7.06 (s, 1H), 6.99 to 6.93 (m, 1H), $\begin{array}{l} 6.78\ to\ 6.75\ (m,\ 1H+1H\times1/2), 6.65\ (m,\ 1H\times1/2), 6.45\ (s,\ 1H), 6.36\ (br\ s,\ 1H\times1/2), 6.20\ (br\ s,\ 1H\times1/2), 4.96\ (s,\ 1H\times1/2), 4.91\ (s,\ 1H\times1/2), 3.62\ to\ 3.42\ (m,\ 12H), 1/2 \ (s,\ 1H\times1/2), 1/2 \ (s,\ 1H\times1$

3.09 to 3.04 (m, 3H), 2.88 (s, 6H), 2.70 to 1.30 (m, 8H), 2.26 (s, 3H \times 1/2), 2.23 (s,

- 3H × 1/2). HRMS: calcd for $C_{24}H_{46}N_{5}O_{75}$ (MH⁺) 668.3112, found 668.3115. 13. *GlyTyr labeled with* **8**. HRMS: calcd for $C_{38}H_{51}N_7O_{10}S$ (MH⁺) 798.3491, found 798.3495. AlaTyrAla labeled with 8. HRMS: calcd for C42H58N8O11S (MH⁺) 883.4018, found 883.4015.
- 14. Lysozyme labeled with 8. ESI-MS: 14,865 (mono-addition product); unmodified 14306
- Myoglobin labeled with 8. ESI-MS: 17,512 (mono-addition product); unmodified 16,953. Modification of myoglobin, which lacks surface-accessible tyrosine,^{8,9b} may have occurred when the protein was denatured during the 4-day reaction time. We previously showed that myoglobin was not modified with the cyclic imine derivative shown in Scheme 1 in aqueous buffer, pH 7.0 at 37 °C for 48 h.⁷