

## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:  
<http://www.tandfonline.com/loi/gpss20>

### A Simple C<sub>2</sub> Symmetric Sulfide for a One Pot Ylide Mediated Asymmetric Conversion of Aldehydes into Epoxides

Karine Julienne , Patrick Metzner , Alfred Greiner & Vivien Henryon

<sup>a</sup> Laboratoire de Chimie Moléculaire et Thio-organique, UMR CNRS, ISMRA-Université , 6, boulevard du Maréchal Juin, 14050 Caen, France

<sup>b</sup> Laboratoire de Chimie Moléculaire et Thio-organique, UMR CNRS, ISMRA-Université , 6, boulevard du Maréchal Juin, 14050 Caen, France

<sup>c</sup> Rhône Poulenc Industrialisation, Centre de Recherches des Carrières , 85, avenue des Frères Penet, 69192 Lyon, France

<sup>d</sup> Rhône Poulenc Industrialisation, Centre de Recherches des Carrières , 85, avenue des Frères Penet, 69192 Lyon, France

Published online: 17 Mar 2008.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## A Simple $C_2$ Symmetric Sulfide for a One Pot Ylide Mediated Asymmetric Conversion of Aldehydes into Epoxides

KARINE JULIENNE<sup>a</sup>, PATRICK METZNER<sup>a\*</sup>, ALFRED GREINER<sup>b</sup>  
and VIVIEN HENRYON<sup>b</sup>


<sup>a</sup>Laboratoire de Chimie Moléculaire et Thio-organique, UMR CNRS, ISMRA-Université, 6, boulevard du Maréchal Juin, 14050 Caen, France. and

<sup>b</sup>Rhône Poulenc Industrialisation, Centre de Recherches des Carrières, 85, avenue des Frères Penet, 69192 Lyon, France

The use of sulfur ylides in their reaction with aldehydes offers an attractive synthesis of oxiranes. Though it was developed in the early 60's, asymmetric versions only appeared in the 90's.<sup>1-5</sup> A major practical advantage of this route to enantioenriched oxiranes is that they are obtained in **one** step from aldehydes and organic halides whereas the oxidation of alkenes adds one step, *i.e.* the synthesis of the alkenes by a Wittig reaction which starts from the same substrates.

Our project was to design a new chiral sulfur ylide with emphasis on **simplicity**, in terms of structure of the chiral auxiliary, preparation, and epoxidation reaction conditions.


We wish to report the efficiency of (2*R*,5*R*)-dimethylthiolane **1** for the asymmetric ylide mediated epoxidation of carbonyl compounds.<sup>6</sup> This sulfide<sup>6,7</sup> was easily synthesised in two steps and 95% yield from commercially available material.




The  $C_2$  symmetry of **1** allows the formation of a single sulfonium salt by reaction with a benzyl halide. After an extensive search of the reaction

\* E-mail: metzner@ismra.fr. Fax: (33)-2-31-45-28-77

conditions, especially the nature of the solvent, we have found an extremely simple one-pot epoxidation procedure of aldehydes to afford oxiranes **2**. The reaction was performed in *t*-BuOH/H<sub>2</sub>O 90:10, at room temperature, with a mineral base (NaOH).



As a representative example, *trans*-(2*S*,3*S*)-2,3-diphenyloxirane was obtained from benzaldehyde and benzyl bromide with **92% chemical yield**, **86% diastereomeric** and **88% enantiomeric excesses**. A range of non racemic epoxides have been synthesised using a variety of aldehydes and alkyl halides. The accompanying model is postulated to explain the stereoselectivity.



As compared to recently reported asymmetric versions of the epoxidation of aldehydes<sup>2</sup> our system has a number of advantages: i) the auxiliary has a low molecular weight and is chemically robust, ii) the reaction involves simple reagents and avoids harsh electrophiles, iii) the procedure is extremely simple to carry out and provides clean products with high ee's. Furthermore, an efficient catalytic version of the reaction (0.1 equiv. of sulfide **1**) has been shown to be possible.

Extension of this method to the synthesis of various oxiranes is in progress. Our new chiral sulfide might also be employed as an auxiliary for other reactions.

## References

- [1] A.-H. Li, L.-X. Dai, and V. K. Aggarwal, *Chem. Rev.*, **97**, 2341 (1997). V. K. Aggarwal, *Synlett*, 329 (1998). A. Solladié-Cavallo and A. Diep-Vohuule, *Tetrahedron: Asymmetry*, **7**, 1783 (1996). T. Durst, *Phosphorus, Sulfur Silicon Relat. Elem.*, **74**, 215 (1993). A.-H. Li, L.-X. Dai, X.-L. Hou, Y.-Z. Huang, and F.-W. Li, *J. Org. Chem.*, **61**, 489 (1996).
- [2] K. Julienne, P. Metzner, V. Henryon, and A. Greiner, *J. Org. Chem.*, **63**, 4532 (1998).
- [3] S. Otten, R. Fröhlich, and G. Haufe, *Tetrahedron: Asymmetry*, **9**, 189 (1998).