A NEW SYNTHESIS OF VITAMIN A

Teruaki MUKAIYAMA and Akihiko ISHIDA Department of Chemistry, Faculty of Science, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113

A new synthesis of vitamin A^{1} was successfully accomplished by reducing retinal, prepared by the reaction of 3-methyl-1-trimethylsiloxy-1,3-butadiene (III) with β -ionylideneacetaldehyde dimethyl acetal (VII) followed by the elimination of methanol with tertiary amine such as 1,5-diazabicyclo[5.4.0]undecene-5. The intermediate, β -ionylideneacetaldehyde (V), was also prepared according to the same procedure starting from β -cyclocitral dimethyl acetal (II).

In the preceding papers, a convenient method for the preparation of δ -alkoxy- α,β -unsaturated aldehydes by the reaction of various acetals with dienoxysilane in the coexistence of TiCl₄ and Ti(0ⁱPr)₄²) and a new method for the preparation of polyenals from δ -alkoxy- α,β -unsaturated aldehydes by the elimination of alcohol with tertiary amine such as 1,5-diazabicyclo[5.4.0]undecene-5 (DBU) or 1,5-diazabicyclo[4.3.0]nonene-5 (DBN) in the presence of molecular sieves 3A or 4A³) were described.

In this communication, we wish to report a new synthesis of vitamin A starting from β -cyclocitral (I) by the repetition of the above mentioned two preparative methods: β -Cyclocitral (I)⁴) was converted to its dimethyl acetal (II) on treatment with HC(OCH₃)₃-CH₃OH in the presence of d-camphorsulfonic acid at -10°C for 10 min in almost quantitative yield. Dienoxysilane (III) reacted instantaneously with II at -40°C in the presence of TiCl₄-Ti(OⁱPr)₄ to afford δ -methoxy- α , β -unsaturated aldehyde (IV) (E:Z = 3:1 mixture) in 70-80% yield.



The treatment of IV with DBU (4 equiv) in the presence of molecular sieves 3A for 5 hr in refluxing CH_2Cl_2 -CH₃CN (1:1) under an argon atmosphere afforded V and

its 2-(Z) isomer (VI) in 56% and 19% yields, respectively. The isomer VI was converted to V in 70% yield on treating with a catalytic amount of iodine $^{5)}$ in abs. $Et_2 O-C_6 H_6$ (1:1) for 4 hr at room temperature.

The synthesis of retinal (IX), a precursor of vitamin A, was also achieved according to the same procedure from V as shown in the following scheme.



Retinal (IX) was reduced with sodium borohydride in methanol at -30°C for 1 hr under an argon atmosphere to give vitamin A (XI) in quantitative yield. Acetylation of XI with acetic anhydride-pyridine under an argon atmosphere gave vitamin A acetate (XII) in 71% yield based on IX.



Spectral data of XII were consistent with those of the authentic sample obtained from commercial source; nmr (δ_{TMS} ppm, CCl₄): 1.00 (s,6H), 1.70 (s,3H), 1.85 (s,3H), 1.90 (s,3H), 1.95 (s,3H), 4.65 (d,2H), 5.55 (t,1H), 5.80 - 6.70 (5H); uv: λ_{max}^{EtOH} 326nm (ϵ 4.45 × 10³); ir: 1740, 1220 cm⁻¹; mass: m/e 328 (M⁺).

References

- O. Isler and P. Schudel, "Advances in Organic Chemistry" 4, 115, Interscience, 1) New York (1963).
- T. Mukaiyama and A. Ishida, Chem. Lett., 319 (1975).
- 3)
- A. Ishida and T. Mukaiyama, Chem. Lett., 519 (1975). L. Colombi, A. Bosshard, H. Schinz, and C. F. Seidel, Helv. Chim. Acta, <u>34</u>, 265 (1951). 4)
- H. B. Henbest, B. L. Shaw, and G. Woods, J. Chem. Soc., 1154 (1952). G. Cainell, G. Cardillo, M. Contento, P. Grasselli, and A. U. Ronchi, Gazz. 5)
- Chim. Ital., <u>103</u>, 117 (1973).

(Received September 22, 1975)