


Synthesis of the Tetrahydropyran Ring Part of a Marine Toxin Polycavernoside-A

Kenshu FUJIWARA, Seiji AMANO, Takahiro OKA, and Akio MURAI*


Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo 060

Construction of the tetrahydropyran ring part of polycavernoside A, which has been isolated as one of toxic principles from the red alga *Polycavernosa tsudai*, is described starting from (*S*)-(-)- and (*R*)-(+)-glycidols.

Continuing the previous paper¹⁾ about the synthetic studies on polycavernoside A (**1**) isolated from *Polycavernosa tsudai* by Yasumoto,²⁾ we describe herein the synthesis of **2** corresponding to the tetrahydropyran ring moiety of **1**.



The synthesis started with protection of commercially available (*S*)-(-)-glycidol³⁾ with TBSCl (Scheme 1). By a modification of the Kotsuki procedure,⁴⁾ tetrahydropyranyl ether of propargyl alcohol was initially lithiated with BuLi and treated with **3** in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ to give the coupling product, which was desilylated and then subjected to acetonide formation to yield the acetylene alcohol **4**. The compound **4** was partially reduced with Lindlar catalyst and the product was oxidized under the Sharpless conditions⁵⁾ using (+)-diethyl tartrate to give (2*S*,3*R*)-2,3-epoxy alcohol **5** in 75% yield. Compound **5** was treated successively with a mixture of MeLi (7 eq) and CuI (4 eq) in THF-ether (2.3:1)⁶⁾ and with NaIO_4 to yield an 1,3-diol **6**, along with an oxidized compound **6**, in 69% and 20% overall yields, respectively. The compound **6** was led to the corresponding monobenzyl ether **7** in a three-step process (TBSCl; BnBr; TBAF). The hydroxyl group of **7** was oxidized under the Swern condition to afford the aldehyde, which was immediately allowed to react with $(\text{CH}_3\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CO}_2\text{CH}_3$, KH, and 18-crown-6 in THF^{6,7)} to afford a separable 5:1 mixture of (*Z*)- and (*E*)-olefin esters. The (*Z*)-olefin ester was isolated and deprotected with PTS to give a diol **8** in 92% yield.




Reagents: a) TBSCl, Imid., DMF, 0 °C, 1 h, 96%; b) propargyl alcohol THP-ether, BuLi, BF₃·OEt₂, THF, -78 °C, 140 min; PTS·H₂O, MeOH, reflux, 9 h; dimethoxypropane, PTS·H₂O, CH₂Cl₂-MeOH (1:4), 20 °C, 25 min, 71%; c) H₂, Lindlar cat., quinoline, EtOH, 20 °C, 3 h, 91%; (+)-DET (1.0 eq), Ti(O^{*i*}Pr)₄ (1.0 eq), TBHP (2.2 eq), MS4A, CH₂Cl₂, -20 °C, 3 days, 75%; d) MeLi (7 eq), CuI (4 eq), THF-ether (2.3:1), -20 °C, 23 h; NaIO₄, THF-H₂O (2:1), 20 °C, 26 min, **6** (69%), **6'** (20%); e) TBSCl, DMAP, NEt₃, CH₂Cl₂, 0 °C, 1.5 h, 91%; *t*-BuOK, BnBr, TBAI, THF, 0 °C, 47 min, 95%; TBAF, THF, 20 °C, 80 min, 79%; f) Swern oxid.; (MeO)₂POCH₂CO₂Me, KH, 18-crown-6, THF, 0 °C, 1 h, then aldehyde, -78 °C, 225 min, (Z)-olefin (78%), (E)-olefin (15%); PTS·H₂O, MeOH, 28 °C, 1.5 h, 92%; g) *t*-BuOK (8 mol%), THF, -20 °C, 20 min, 92% (**2**:**9**=6.7:1); h) *t*-BuOK (1 eq), MeOH, 25 °C, 11 h, 91% from **8**.

Scheme 1.

In order to optimize the reaction conditions for the next intramolecular Michael cyclization of the olefin diol **8**, the enantiomeric (*E*)-isomer **10** was subjected to the reaction under the various conditions (Table 1). First of all, *t*-BuOK (1.5 eq) was employed in THF at rather low temp (Run 1).⁸⁾ Although the ester part was predominantly hydrolyzed, the desired cyclized compound **11** was obtained in totally 45% yield. It is to be noted that in this case the isomeric cyclized product **12** was not detected in even a trace amount. Next, use of MeONa as a rather weak base led to formation of an 1:1 mixture of **11** and **12** (Run 2). When a catalytic

Table 1.




Run	Conditions	Yield of 11 ^{a)}	Yield of 12 ^{a)}
1	10 , <i>t</i> -BuOK (1.5 eq), THF -20 °C, 12 min → 20 °C, 75 min	40% (R=H), ^{b)} 5% (R=CH ₂ CH ₃)	—
2	10 , NaH (1.5 eq), MeOH 20 °C, 15 h → 40 °C, 9.5 h	42% (R=CH ₃)	42% (R=CH ₃)
3	10 , <i>t</i> -BuOK (0.1 eq), THF -20 °C, 23 min → 20 °C, 14 min	34% (R=CH ₂ CH ₃) ^{c)}	51% (R=CH ₂ CH ₃) ^{c)}
4	12 , <i>t</i> -BuOK (1.5 eq), THF -20 °C, 15 min → 20 °C, 1 h	39% (R=H) ^{b)}	13% (R=H) ^{b)}

a) The respective yields of **11** and **12** were estimated by ¹H NMR spectra; b) The compound was isolated as its methyl ester on treatment with CH₂N₂ in ether; c) The compound was isolated as its methyl ester on reaction with *t*-BuOK (1.5 eq) in MeOH.

amount of *t*-BuOK was used, the relative yields of **11** and **12** amounted to 34% and 51%, respectively (Run 3). These results reveal that the comparable ratio of **11** and **12** would be formed kinetically followed by gradual decomposition of the isomer **12** with an excess amount of *t*-BuOK. In fact, treatment of the less stable **12** with *t*-BuOK (1.5 eq) in THF effected equilibration, leading to the 3:1 ratio of **11** and **12** with only a total 52% recovery (Run 4). Eventually, (*E*)-olefin alcohol **10** gave rise to only a limited amount of the desired product **11**. Accordingly, we attempted to check the reaction with the corresponding (*Z*)-olefin diol **8** (Scheme 1). Compound **8** was treated with *t*-BuOK (8 mol%) in THF at -20 °C for 20 min, the intramolecular Michael addition reaction was effected to give a 6.7:1 mixture of the desired cyclized product **2** and its dimer **9** in 92% yield.⁹⁾ Compound **9** was smoothly converted to **2** with *t*-BuOK (1 eq) and the overall yield of **2** amounted to 91% from **8**. The exclusive formation of **2** could be rationalized in terms of the transition state closing to the cyclization (Scheme 2). The transition state **8A** leading to **2** might be considerably favored comparing to another rotated isomer **8B** leading to the plausible isomer **13** on the basis of steric repulsion between methoxycarbonyl group and the hydrogen at C-5 position.¹⁰⁾

The enantioisomer corresponding to **2** has been also synthesized starting from (*R*)-(+)-glycidol.^{3,11)} The synthetic study on the respective connection of the tetrahydropyran **2** and its enantiomer with the isomers of the tetrahydrofuran moieties¹⁾ is now in progress in our laboratories.



Scheme 2.

The authors thank financial support by the Grant-in Aid for Scientific Research on Priority Areas No. 06225201 from the Ministry of Education, Science and Culture, Japan.

References

- 1) N. Hayashi, T. Mine, K. Fujiwara, and A. Murai, the previous paper.
- 2) M. Y.-Yamashita, R. L. Haddock, and T. Yasumoto, *J. Am. Chem. Soc.*, **115**, 1147 (1993).
- 3) The reagent was kindly provided by DAISO Co., Ltd.
- 4) H. Kotsuki, Y. Ushio, I. Kadota, and M. Ochi, *J. Org. Chem.*, **54**, 5153 (1989).
- 5) Y. Gao, R. M. Hanson, J. M. Klunder, S. Y. Ko, H. Masamune, and K. B. Sharpless, *J. Am. Chem. Soc.*, **109**, 5765 (1987).
- 6) Cf., H. Nagaoka and Y. Kishi, *Tetrahedron*, **37**, 3873 (1981).
- 7) M. Miyashita, K. Yoshihara, K. Kawamine, M. Hoshino, and H. Irie, *Tetrahedron Lett.*, **34**, 6285 (1993).
- 8) Cf., K. Mori, N. Hikage, A. Inagaki, S. Mori, K. Nomura, and E. Yoshii, *J. Org. Chem.*, **57**, 2888 (1992).
- 9) **2**; colorless needles, mp 88-89 °C, $[\alpha]_D^{23} -50.9^\circ$ (c 1.01, CHCl₃).
- 10) For the intramolecular cyclization leading to tetrahydropyran derivatives from the linear starting compounds having ω-hydroxy-α,β-unsaturated carboxylic esters, see B. Fraser-Reid, R. D. Dawe, and D. B. Tulshian, *Can. J. Chem.*, **57**, 1746 (1979); R. D. Dawe and B. Fraser-Reid, *J. Org. Chem.*, **49**, 522 (1984); V. S. Martin, M. T. Nunez, M. A. Ramirez, and M. A. Soler, *Tetrahedron Lett.*, **31**, 763 (1990); T. D. Aicher, K. R. Buszek, F. G. Fang, C. J. Forsyth, S. H. Jung, Y. Kishi, and P. M. Scola, *ibid.*, **33**, 1549 (1992); V. S. Martin and J. M. Palazon, *ibid.*, **33**, 2399 (1992); J. M. Palazon, M. A. Soler, M. A. Ramirez, and V. S. Martin, *ibid.*, **34**, 5467 (1993).
- 11) Enantiomer of **2**; colorless needles, mp 89-90 °C, $[\alpha]_D^{24} +51.0^\circ$ (c 1.01, CHCl₃).

(Received August 22, 1994)