This article was downloaded by: [Stanford University Libraries] On: 19 September 2012, At: 03:34 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lsyc20

An Improved Preparation Method of Benzyl and Thenyl Triphenylphosphonium Salts

Jian-Xin Zhang $^{\rm a}$, Philippe Dubois $^{\rm a}$ & Robert Jérôme $^{\rm a}$

^a Center for Education and Research on Macromolecules (CERM), University of Liège, Sart-Tilman(B6), 4000, Liège, BELGIUM

Version of record first published: 21 Aug 2006.

To cite this article: Jian-Xin Zhang, Philippe Dubois & Robert Jérôme (1996): An Improved Preparation Method of Benzyl and Thenyl Triphenylphosphonium Salts, Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry, 26:16, 3091-3095

To link to this article: http://dx.doi.org/10.1080/00397919608004613

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

AN IMPROVED PREPARATION METHOD OF BENZYL AND THENYL TRIPHENYLPHOSPHONIUM SALTS

Jian-Xin Zhang, Philippe Dubois*, and Robert Jérôme

Center for Education and Research on Macromolecules (CERM), University of Liège, Sart-Tilman(B6), 4000 Liège BELGIUM

<u>Abstract</u>: New benzyl and thenyl phosphonium salts have been prepared by an improved method in solution from the parent alcohols and triphenylphosphine hydrobromide. Benzyl-type alcohols containing a tertiary amine can also be converted within high yields to phosphonium salts by removing the formed water by azeotropic distillation. Solvent (methylene chloride, chloroform and acetonitrile) and temperature have been chosen in close relation to the alcohol reactivity. Purification is straightforward and yields are relatively high.

The Wittig reaction is a well known and very efficient method for the incorporation of a carbon-carbon double bond in an organic molecule¹. This opportunity is largely exploited in the field of organic conjugated materials²⁻⁴. The basic reagent in Wittig reaction, i.e. alkyl triphenylphosphonium salt, is usually prepared from the parent halide by reaction with triphenylphosphine. Although the quaternization reaction of triphenylphosphine usually works well, its application is limited to the halides that are stable enough to be properly worked up. For instance, 4-dimethylaminobenzyl triphenylphosphonium chloride cannot be prepared by using the traditional method.

^{*}to whom correspondence should be addressed

Copyright © 1996 by Marcel Dekker, Inc.

This paper will report on the improvement of the Hamanaka's method⁵ in order to prepare phosphonium salts of various aromatic compounds by reacting benzyl or thenyl alcohols with triphenylphosphonium hydrobromide (Scheme 1). A special attention will be paid to aromatic derivatives substituted by a tertiary amine, for which the synthesis of the phosphonium salts has never been reported, at least at our best knowledge.

2-Thiophenemethanol 1a has been reacted with 2 in refluxing methylene chloride for 4 hours. As shown in Table 1, 52% of 2-thenyl triphenylphosphonium bromide 3a have been recovered by precipitation in diethyl ether.



Scheme 1

 Table 1. Synthesis of Phosphonium Bromides 3 from the Parent Alcohols 1 and

 Triphenylphosphine Hydrobromide 2 with Removing the Formed Water by

Entry ⁽¹⁾	Phosphonium bromide	Solvent	Refluxing time (hr)	Yield ⁽²⁾ (%)
a	3 a	CH ₂ Cl ₂	4	52 ⁽³⁾
	1	CH_2Cl_2	1	99
		CHCl ₃	1	100
b	3b	CH ₂ Cl ₂	1	96
с	3c	CH ₂ Cl ₂	1	97
		CHCl ₃	1	100
d	3d	CHCl ₃	2	100
e	3e	CH ₂ Cl ₂ - CH ₃ CN	1	82
		CH ₃ CN	2.5	81
f	3f	CH ₂ Cl ₂	6	(4)
		CH ₃ CN	2	82
g	3g + (5)	CH ₃ CN	1	97 ⁽⁶⁾
h	3h	CH ₃ CN	2	100
i	3i	CH ₃ CN	1	91
j	3j	CH ₃ CN	1	99

(1) see Scheme 1; (2) yield of the isolated product; (3) without solvent distillation; (4) no reaction; (5) bis-substituted phosphonium; (6) yield calculated for mono-substitution.

In order to shift the reaction equilibrium toward formation of the expected product, the azeotropic distillation of the solvent has been carried out all along the reaction and very high yields, even quantitative in some cases, have been observed, as shown in Table 1. Various solvents, i.e. methylene chloride, chloroform and acetonitrile, have been

selected for their ability to dissolve the final phosphonium salt and to be distilled off in the required reaction temperature range. It is worth pointing out that the removal of water either by azeotropic distillation or with drying agents has been proposed by Pommer et al. for the preparation of phosphonium salts from alcohols, e.g. β -ionol, and triphenyl-phosphine hydrohalides⁶. However only reactions involving non functionalized allylic alcohols were reported.

Benzyl and thenyl phosphonium salts **3c-f** have also been successfully recovered within high yields and purity even tough a clear difference in the reaction yields is observed when benzyl alcohols and thenyl alcohols in the series of dimethylamino compounds are compared (Entry c to f in Table 1), which might reflect the higher reactivity of benzyl alcohols compared to thenyl analogues.

General Procedure to Prepare Benzyl and Thenyl Phosphonium Bromide 3:

Triphenylphosphine hydrobromide 2 was prepared according to the procedure reported by Hercouet⁷. In a flask equipped with a cooler, 0.1mole of alcohol and 0.09mole of **2** were added to ca. 100 ml of a suitable solvent (see Table 1) under magnetic stirring. After refluxing for the desired period of time, the cooler was replaced by a distillation equipment. The solvent was distilled off slowly until dryness for ca. 1 hour. The crude product was then solubilized in chloroform, and the pure phosphonium bromide was recovered by precipitation in diethyl ether, washed two times with ether, and isolated by filtration. ACKNOWLEDGMENT: The authors are indebted to the "Ministère de la Région Wallonne" for a research grant (Matériaux nouveaux à architecture intelligente) and the "Fonds National de la Recherche Scientifique (FNRS)" for financial support. Ph. D. is Research Associate of the Belgian FNRS. They also thank the "Services des Affaires Scientifiques Techniques et Culturelles" (Pôles d'Attraction Interuniversitaires : Polymères).

REFERENCES AND NOTES:

- 1. Maryanoff, B. and Reitz, A. Chem. Rev. 1989, 89, 863-927;
- 2. Koβmehl, G. and Hoppe, F. D. Liq. Cryst. 1993, 15, 383-393;
- 3. Effenberger, F.; Meller, P.; Ringsdorf, H. and Schlosser, H. Adv. Mater. 1991, 3, 555-558;
- S'heeren, G.; Persoons, A.; Rondou, P.; Wiersma, J.; van Beylen, M. and Samyn, C. Makromol. Chem, 1994, 195, 1733-1744;
- 5. Hamanaka, N.; Kosuge, S.; and Iguchi, S. Synlett 1991, 139-140;
- Sarnecki, W. and Pommer H. Ger. Pat. 1,060,386 (to BASF) [C.A., 1961, 55, 4577b]; see also Organic Reactions, vol. 14, p. 392;
- 7. Hercouet, A. and Le Corre, M. Synthesis 1988, 157-158;

(Received in The Netherlands 01 March 1996)