

Efficient and Stable Photocatalytic Systems for Reductive
Dechlorination of *p*-Chlorobiphenyl by Sodium Borohydride

Masashi ISHIKAWA[†] and Shunichi FUKUZUMI*

Department of Applied Chemistry, Faculty of Engineering,
Osaka University, Suita, Osaka 565

Acridine Derivatives (9,10-dihydro-10-methylacridine and acriflavine) act as efficient and stable photocatalysts for reductive dechlorination of *p*-chlorobiphenyl as well as dehalogenation of bromochlorobenzenes with sodium borohydride in acetonitrile/H₂O (9:1 v/v) at 298 K. In the case of acriflavine, the limiting quantum yield at irradiation wavelength $\lambda = 350$ nm has reached 0.63.

Photochemical reductive dechlorination of chlorinated compounds has recently been studied extensively, in part due to their role as environmental pollutants.¹⁻⁷⁾ Although several photocatalytic systems for reductive dechlorination have so far been reported,^{3,4)} it is desired to develop much more efficient and stable photocatalysts which have sufficient spectral overlap with the solar spectrum.

This study reports efficient and stable photocatalytic systems for reductive dechlorination of *p*-chlorobiphenyl, which is known to be most difficult to be reduced among PCBs,⁷⁾ as well as dehalogenation of bromochlorobenzenes using sodium borohydride (NaBH₄) and acridine derivatives as a reductant and photocatalysts, respectively.

No appreciable photoreduction of *p*-chlorobiphenyl (ClBP) by NaBH₄ occurs in the absence of photocatalyst in a mixture of acetonitrile and H₂O (MeCN/H₂O, 9:1 v/v) under the irradiation of light from a Xenon lamp as shown in Fig. 1 (part a).⁸⁾ When 9,10-dihydro-10-methylacridine (AcrH₂) or acriflavine (AFH⁺) is added to this system at 298 K, each species acts as an efficient photocatalyst for reductive dechlorination of *p*-chlorobiphenyl with NaBH₄ to yield biphenyl as shown in Fig. 1 (part b or c), where the yield of biphenyl was determined by glc. No appreciable photo-

[†]Present address: Laboratory of Organic Chemistry, Osaka Women's University, Sakai, Osaka 590.

degradation of the catalysts was observed during the photocatalytic reaction, since the concentration of AcrH_2 remained unchanged. When AcrH_2 is replaced by the oxidized form, 10-methylacridinium perchlorate ($\text{AcrH}^+\text{ClO}_4^-$), essentially the same result is obtained, since AcrH^+ is immediately reduced by NaBH_4 to yield AcrH_2 selectively.⁹⁾ Thus, $\text{AcrH}_2/\text{AcrH}^+$ redox pair acts as an efficient and stable photocatalyst for dechlorination of ClBP with NaBH_4 . Efficient dehalogenation of *o*-, *m*-, *p*-bromochlorobenzenes also occurs by using the present photocatalytic system.¹⁰⁾

Since the one-electron oxidation potential (E_{ox}^0) of the singlet excited state

$^1\text{AcrH}_2^*$ is known to be largely negative ($E_{\text{ox}}^0 = -3.1 \text{ V vs. SCE}$),¹¹⁾ $^1\text{AcrH}_2^*$ may act as a very strong reductant. In fact, the fluorescence of $^1\text{AcrH}_2^*$ is readily quenched by ClBP. From the Stern-Volmer plot is obtained the Stern-Volmer constant (K_{SV}) as $1.2 \times 10^2 \text{ dm}^3 \text{ mol}^{-1}$. The fluorescence lifetime (τ) of $^1\text{AcrH}_2^*$ in $\text{MeCN}/\text{H}_2\text{O}$ (9:1 v/v) was determined as 7.0 ns by using the single photon counting technique. Thus, the quenching rate constant ($k_q = K_{\text{SV}}\tau^{-1}$) is obtained as $1.7 \times 10^{10} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ which is close to the diffusion rate constant.¹⁰⁾ The K_{SV} value for the fluorescence quenching of the reduced form of acriflavine ($^1\text{AFH}_2^*$) by ClBP is also obtained as $3.6 \times 10^1 \text{ dm}^3 \text{ mol}^{-1}$ which is smaller than the K_{SV} value of $^1\text{AcrH}_2^*$ ($1.2 \times 10^2 \text{ dm}^3 \text{ mol}^{-1}$).

The quantum yields (Φ) of the photocatalytic dechlorination of ClBP were determined using a ferrioxalate actinometer. The Φ values in the presence of AcrH_2 under irradiation of light ($\lambda = 320 \text{ nm}$) were constant with the change of both AcrH_2 and NaBH_4 concentrations. On the other hand, the quantum yield increased with an increase in the concentration of ClBP





Fig. 1. Photodechlorination of *p*-chlorobiphenyl (0.10 mol dm^{-3}) with NaBH_4 (1.0 mol dm^{-3}) in $\text{MeCN}/\text{H}_2\text{O}$ (9:1 v/v) (a) in the absence of catalyst (Δ), (b) in the presence of AcrH_2 ($2.0 \times 10^{-2} \text{ mol dm}^{-3}$) ($\circ$) and (c) AFH^+ ($2.0 \times 10^{-2} \text{ mol dm}^{-3}$) ($\bullet$) under the irradiation of light from a Xenon lamp. Plots of yield of biphenyl vs. irradiation time.


to approach a limiting value (Φ_{∞}) in the high concentrations in accordance with Eq. 1. Then, from the linear plot between Φ^{-1} and $[\text{ClBP}]^{-1}$ the

$$\Phi^{-1} = \Phi_{\infty}^{-1} [1 + (K_{\text{Obs}}[\text{ClBP}])^{-1}] \quad (1)$$

Φ_{∞} and K_{Obs} values are obtained as 0.42 and $1.6 \times 10^2 \text{ dm}^3 \text{ mol}^{-1}$, respectively. In the absence of NaBH_4 as well, *p*-chlorobiphenyl was reduced by AcrH_2 in $\text{MeCN}/\text{H}_2\text{O}$ (9:1 v/v) under irradiation of light ($\lambda = 320 \text{ nm}$) to yield AcrH^+ and biphenyl, Eq. 2. The Φ values in the absence of NaBH_4



showed the same dependence on the concentration of ClBP as Eq. 1, and the Φ_{∞} and K_{Obs} values were determined as 0.61 and $1.2 \times 10^2 \text{ dm}^3 \text{ mol}^{-1}$, respectively, which agree reasonably well as those obtained in the presence of NaBH_4 (0.42 and $1.6 \times 10^2 \text{ dm}^3 \text{ mol}^{-1}$, respectively).¹¹⁾ When AcrH_2 is replaced by AFH^+ in the photocatalytic reductive dechlorination of ClBP with NaBH_4 , the limiting quantum yield ($\Phi_{\infty} = 0.63$) becomes larger than the case of AcrH_2 ($\Phi_{\infty} = 0.42$) under irradiation of light of longer wavelength ($\lambda = 350 \text{ nm}$), which has the sufficient overlap with the solar spectrum for the practical use. The K_{Obs} values obtained from the Φ dependence on the concentration of ClBP (Eq. 1) in the $\text{AcrH}_2/\text{AcrH}^+$ - and $\text{AFH}_2/\text{AFH}^+$ -catalyzed photo-dechlorination with NaBH_4 ($K_{\text{Obs}} = 1.6 \times 10^2$ and $3.3 \times 10^1 \text{ dm}^3 \text{ mol}^{-1}$, respectively) agree well with the K_{SV} values of $^1\text{AcrH}_2^*$ and $^1\text{AFH}_2^*$ ($K_{\text{SV}} = 1.2 \times 10^2$ and $3.6 \times 10 \text{ dm}^3 \text{ mol}^{-1}$, respectively). Thus, the photocatalytic dechlorination of *p*-chlorobiphenyl by NaBH_4 may proceed via the reductive dechlorination of ClBP by the singlet excited states $^1\text{AcrH}_2^*$ (or $^1\text{AFH}_2^*$), followed by the thermal reduction of AcrH^+ (or AFH^+) by NaBH_4 to regenerate AcrH_2 (or AFH_2) as shown below.



References

- 1) "PCB Poisoning and Pollution," ed by K. Higuchi, Academic Press, Tokyo (1976); D. G. Ackerman, L. L. Scinto, P. S. Bakshi, R. G. Delumyea, R. J. Johnson, G. Richard, A. M. Takata, and E. M. Sworzyn, "Destruction and Disposal of PCBs by Thermal and Non-Thermal Methods," Noyes, Park Ridge (1983); I. Mori, Kankyo Gijutsu, 14, 588 (1985).
- 2) G. A. Epling, W. M. McVicar, and A. Kumar, Chemosphere, 17, 1355 (1988).
- 3) M. Ohashi, K. Tsujimoto, and K. Seki, J. Chem. Soc., Chem. Commun., 1973, 384; K. Tsujimoto, S. Tasaka, and M. Ohashi, *ibid.*, 1975, 758; M. Ohashi and K. Tsujimoto, Chem. Lett., 1983, 423; Y. Tanaka, T. Uryu, M. Ohashi, and K. Tsujimoto, J. Chem. Soc., Chem. Commun., 1987, 1703.
- 4) J. Ph. Soumillion, P. Vandereecken, and F. C. De Schryver, Tetrahedron Lett., 30, 697 (1989).
- 5) N. J. Bunce, J. P. Landers, J. A. Langshaw, and J. S. Nakai, Environ. Sci. Technol., 23, 213 (1989); N. J. Bunce, C. T. DeSchutter, and E. J. Toone, J. Chem. Soc., Perkin Trans. 2, 1983, 859.
- 6) P. K. Freeman, R. Srinivasa, J.-A. Campbell, and M. L. Deinzer, M. L. J. Am. Chem. Soc., 108, 5531 (1986).
- 7) G. A. Epling and E. Florio, J. Chem. Soc., Chem. Commun., 1986, 185; G. A. Epling and E. Florio, Tetrahedron Lett., 27, 675 (1986).
- 8) Typically, an MeCN/H₂O (9:1 v/v) solution (0.50 cm³) containing ClBP (0.10 mol dm⁻³) and NaBH₄ (1.0 mol dm⁻³) was added to a quartz cuvette (1 mm i.d) and deoxygenated by a stream of argon. The photolysis with a Ushio model U1-501 Xenon lamp results in no appreciable reaction under our experimental conditions, although photolysis of ClBP and NaBH₄ with a low pressure mercury lamp is reported to result in the dechlorination of ClBP.⁷⁾
- 9) R. M. G. Roberts, D. Ostovic, and M. M. Kreevoy, Faraday Discuss. Chem. Soc., 74, 257 (1982).
- 10) The reductive debromination of bromochlorobenzenes proceeds more efficiently than the dechlorination, since chlorobenzene is obtained as a major product; e.g., the product ratio of PhCl to PhBr was 8:1 in the case of *o*-BrC₆H₄Cl.
- 11) S. Fukuzumi and T. Tanaka, "Photoinduced Electron Transfer," ed by M. A. Fox and M. Chanon, Elsevier, Amsterdam (1988), Part C, Chap. 10.
- 12) The limiting quantum yields in the presence of NaBH₄ may be larger than 0.42, since the photolysis in the presence of NaBH₄ results in the formation of H₂ which bubbles up from the reactant solution, causing the slight decrease in the light absorption.

(Received April 9, 1990)