

Preparation, Spectroscopic and Structural Characterization of η^1 -N and η^2 -Se,Se' Complexes of a $P_2N_4Se_2$ Ring

Tristram Chivers,^{*a} Daniel D. Doxsee,^a Robert W. Hiltz,^a Auke Meetsma,^b Masood Parvez^a and Johan C. van de Grampel^{*b}

^a Department of Chemistry, The University of Calgary, Calgary, Alberta T2N 1N4, Canada

^b Department of Polymer Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands


The reaction of 1,5- $Ph_4P_2N_4Se_2$, **1**, prepared by the treatment of $Ph_2PN_2(SiMe_3)_3$ with a mixture of $SeCl_4$ and Se_2Cl_2 in acetonitrile, with $[PtCl_2(PEt_3)]_2$ gives the adducts $[PtCl_2(PEt_3)]_n[\eta^1\text{-N-}Ph_4P_2N_4Se_2]$ (**2**, $n = 1$; **3**, $n = 2$) in which the $P_2N_4Se_2$ ring is shown by an X-ray structural determination of **3** to contain a transannular Se–Se interaction; the oxidative addition of **1** or **2** to $Pt(C_2H_4)(PPh_3)_2$ produces the corresponding η^2 -Se,Se' complexes.

Several interesting reactions of Se_4N_4 have been reported recently.^{1–5} Without exception these reactions result in breakdown of the heterocyclic ring. For example, the treatment of Se_4N_4 with $[PtCl_2(PMe_2Ph)]_2$ or $Pt(PPh_3)_3$ produces the complexes $Pt(Se_3N)Cl(PMe_2Ph)_4$ or $[Pt(Se_2N_2)(PPh_3)]_2$,⁵ which contain the chelating ligands Se_3N^- and $Se_2N_2^{2-}$, respectively. Furthermore, Se_4N_4 is difficult to handle owing to its explosive nature and insolubility in organic solvents. By contrast, the 1,5-diphosphadiselenatetrazocine **1** is non-explosive, soluble in organic solvents, and contains a structurally informative ^{31}P NMR probe.⁶ By analogy with the chemistry of the sulfur analogue,^{7,8} the hybrid inorganic heterocycle **1** should provide a good model for investigating reactions of unsaturated selenium–nitrogen systems. However, **1** has only been obtained in very low yields and the existence of the cross-ring bond has not been unequivocally established owing


to the tendency of this compound to deposit red selenium in the solid state or in solution.⁶

We report here a high yield synthesis of **1** and the reaction of **1** with $[PtCl_2(PEt_3)]_2$ to give the *N*-bonded 1:1 and 1:2 adducts, **2** and **3**. We also establish the presence of a transannular Se–Se interaction in **3** by X-ray crystallography. The oxidative addition of $Pt(PPh_3)_2(C_2H_4)$ to **1** or **2** produces the corresponding η^2 -Se,Se' adducts **4** and **5**, respectively.

The eight-membered ring **1** is readily obtained by the cyclocondensation reaction between $Ph_2PN_2(SiMe_3)_3$ and a mixture of selenium halides designed to imitate 'SeCl₃' [eqn. (1)]. The slow addition of $Ph_2PN_2(SiMe_3)_3$ (21.5 mmol) in acetonitrile (100 ml) to a mixture of $SeCl_4$ (14.3 mmol) and Se_2Cl_2 (3.6 mmol) suspended in acetonitrile (150 ml) at 23 °C produced (after 16 h) a yellow–brown precipitate and a yellow solution. The volume of the solution was reduced to 50 ml and



the precipitate was dried under vacuum. Extraction of this precipitate with dichloromethane (3×40 ml) gave **1** in 85% yield (9.1 mmol) as a yellow solid identified by ^{31}P and ^{77}Se NMR spectra.^{6†}



The addition of **1** to the dimer $[\text{PtCl}_2(\text{PEt}_3)]_2$ in a 2 : 1 molar ratio in CH_2Cl_2 at -78°C , followed by warming to room temp. for 20 min, produces a mixture of the 1 : 1 adduct, **2**, and the 2 : 1 adduct, **3**, in the approximate molar ratio 7 : 2 as determined by ^{31}P NMR spectroscopy. A pure sample of the monoadduct was obtained by layering hexane on to a solution of the mixture of **2** and **3** in dichloromethane.[†] The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **2** shows three resonances at δ 2.4 (d, $^3J_{\text{P}_\text{A}-\text{P}_\text{B}}$ 5, $^1J_{\text{P}_\text{A}-\text{Pt}}$ 3619 Hz), 109.3 (d, $^4J_{\text{P}_\text{A}-\text{P}_\text{B}}$ 16, $^2J_{\text{P}_\text{B}-\text{Se}_\text{A}}$ 93 Hz) and 114.3 (dd, $^4J_{\text{P}_\text{A}-\text{P}_\text{B}}$ 16, $^3J_{\text{P}_\text{A}-\text{P}_\text{X}}$ 5, $^2J_{\text{P}_\text{A}-\text{Pt}}$ 94 Hz). The low-frequency resonance is readily assigned to the Et_3P ligand (P_X) on the basis of the large P–Pt coupling constant, whereas the two high-frequency resonances can be attributed to the inequivalent phosphorus atoms, P_A and P_B of the heterocyclic ring, [cf. $\delta(^{31}\text{P})$ 113.4 for **1**].⁶ The ^{77}Se NMR spectrum of **2** shows two resonances for the inequivalent Se atoms at δ 1270 ($^2J_{\text{Se}_\text{A}-\text{P}_\text{B}}$ 93, $^2J_{\text{Se}_\text{A}-\text{P}_\text{A}}$ 85 Hz) and 1258 ($^2J_{\text{Se}_\text{B}-\text{P}}$ 96, $^2J_{\text{Se}_\text{B}-\text{P}}$ 60 Hz) (ref. SeMe_2 δ 0), which appear as a pseudotriplet and a doublet of doublets, respectively.

The diadduct **3** can be obtained in virtually quantitative yield by carrying out the reaction of **1** with $[\text{PtCl}_2(\text{PEt}_3)]_2$ in a 1 : 1 molar ratio.[†] The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **3** exhibits two singlets at δ 3.0 ($^1J_{\text{P}-\text{Pt}}$ 3654 Hz) and 113.2 ($^2J_{\text{P}-\text{Se}}$ 69 and 89 Hz) attributable to the Et_3P ligands and equivalent PPh_2 groups, respectively. The ^{77}Se NMR spectrum of **3** exhibits a five line AA'X pattern (where X is ^{77}Se for the most abundant isotopomer) centred at δ 1337 with $|J_{\text{P}_\text{A}-\text{Se}} + J_{\text{P}_\text{A}'-\text{Se}}|$ 160 Hz. These NMR data are uniquely consistent with a distal arrangement for the $\text{PtCl}_2(\text{PEt}_3)$ groups in the 2 : 1 adduct. Furthermore, the anomalously high-frequency ^{31}P NMR chemical shifts of the PPh_2 groups observed for **2** and **3** suggests strongly that N-coordination does not disrupt the Se–Se transannular interaction in these adducts.^{9,10}

The structure of **3** was established by X-ray crystallography. Yellow crystals of **3** were obtained by recrystallization from




Fig. 1 ORTEP plot (50% probability ellipsoids) for $[\eta^1\text{-PtCl}_2(\text{PEt}_3)]_2(1,5\text{-Ph}_4\text{P}_2\text{N}_4\text{Se}_2)$, **3**; selected bond lengths (Å) and bond angles ($^\circ$): Pt(1)–P(1) 2.229(3), Pt(1)–Cl(1) 2.319(3), Pt(1)–Cl(2) 2.301(3), Pt(1)–N(1) 2.162(9), Se(1)–Se(1)* 2.594(2), Se(1)–N(1) 1.753(9), Se(1)–N(2)* 1.717(8), P(2)–N(1) 1.673(9), P(2)–N(2) 1.64(1), Cl(2)–Pt(1)–P(1) 91.5(1), Cl(2)–Pt(1)–N(1) 87.5(3), Cl(1)–Pt(1)–P(1) 94.2(1), Cl(1)–Pt(1)–N(1) 86.9(3), Se(1)*–Se(1)–N(1) 91.7(3), Se(1)–Se(1)*–N(2) 89.9(3), N(1)–Se(1)–N(2)* 106.0(4), N(1)–P(2)–N(2) 110.9(4), Pt(1)–N(1)–Se(1) 108.7(4), Pt(1)–N(1)–P(2) 117.2(5), Se(1)–N(1)–P(2) 118.1(6), Se(1)*–N(2)–P(2) 122.4(5). The asterisks indicate symmetry-related atoms.

CH_2Cl_2 –hexane at 0°C .[‡] The molecular geometry and atomic numbering scheme are shown in Fig. 1. The structural determination confirms that the equivalent $\text{PtCl}_2(\text{PEt}_3)$ groups are $\eta^1\text{-N}$ bonded to the $\text{P}_2\text{N}_4\text{Se}_2$ ring in distal positions. The most interesting feature of the structure of **3** is the transannular Se–Se interaction. The Se–Se distance is 2.594(2) Å (cf. 2.34 Å for the Se–Se single bond in Se_8).¹¹ Based on the observation that the formation of N-bonded adducts results in a small contraction of the S–S distance for the sulfur analogue of **1**,^{9,10} we tentatively estimate a value of 2.65–2.70 Å for the Se–Se interaction in **1**. This can be compared with the values of 2.748(9) and 2.84(1) Å found for the analogous intramolecular $\pi^*-\pi^*$ interactions¹² in S_4N_4 ¹³ and Se_8^{2+} ,¹⁴ respectively. The Se–N and P–N bonds involving the coordinated nitrogen atom N(1) in **3** are slightly longer than the corresponding bonds to N(2). The Se(1)–N(2) distance of 1.717(8) Å is significantly shorter than the average value of 1.777(3) Å observed for $d(\text{Se}^{\text{IV}}-\text{N})$ in 1,5- $\text{Ph}_4\text{P}_2\text{N}_4\text{Se}_2\text{Me}_2$ ⁶ (cf. 1.64 Å predicted for $\text{Se}^{\text{IV}}=\text{N}$ double bond).¹³ The coordinated nitrogen atom N(1) is removed significantly from the Se_2N_2 plane as indicated by the dihedral angles N(1)–Se(1)–Se(1)*–N(1)* = 129.0(6), N(2)*–Se(1)–Se(1)–N(2)* = –83.0(6), and N(1)–Se(1)–Se(1)*–N(2)* = 23.0(4) $^\circ$. The sum of the angles at N(1) is 344 $^\circ$. The geometry at the platinum(II) centres is approximately square planar with bond angles in the range 86.9(3)–94.2(1) $^\circ$.

The eight-membered ring **1** readily undergoes oxidative-addition with $\text{Pt}(\text{C}_2\text{H}_4)(\text{PPh}_3)_2$ at 0°C in toluene to give **4** in 70% yield.[†] The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **4** consists of two singlets (with ^{195}Pt and ^{77}Se satellites) at δ 17.6 ($^1J_{\text{P}-\text{Pt}}$ 2980,

[‡] Crystal data for $\text{C}_{36}\text{H}_{50}\text{Cl}_4\text{P}_4\text{N}_4\text{Se}_2\text{Pt}_2$ **3**: $M = 1352.62$, monoclinic, space group $\text{C}2/c$; at 150(1) K, $a = 25.150(10)$, $b = 12.090(3)$, $c = 14.597(5)$ Å, $\beta = 96.73(4)^\circ$, $V = 4407(3)$ Å³, $Z = 4$, $\mu(\text{Mo-K}\alpha) = 8.464 \text{ mm}^{-1}$, $F(000) = 2584$, $D_c = 2.038 \text{ g cm}^{-3}$. Intensity data were collected at 150(1) K on a Rigaku AFC6S diffractometer operating in the ω – 2θ mode and using Mo-K α radiation ($\lambda = 0.71069$ Å, graphite monochromator) and were corrected for absorption. Total number of unique reflections 4121, of which 2915 had $I > 3\sigma(I)$. Crystal dimensions $0.45 \times 0.55 \times 0.15$ mm. The structure was solved by direct methods and was refined by full-matrix least-squares calculations to R 0.037 and R_w 0.034. Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.

[†] Satisfactory C, H, N analyses were obtained for **1**, **2**, **3** and **4**.

$^2J_{\text{P-Se}}$ 140 Hz) and 46.8 ($^3J_{\text{P-Pt}}$ 424, $^2J_{\text{P-Se}}$ 71 Hz) attributable to the equivalent Ph_3P ligands and Ph_2P groups, respectively. The ^{77}Se NMR spectrum of **4** is characteristic of an $\text{A}_2\text{BB}'\text{M}$ spin system (where M is ^{77}Se in the most abundant isotopomer for which the Ph_3P phosphorus atoms are magnetically inequivalent). The simulation of this spectrum provides the following coupling constants: $^2J_{\text{P}_\text{A-Se}}$ 71, $^2J_{\text{P}_\text{B-Se}}$ 137 $^2J_{\text{P}_\text{B'-Se}}$ 28, and $^2J_{\text{P}_\text{B-P}_\text{B'}}$ 8 Hz.¹⁵ Thus, the NMR data are consistent with an $\eta^2\text{-Se,Se}'$ bonding mode for **4** analogous to that established by X-ray crystallography for the corresponding $\eta^2\text{-S,S}'$ complex.^{8b} Finally we note that **2** also undergoes oxidative-addition with $\text{Pt}(\text{C}_2\text{H}_4)(\text{PPh}_3)_2$ to give complex **5** in which the $\text{P}_2\text{N}_4\text{Se}_2$ ring acts as a $\mu^2, \eta^3\text{-N,Se,Se}'$ ligands on the basis of ^{31}P NMR spectroscopy. Details will be given in the full account of this work.

In summary, the high-yield synthesis of **1** has facilitated studies of the chemistry of this heterocycle. Initial investigations indicate multimode coordination behaviour for this interesting ligand and the combination of ^{31}P and ^{77}Se NMR spectroscopy provides a powerful structural probe for metal complexes. The presence of a transannular Se-Se interaction in N -bonded platinum(II) adducts of **1** has been established.

We thank the NSERC (Canada) for financial support.

Received, 17th June 1992; Com. 2/03164G

References

- 1 T. M. Klapötke, in *The Chemistry of Inorganic Ring Systems*, ed. R. Steudel, Elsevier, Amsterdam, 1992, pp. 409–427.
- 2 J. Adel, A. El-Kholi, W. Willing, U. Müller and K. Dehnicke, *Chimia*, 1988, **42**, 70; J. Adel and K. Dehnicke, *Chimia*, 1988, **42**, 413.
- 3 E. G. Awere, J. Passmore, P. S. White and T. Klapötke, *J. Chem. Soc., Chem. Commun.*, 1989, 1415; P. K. Gowik, T. M. Klapötke and T. S. Cameron, *J. Chem. Soc., Dalton Trans.*, 1991, 1433.
- 4 P. F. Kelly, A. M. Z. Slawin, D. J. Williams and J. D. Woollins, *J. Chem. Soc., Chem. Commun.*, 1989, 408.
- 5 P. F. Kelly, A. M. Z. Slawin, D. J. Williams and J. D. Woollins, *Polyhedron*, 1990, **9**, 1567.
- 6 T. Chivers, D. D. Doxsee and J. F. Fait, *J. Chem. Soc., Chem. Commun.*, 1989, 1703.
- 7 N. Burford, T. Chivers, M. N. S. Rao and J. F. Richardson, *Inorg. Chem.*, 1984, **23**, 1946; T. Chivers, G. Dénès, S. W. Liblong and J. F. Richardson, *Inorg. Chem.*, 1989, **29**, 3683.
- 8 (a) T. Chivers, K. S. Dhathathreyan and T. Ziegler, *J. Chem. Soc., Chem. Commun.*, 1989, 86; (b) T. Chivers, M. Edwards, A. Meetsma, J. C. van de Grampel and A. van der Lee, *Inorg. Chem.*, 1992, **30**, 2156.
- 9 T. Chivers, G. Y. Dénès, S. W. Liblong and J. F. Richardson, *Inorg. Chem.*, 1989, **28**, 3683.
- 10 T. Chivers and R. W. Hilt, *Inorg. Chem.*, submitted for publication.
- 11 P. Cherin and P. Unger, *Acta Crystallogr., Sect. B*, 1972, **28**, 313.
- 12 N. Burford, J. Passmore and J. C. P. Sanders, in *From Atoms to Polymers—Isoelectronic Analogies*, eds. J. F. Liebman and A. Greenberg, VCH, Weinheim, 1989, p. 53.
- 13 H. Bärnighausen, T. v. Volkman and J. Jander, *Acta Crystallogr.*, 1966, **21**, 571.
- 14 R. K. McMullen, D. J. Prince and J. D. Corbett, *Inorg. Chem.*, 1971, **10**, 1749.
- 15 PANIC NMR simulation program supplied by Brüker.