

# Highly Selective and Divergent Acyl and Aryl Cross-Couplings of Amides via Ir-Catalyzed C–H Borylation/N–C(O) Activation

Pengcheng Gao and Michal Szostak\*

Cite This: https://dx.doi.org/10.1021/acs.orglett.0c02105



Ar

Ŕ,

Pd catalyst

-CO -NR'R'

catalyst, Bapina

Article Recommendations

ACCESS

# III Metrics & More

**ABSTRACT:** Herein, we demonstrate that amides can be readily coupled with nonactivated arenes via sequential Ir-catalyzed C–H borylation/N–C(O) activation. This methodology provides facile access to biaryl ketones and biaryls by the sterically controlled Ir-catalyzed C–H borylation and divergent acyl and decarbonylative amide N–C(O) and C–C activation. The methodology diverts the traditional acylation and arylation regioselectivity, allowing us to directly utilize readily available arenes and amides to produce valuable ketone and biaryl motifs.

**N** ondirected activation of C–H bonds is among the most important tools for transforming functional groups in organic synthesis.<sup>1</sup> The value of nondirected C–H functionalizations lies in the fact that these transformations enable the construction of complex fragments by exploiting readily available arenes (Figure 1A).<sup>2</sup> One of the most promising applications of these methods is Ir-catalyzed C–H borylation, wherein the development of sterically and ligand-controlled C–H functionalizations enables transformative fragment



 ${\bf B};$  This work: highly selective and divergent acyl and aryl coupling of non-activated arenes with amides



■ first tandem borylation/acylation diverting Friedel-Crafts selectivity

Figure 1. (A) Meta-borylation in organic synthesis. (B) Present work.

couplings by exploiting new routes to organoboron building blocks of broad relevance to pharmaceuticals, functional materials, and the synthesis of fine chemicals.<sup>2–5</sup>

divergent C\_H/N\_C

activations

Divergent AcvI and ArvI Coupling of Non-Activated Arenes with Amides

First tandem Borylation/Acylation diverting Friedel-Crafts Selectivity

R" acyclic, twisted

R'

**SUPPORTING Information** 

Ar

Ŕ.

catalyst, Bapina

Pd catalyst

-NR'R

Simultaneously, recent years have witnessed the development of powerful strategies for activation of amide N–C(O) bonds,<sup>6</sup> whereby the traditional amidic resonance ( $n_N \rightarrow \pi^*C=O$  delocalization)<sup>7,8</sup> is diverted into facile oxidative addition of the N–C(O) amide bond by ground-state destabilization and diminution of amidic resonance.<sup>9–12</sup> The value of amide cross-coupling lies in the fact that amides are among the most ubiquitous functional groups in organic synthesis, serve as essential scaffolds in polymers, and most importantly constitute key linkages in peptides and proteins.<sup>6–8</sup> A prominent feature of amide bond cross-coupling is that the acyl-metal intermediate formed after metal insertion can undergo CO extrusion, resulting in an overall N–C(O)/C–C bond activation to afford an aryl-metal, which allows for the direct installation of arenes by deamidative cleavage.<sup>13</sup>

Recently, we became interested in establishing the synergistic merger of nondirected C–H activation with amide N–C(O) cross-coupling (Figure 1B). The utilization of readily available arenes<sup>1–5</sup> as an abundant source of coupling partners together with the selective breaking of nitrogen–carbon bonds in amides<sup>6–8</sup> hold significant promise to define new paradigms in the transformation of functional groups crucial to many synthetic and biological processes.<sup>14</sup>

Received: June 25, 2020



#### pubs.acs.org/OrgLett

Letter

Scheme 1. Acylation of Arylboronate Esters with Amides by Sequential C-H/N-C(O) Activation: Amide Scope<sup>a</sup>



<sup>*a*</sup>Conditions: **2** (2.0 equiv),  $B_2pin_2$  (0.75 equiv),  $[Ir(cod)(OMe)]_2$  (0.5 mol %), dtbpy (1 mol %), THF, 80 °C, 24 h, **1** (1.0 equiv), [Pd(IPr)(cin)Cl] (3 mol %),  $K_2CO_3$  (3.0 equiv), THF: $H_2O$  = 9:1 (0.20 M), 23 °C, 16 h. <sup>*b*</sup> $Cs_2CO_3$  (3.0 equiv), 60 °C.



Scheme 2. Acylation of Arylboronate Esters with Amides by Sequential C-H/N-C(O) Activation: Arene Scope<sup>a</sup>

<sup>*a*</sup>Conditions: see Scheme 1. <sup>*b*</sup>[Pd(IPr)(cin)Cl] (5 mol %). <sup>*c*</sup>Cs<sub>2</sub>CO<sub>3</sub>, 60 °C. <sup>*d*</sup>KOH, 60 °C. <sup>*e*</sup>Cs<sub>2</sub>CO<sub>3</sub>, THF, 80 °C. <sup>*f*</sup>Step 1: hexane. <sup>*g*</sup>N-Ms-N-phenylbenzamide.

Herein, we detail the successful development of highly selective and divergent acyl and aryl cross-couplings of amides with nonactivated arenes via C-H/N-C(O) activation. This methodology connects the classic sterically hindered Ircatalyzed C–H borylation<sup>2,3</sup> with the biorelevant manifold of amide bond activation<sup>6–8</sup> to provide straightforward access to

biaryl ketones and biaryls by two inert bond activation events and diverting the traditional acylation and arylation regioselectivity. The method shows excellent functional group tolerance, chemo- and regioselectivity. We demonstrate the synthetic utility by late-stage derivatization of pharmaceuticals and conjugative cross-coupling of bioactive molecules. The

# Scheme 3. Arylation of Arylboronate Esters with Amides by Sequential C-H/N-C(O)/C-C Activation: Scope<sup>a</sup>



<sup>*a*</sup>Conditions: **2** (2.0 equiv),  $B_2pin_2$  (0.6 equiv),  $[Ir(cod)(OMe)]_2$  (0.5 mol %), dtbpy (1 mol %), THF, 80 °C, 24 h, **1** (1.0 equiv),  $[Pd(dppb)Cl_2]$  (5 mol %), NaHCO<sub>3</sub> (4 equiv), MeB(OH)<sub>2</sub> (1.5 equiv), dioxane (0.125 M), H<sub>2</sub>O, 160 °C, 16 h. <sup>*b*</sup>2-Methyl-N-Ms-N-phenylbenzamide. See SI for details.

synergistic merger with nonactivated arenes opens the door to routinely utilize amides as acyl and aryl cross-coupling electrophiles in a wide range of chemical processes.<sup>1–14</sup>

The arylation of N,N-Boc-benzamide, which is readily prepared in a single step from benzamide,<sup>7a</sup> with 4-tol-Bpin was selected as the starting point of our study to identify conditions for the cross-coupling of boronic esters with amides (Table S1, Supporting Information). Encouragingly, we found that the broadly applicable, air- and moisture-stable [Pd(IPr)-(cin)Cl] (Neolyst CX31, cin = cinnamyl)<sup>15</sup> serves as an efficient catalyst for this coupling to give the model biaryl ketone product in 98% yield (entry 13). Notably, this crosscoupling with boronic esters was found to be sensitive to the amount of water and temperature, controlling the release of aryl boronic acid,<sup>16</sup> with the best results obtained in THF:H<sub>2</sub>O (9:1) at 23 °C (cf. Table S2 for arylation). The key difference between the two Pd catalytic systems (Table S1 vs Table S2) is that Pd-NHCs facilitate oxidative addition/reductive elimination steps due to strong  $\sigma$ -donation and flexible bulk,<sup>9a</sup> respectively, while Pd-phosphine systems in the presence of weak base favor decarbonylation due to slowing the transmetalation relative to the CO deinsertion step.<sup>13</sup> Note that the substrates are presented on schemes to reflect how the reactions were carried out. [Pd(IPr)(cin)Cl)] is the catalyst of choice for amide bond Suzuki cross-coupling.9a,15

With the identified conditions in hand, we explored the scope of the sequential C–H/N–C(O) activation (Scheme 1). The borylation was conducted with 0.75 equiv of  $B_2pin_2$  in the presence of 0.50 mol % of  $[Ir(cod)(OMe)]_2$  and 1.0 mol % of dtbpy.<sup>2a,b</sup> Arylboronate esters were subjected to the C(O)–N amide cross-coupling after removing volatiles and addition of amide derivatives. 1,3-Dimethoxybenzne was used for the borylation reaction as the model arene substrate.<sup>2a,b</sup> As shown

in Scheme 1, the scope of the reaction with respect to amide precursors is very broad and encompasses a wide variety of functional groups. Electron-rich (3a, 3c, 3e, 3t), sterically hindered (3b, 3o, 3p), and electron-deficient (3d, 3f, 3g, 3h) groups were well-tolerated. Electrophilic functional groups, such as ester (3f), carbonate (3j, 3p), cyano (3k), and nitro (31), which would be problematic with classical Weinreb amides, gave the desired ketone products in high yields. Furthermore, heterocyclic amides, including thiophene (3i), pyrazine (3q), and pyridine (3r) readily underwent coupling. Of note, the latter example (3r) represents a direct activation of vitamin B3 (nicotinamide), while 4-hydroxybenzamide (3i) is a common amide pharmaceutical intermediate, demonstrating potential pharmaceutical applications (vide infra). Notably, in all cases, the meta-substituted product was formed with exquisite selectivity, which is in sharp contrast to the traditional acylation with arenes giving ortho products.

The scope of the arene was also extensively studied (Scheme 2). Pleasingly, we found that this C-H/N-C(O) acylation is compatible with a broad range of 1,3-disubstitued arenes, giving the cross-coupling products in good to excellent yields. Acylation of challenging electron-deficient organoboranes substituted with ester groups (**3u**, **3x**, **3y**) is well-tolerated. Pivalates (**3v**), alkyl groups (**3w**), ethers (**3z**, **3aa**), ketones (**3ab**), heterocycles (**3ac**, **3ad**), amides (**3af**), amines (**3ag**), and fluorinated arenes (**3ah**, **3ai**) afford the desired ketone products in high yields, in all cases featuring exclusive arylation regioselectivity. The divergent chemoselectivity for the activated amide is noteworthy (**3af**). Notably, this method could be employed for the direct derivatization of esters of natural products and pharmaceuticals with complex architecture as illustrated by the esters of fructose (**3aj**), menthol

## Scheme 4. Coupling of Arenes with Amides by C-H/N-C(O) Activation: Complex Arenes and Amides<sup>a</sup>



<sup>a</sup>Conditions: **2** (2.0 equiv),  $B_{2}pin_{2}$  (0.75 equiv),  $[Ir(cod)(OMe)]_{2}$  (0.5 mol %), dtbpy (1 mol %), THF, 80 °C, 24 h, **1** (1.0 equiv), [Pd(IPr)(cin)Cl] (3 mol %),  $K_{2}CO_{3}$  (3.0 equiv), THF: $H_{2}O$  = 9:1 (0.20 M), 23 °C, 16 h.

(3ak), *trans*-androsterone (3al), cholesterol (3am), as well as adapalene (3an) and febuxostat (3ao).

The applicability to various amides was explored (Figure S1). This C-H/N-C(O) activation works with *N*-acyl-glutarimides (1w), *N*-acyl amides (1x), *N*-Ts-sulfonamides (1y), and *N*-Boc-carbamates (1z), which expands the scope of the amide component to N-cyclic and N-acyclic amides.<sup>6-9</sup>

Encouraged by the success of C-H/N-C(O) acylative coupling, we considered the possibility of developing divergent aryl coupling by triple C-H/N-C(O)/C-C bond activation (Scheme 3). For this type of cross-coupling, the control of the rate of boronic acid release is critical to match with decarbonylation of the acyl metal.<sup>13</sup> Although the initial results gave no or little conversion to the biaryl product, we discovered that the use of the MeB(OH)<sub>2</sub> additive facilitates the organoboronate release under these conditions,<sup>16</sup> leading to the efficient coupling (Table S2).

We examined the generality of the C-H/N-C(O)/C-C biaryl coupling (Scheme 3). This method is compatible with diverse substituents on both reaction components, including esters (4a, 4d, 4e, 4g), heterocycles (4b), electronically deactivated substrates (4c), halides (4f), tosylates (4h), ketones (4j), and sterically hindered substrates (4m), delivering the biaryl products in good to high yields. Notably, this method delivers functional handles for further derivatization, including electrophilic groups.

Most crucially, the synthetic potential of this biaryl C–H/N-C(O)/C-C coupling was demonstrated in the rapid latestage modification of esters of natural products, such as menthol (4q), fructose (4r), and *trans*-androsterone (4s), thus underscoring the synthetic potential of this activation platform.

An assessment of various amides in the biaryl coupling was performed (Figure S2, Supporting Information). Pleasingly, N-cyclic (1w) as well as N-acyclic amides, such as N-acyl (1x), N-Ts-sulfonamide (1y), and N-Ms-sulfonamide (1ak), can be utilized in this coupling protocol to deliver biaryls from arenes and amides in good yields.

Finally, we applied our C-H/N-C(O) activation technology to conjugative coupling of bioactive molecules (Scheme 4). Thus, the direct derivatization of complex arenes and amides is readily accomplished to deliver conjugates between

complex bioactive molecules and amides under standard conditions (3aq-3ar).<sup>1b</sup>

In conclusion, an expedient method for the cross-coupling of nonactivated arenes with amides via sequential Ir-catalyzed C– H borylation/N–C(O) activation has been developed. This reaction provides facile access to biaryl ketones and biaryls by exploiting regioselective C–H activation with the divergent reactivity of the amide bond by acyl and decarbonylative pathways. This methodology tolerates a wide variety of functional groups, including various amides and direct derivatization of natural products and pharmaceuticals with intricate architectures. Based on this methodology, we demonstrated conjugative coupling between complex biomolecules. This study opens the door for arene–amide coupling in a wide variety of chemical processes.

#### ASSOCIATED CONTENT

#### Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.orglett.0c02105.

Procedures and analytical data (PDF)

#### AUTHOR INFORMATION

### **Corresponding Author**

Michal Szostak – Department of Chemistry, Rutgers University, Newark, New Jersey 07102, United States; Orcid.org/0000-0002-9650-9690; Email: michal.szostak@rutgers.edu

#### Author

**Pengcheng Gao** – Department of Chemistry, Rutgers University, Newark, New Jersey 07102, United States

Complete contact information is available at: https://pubs.acs.org/10.1021/acs.orglett.0c02105

#### Notes

The authors declare no competing financial interest.

# ACKNOWLEDGMENTS

Rutgers University and the NSF (CAREER CHE-1650766) are gratefully acknowledged for support. The 500 MHz spectrometer was supported by the NSF-MRI grant (CHE-1229030). P.G. thanks the China Scholarship Council (No. 201606240022) for a fellowship. We thank Dr. Shicheng Shi (Rutgers University) for preliminary studies. We thank Mr. Qi Wang and Prof. Hao Chen (NJIT) for assistance with HRMS measurements.

# REFERENCES

(1) For representative reviews, see: (a) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. The medicinal chemist's toolbox for late stage functionalization of drug-like molecules. *Chem. Soc. Rev.* **2016**, *45*, 546–576. (b) Blakemore, D. C.; Castro, L.; Churcher, I.; Rees, D. C.; Thomas, A. W.; Wilson, D. M.; Wood, A. Organic Synthesis Provides Opportunities to Transform Drug Discovery. *Nat. Chem.* **2018**, *10*, 383–394.

(2) For reviews, see: (a) Hartwig, J. F. Borylation and Silylation of C-H Bonds: A Platform for Diverse C-H Bond Functionalizations. *Acc. Chem. Res.* **2012**, *45*, 864–873. (b) Mkhalid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F. C-H Activation for the Construction of C-B Bonds. *Chem. Rev.* **2010**, *110*, 890–931.

(3) For representative examples, see: (a) Maleczka, R. E., Jr.; Shi, F.; Holmes, D.; Smith, M. R., III C-H Activation/Borylation/ Oxidation: A One-Pot Unified Route To Meta-Substituted Phenols Bearing Ortho-/Para-Directing Groups. J. Am. Chem. Soc. 2003, 125, 7792-7793. (b) Robbins, D. W.; Hartwig, J. F. Sterically controlled alkylation of arenes through iridium-catalyzed C-H borylation. Angew. Chem., Int. Ed. 2013, 52, 933-937. (c) Murphy, J. M.; Tzschucke, C. C.; Hartwig, J. F. One-Pot Synthesis of Arylboronic Acids and Aryl Trifluoroborates by Ir-Catalyzed Borylation of Arenes. Org. Lett. 2007, 9, 757-760. (d) Robbins, D. W.; Hartwig, J. F. A C-H Borylation Approach to Suzuki-Miyaura Coupling of Typically Unstable 2-Heteroaryl and Polyfluorophenyl Boronates. Org. Lett. 2012, 14, 4266-4269. (e) Joliton, A.; Carreira, E. M. Ir-Catalyzed Preparation of SF5-Substituted Potassium Aryl Trifluoroborates via C-H Borylation and Their Application in the Suzuki-Miyaura Reaction. Org. Lett. 2013, 15, 5147-5149. (f) Hong, B.; Li, C.; Wang, Z.; Chen, J.; Li, H.; Lei, X. Enantioselective Total Synthesis of (-)-Incarviatone A. J. Am. Chem. Soc. 2015, 137, 11946-11949. (g) Murphy, J. M.; Liao, X.; Hartwig, J. F. Meta Halogenation of 1,3-Disubstituted Arenes via Iridium-Catalyzed Arene Borylation. J. Am. Chem. Soc. 2007, 129, 15434-15435. (h) Liskey, C. W.; Liao, X.; Hartwig, J. F. Cyanation of Arenes via Iridium-Catalyzed Borylation. J. Am. Chem. Soc. 2010, 132, 11389-1391. (i) Feng, Z.; Min, Q. Q.; Fu, X. P.; An, L.; Zhang, X. Chlorodifluoromethane-triggered formation of difluoromethylated arenes catalysed by palladium. Nat. Chem. 2017, 9, 918-923.

(4) For further examples, see: (a) Cho, J. Y.; Tse, M. K.; Holmes, D.; Maleczka, R. E., Jr.; Smith, M. R., III Remarkably Selective Iridium Catalysts for the Elaboration of Aromatic C-H Bonds. Science 2002, 295, 305-308. (b) Ishiyama, T.; Takagi, J.; Ishida, K.; Miyaura, N.; Anastasi, N. R.; Hartwig, J. F. Mild Iridium-Catalyzed Borylation of Arenes. High Turnover Numbers, Room Temperature Reactions, and Isolation of a Potential Intermediate. J. Am. Chem. Soc. 2002, 124, 390-391. (c) Chotana, G. A.; Rak, M. A.; Smith, M. R., III Sterically Directed Functionalization of Aromatic C-H Bonds: Selective Borylation Ortho to Cyano Groups in Arenes and Heterocycles. J. Am. Chem. Soc. 2005, 127, 10539-10544. (d) Boller, T. M.; Murphy, J. M.; Hapke, M.; Ishiyama, T.; Miyaura, N.; Hartwig, J. F. Mechanism of the Mild Functionalization of Arenes by Diboron Reagents Catalyzed by Iridium Complexes. Intermediacy and Chemistry of Bipyridine-Ligated Iridium Trisboryl Complexes. J. Am. Chem. Soc. 2005, 127, 14263-14278. (e) Eliseeva, M. N.; Scott, L. T. Pushing the Ir-Catalyzed C-H Polyborylation of Aromatic Compounds to Maximum Capacity by Exploiting Reversibility. J. Am.

Chem. Soc. 2012, 134, 15169–15172. (f) Preshlock, S. M.; Ghaffari, B.; Maligres, P. E.; Krska, S. W.; Maleczka, R. E., Jr.; Smith, M. R., III High-Throughput Optimization of Ir-Catalyzed C–H Borylation: A Tutorial for Practical Applications. J. Am. Chem. Soc. 2013, 135, 7572–7582.

(5) Fyfe, J. W. B.; Watson, A. J. B. Recent Developments in Organoboron Chemistry: Old Dogs, New Tricks. *Chem.* **2017**, *3*, 31–55.

(6) Greenberg, A.; Breneman, C. M.; Liebman, J. F. *The Amide Linkage: Structural Significance in Chemistry, Biochemistry and Materials Science*; Wiley-VCH: New York, 2003.

(7) For studies on amide bond ground-state destabilization pertinent to N-C activation, see: (a) Szostak, R.; Shi, S.; Meng, G.; Lalancette, R.; Szostak, M. Ground-State Distortion in N-Acyl-tert-butylcarbamates (Boc) and N-Acyl-tosylamides (Ts): Twisted Amides of Relevance to Amide N-C Cross-Coupling. J. Org. Chem. 2016, 81, 8091-8094. (b) Meng, G.; Shi, S.; Lalancette, R.; Szostak, R.; Szostak, M. Reversible Twisting of Primary Amides via Ground State N-C(O) Destabilization: Highly Twisted Rotationally Inverted Acyclic Amides. J. Am. Chem. Soc. 2018, 140, 727-734. (c) Liu, C.; Shi, S.; Liu, Y.; Liu, R.; Lalancette, R.; Szostak, R.; Szostak, M. The Most Twisted Acyclic Amides: Structures and Reactivity. Org. Lett. 2018, 20, 7771-7774.

(8) For selected examples, see: (a) Roughley, S. D.; Jordan, A. M. The medicinal chemist's toolbox: an analysis of reactions used in the pursuit of drug candidates. J. Med. Chem. 2011, 54, 3451-3479.
(b) Kaspar, A. A.; Reichert, J. M. Future Directions for Peptide Therapeutics Development. Drug Discovery Today 2013, 18, 807-817. (c) Marchildon, K. Polyamides: Still Strong After Seventy Years. Macromol. React. Eng. 2011, 5, 22-54.

(9) For representative reviews on N–C bond activation, see: (a) Shi, S.; Nolan, S. P.; Szostak, M. Well-Defined Palladium(II)-NHC (NHC = N-Heterocyclic Carbene) Precatalysts for Cross-Coupling Reactions of Amides and Esters by Selective Acyl CO–X (X = N, O) Cleavage. Acc. Chem. Res. **2018**, 51, 2589–2599. (b) Liu, C.; Szostak, M. Decarbonylative Cross-Coupling of Amides. Org. Biomol. Chem. **2018**, 16, 7998–8010. (c) Bourne-Branchu, Y.; Gosmini, C.; Danoun, G. N-Boc-Amides in Cross-Coupling Reactions. Chem. - Eur. J. **2019**, 25, 2663–2674. (d) Chaudhari, M. B.; Gnanaprakasam, B. Recent Advances in the Metal-Catalyzed Activation of Amide Bonds. Chem. - Asian J. **2019**, 14, 76–93.

(10) For selected examples, see: (a) Meng, G.; Szostak, M. Sterically Controlled Pd-Catalyzed Chemoselective Ketone Synthesis via N-C Cleavage in Twisted Amides. Org. Lett. 2015, 17, 4364-4367. (b) Meng, G.; Shi, S.; Szostak, M. Palladium-Catalyzed Suzuki-Miyaura Cross-Coupling of Amides via Site-Selective N-C Bond Cleavage by Cooperative Catalysis. ACS Catal. 2016, 6, 7335-7339. (c) Hie, L.; Fine Nathel, N. F.; Shah, T. K.; Baker, E. L.; Hong, X.; Yang, Y.-F.; Liu, P.; Houk, K. N.; Garg, N. K. Conversion of Amides to Esters by the Nickel-Catalysed Activation of Amide C-N Bonds. Nature 2015, 524, 79-83. (d) Ni, S.; Zhang, W.; Mei, H.; Han, J.; Pan, Y. Ni-Catalyzed Reductive Cross-Coupling of Amides with Aryl Iodide Electrophiles via C-N Bond Activation. Org. Lett. 2017, 19, 2536-2539. (e) Amani, J.; Alam, R.; Badir, S.; Molander, G. A. Synergistic Visible-Light Photoredox/Nickel-Catalyzed Synthesis of Aliphatic Ketones via N-C Cleavage of Imides. Org. Lett. 2017, 19, 2426-2429. (f) Reina, A.; Krachko, T.; Onida, K.; Bouyssi, D.; Jeanneau, E.; Monteiro, N.; Amgoune, A. Development and Mechanistic Investigations of a Base-Free Suzuki-Miyaura Cross-Coupling of  $\alpha, \alpha$ -Difluoroacetamides via C-N Bond Cleavage. ACS Catal. 2020, 10, 2189-2197. (g) Buchspies, J.; Szostak, M. Recent Advances in Acyl Suzuki Cross-Coupling. Catalysts 2019, 9, 53.

(11) For representative decarbonylative coupling, see: (a) Meng, G.; Szostak, M. General Olefin Synthesis by the Palladium-Catalyzed Heck Reaction of Amides: Sterically Controlled Chemoselective N–C Activation. *Angew. Chem., Int. Ed.* **2015**, *54*, 14518–14522. (b) Shi, S.; Meng, G.; Szostak, M. Synthesis of Biaryls through Nickel-Catalyzed Suzuki-Miyaura Coupling of Amides by Carbon-Nitrogen Bond Cleavage. *Angew. Chem., Int. Ed.* **2016**, *55*, 6959–6963. (c) Meng, G.; Szostak, M. Rhodium-Catalyzed C–H Bond Functionalization with Amides by Double C–H/C–N Bond Activation. Org. Lett. 2016, 18, 796–799. (d) Yue, H.; Guo, L.; Liao, H. H.; Cai, Y.; Zhu, C.; Rueping, M. Catalytic Ester and Amide to Amine Interconversion: Nickel-Catalyzed Decarbonylative Amination of Esters and Amides by C–O and C–C Bond Activation. Angew. Chem., Int. Ed. 2017, 56, 4282–4285. (e) Yue, H.; Guo, L.; Lee, S. C.; Liu, X.; Rueping, M. Selective Reductive Removal of Ester and Amide Groups from Arenes and Heteroarenes through Nickel-Catalyzed C–O and C–N Bond Activation. Angew. Chem., Int. Ed. 2017, 56, 3972–3976. (f) Shi, S.; Szostak, M. Decarbonylative Cyanation of Amides by Palladium Catalysis. Org. Lett. 2017, 19, 3095–3098. (g) Zhou, P. X.; Shi, S.; Wang, J.; Zhang, Y.; Li, C.; Ge, C. Palladium/copper-catalyzed decarbonylative heteroarylation of amides via C–N activation. Org. Chem. Front. 2019, 6, 1942–1947.

(12) For studies on N-C activation, see: (a) For a biomimetic esterification, see: Wybon, C. C. D.; Mensch, C.; Hollanders, K.; Gadais, C.; Herrebout, W. A.; Ballet, S.; Maes, B. U. W. Zn-Catalyzed tert-Butyl Nicotinate-Directed Amide Cleavage as a Biomimic of Metallo-Exopeptidase Activity. ACS Catal. 2018, 8, 203-218. (b) For a Cr-catalyzed N-C activation, see: Chen, C.; Liu, P.; Luo, M.; Zeng, X. Kumada Arylation of Secondary Amides Enabled by Chromium Catalysis for Unsymmetric Ketone Synthesis under Mild Conditions. ACS Catal. 2018, 8, 5864–5858. For a  $\sigma$  bond N–C activation, see: (c) Zhang, Z. B.; Ji, C. L.; Yang, C.; Chen, J.; Hong, X.; Xia, J. B. Nickel-Catalyzed Kumada Coupling of Boc-Activated Aromatic Amines via Nondirected Selective Aryl C-N Bond Cleavage. Org. Lett. 2019, 21, 1226-1231. (d) Ni, S.; Li, C.-X.; Mao, Y.; Han, J.; Wang, Y.; Yan, H.; Pan, Y. Ni-catalyzed deaminative cross-electrophile coupling of Katritzky salts with halides via C-N bond activation. Sci. Adv. 2019, 5, 9516. (e) Hu, J.; Wang, M.; Pu, X.; Shi, Z. Nickelcatalysed retro-hydroamidocarbonylation of aliphatic amides to olefins. Nat. Commun. 2017, 8, 14993. For a recent pertinent C-H borylation, see: (f) Lv, J.; Chen, X.; Xue, X. S.; Zhao, B.; Liang, Y.; Wang, M.; Jin, L.; Yuan, Y.; Han, Y.; Zhao, Y.; Lu, Y.; Zhao, J.; Sun, W. Y.; Houk, K. N.; Shi, Z. Metal-free directed sp2 -C-H borylation. Nature 2019, 575, 336-340.

(13) Zhou, T.; Ji, C.-L.; Hong, X.; Szostak, M. Palladium-Catalysed Decarbonylative Suzuki-Miyaura Cross-Coupling of Amides by Carbon-Nitrogen Activation. *Chem. Sci.* **2019**, *10*, 9865–9871.

(14) (a) Carey, F. A.; Sundberg, R. J. Advanced Organic Chemistry: Part B: Reactions and Synthesis; Springer: New York, 2001. (b) Smith, M. B.; March, J. Advanced Organic Chemistry; Wiley: New York, 2007. (15) Marion, N.; Navarro, O.; Mei, J.; Stevens, E. D.; Scott, N. M.; Nolan, S. P. Modified (NHC)Pd(allyl)Cl (NHC = N-heterocyclic carbene) complexes for room-temperature Suzuki-Miyaura and Buchwald-Hartwig reactions. J. Am. Chem. Soc. 2006, 128, 4101– 4111.

(16) Hinkes, S. P. A.; Klein, C. D. P. Virtues of Volatility: A Facile Transesterification Approach to Boronic Acids. *Org. Lett.* **2019**, *21*, 3048–3052.