

Tetrahedron Letters 40 (1999) 4909-4912

TETRAHEDRON LETTERS

Resin-bound Dendrimers as High Loading Supports for Solid Phase Chemistry

Anju Mahajan, Siri Ram Chhabra and Weng C. Chan*

School of Pharmaceutical Sciences, University of Nottingham, University Park, Nottingham NG7 2RD, England

Received 7 April 1999; accepted 6 May 1999

Abstract

The synthesis of a *tris*-Boc tri-amino acid and its utility in the generation of dendrimers as inert scaffolds on solid supports is described. These high loading amine functionalized resins were successfully used in the syntheses of small molecules. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords : Amino acids and derivatives; dendrimers; solid phase synthesis

Dendrimer research has evolved into an area of considerable interest in recent years. Dendrimers are highly branched macromolecules built generationally from a central core by stepwise and controlled processes. These dendrimeric structures differ from typical polymers in possessing well defined molecular architectures. Thus, dendrimers have found widespread applications as catalyst carriers, drug delivery systems, complexation agents, *etc.* [1-4].

The solution phase synthesis of dendrimers is either by the 'divergent approach' involving repetitive multi-step synthesis [5-7] or by the 'convergent method' where small fragments of the dendrimer are assembled on a multi-functional core [8-10]. However, both these approaches suffer from the disadvantage of repetitive time-consuming purification. In contrast, dendrimer synthesis on solid supports [2,3] makes the purification procedure simpler as excess reagents ensure reaction completion and can be easily washed off from the polymer bound dendrimer molecule.

The naturally occurring amino acid lysine has been used as a building block in the generation of inert dendrimeric scaffolds, which are commercially available for the synthesis of highly immunogenic multiple antigenic peptides [11]. It was anticipated that the unique triamino acids described in the previous paper [12] could be exploited for the generation of dendritic scaffolds on solid supports. We envisaged that repeat coupling of the protected triamino acid 4 onto an amine functionalized resin would lead to a geometrical increase in the number of reactive amine functionalities giving extremely high loading per reaction bead.

The synthesis of 4 was readily accomplished by acylating the secondary amine

e-mail: weng.chan@nottingham.ac.uk

functionality in the tri-amino ester 1 with di-*tert*-butyl dicarbonate followed by saponification of the methyl ester to give 2. Catalytic hydrogenolysis gave the *bis*-Boc amino acid 3. Finally, acylation of 3 at the α -amino position by treatment with (Boc)₂O in presence of triethylamine afforded the required *tris*-Boc tri-amino acid 4¹ (Scheme 1). Attempts to synthesize 4 from 2 by one-pot catalytic hydrogenolysis in the presence of (Boc)₂O was not successful as the product obtained was heavily contaminated with the N^{α} . N^{α} -diBoc derivative of 3.



Scheme1 Reagents and conditions : (i) $(Boc)_2O$; (ii) NaOH, MeOH–water; (iii) H₂ / 10% Pd-C, MeOH; (iv) $(Boc)_2O$, Et₃N.

The *tris*-Boc amino acid **4** was readily attached to aminomethylated PS resin (0.8 mmol/g) preswollen in DMF, *via* the DIPCDI-HOAt mediated coupling (Scheme 2). Complete removal of the three Boc protecting groups was effected by treatment of the above resin with 50% TFA-DCM in the presence of TES to give **5**, the first generation amines.



Scheme 2 Reagents and conditions : (i) HOAt–DIPCDI, DMF; (ii) 50% TFA–DCM, TES; (iii) Rink linker, HOAt–DIPCDI, DMF; (iv) 4, HATU–DIPEA,DMF.

¹ Spectral data of N^2 , N^4 , N^6 -tris-tert-butoxycarbonyl-2(S), 6-diamino-4-azahexanoic acid 4:

 $[\]nu_{max}$ 1701 (C=O), 3373 (NH) cm⁻¹; δ_{H} (250 MHz, CDCl₃) 1.41, 1.42, 1.45 (27 H, 3 x s, 3 (CH₃)₃), 2.85-3.89 (6 H, m, 3 CH₂), 3.90-4.78 (1H, m, α -H), 4.88-5.50 (1 H, m, NH), 5.66-6.40 (1H, m, NH), 8.57 (1 H, br s, COOH); δ_{C} (62.9 MHz, CDCl₃) 28.18, 28.27 (3 (CH₃)₃), 39.17 (CH₂), 47.88 (CH₂), 48.95 (CH₂), 52.69 (CH), 79.46, 79.97, 80.94 (3 *C*(CH₃)₃), 155.85 (3 *COC*(CH₃)₃), 173.33 (COOH); *m/z* (FAB) 470 (M+Na), 448 (M+H); HRMS (FAB) 470.2472 (M+Na, C₂₀H₃₇N₃O₈Na requires *m/z* 470.2478), 448.2656 (M+H, C₂₀H₃₈N₃O₈ requires *m/z* 448.2659).

The Rink amide linker [13] was then coupled onto resin 5 to give the derivatised PS resin 6. The efficiency of all coupling steps and deprotections was monitored using the TNBS amine test [14]. Significantly, the resin 6 exhibited substitution of 0.9 mmol/g (89%) based on spectrophotometric determination of the Fmoc-derived chromophore liberated upon treatment with 20% piperidine in DMF using $\varepsilon_{230} = 5253 \text{ M}^{-1} \text{ cm}^{-1}$.

Furthermore, resin 5 was re-exposed to the *tris*-Boc amino acid 4 using standard coupling conditions, which when followed by TFA treatment gave the second generation amines 7. Attachment of the Rink linker to the resin 7 using standard coupling conditions was monitored by a negative amine test. Once again, the resin 8 exhibited an excellent loading of 1.17 mmol/g (84%) based on UV analysis of Fmoc deprotection product, dibenzofulvene-piperidine adduct.

Thus, the high loading dendrimeric resins of 1st and 2nd generation amines were readily obtained, particularly coupling of the linker at the secondary amine functionality was easily achieved. Although the loading per gram of the resin does not increase substantially due to the mass of the compound attached to the resin, the loading per reaction bead increases geometrically from one generation to the next.

To assess the physical appearance of the resin beads, the unmodified aminomethylated resin beads and the second generation dendrimer beads were swollen in DCM and analyzed under an optical microscope with 100 x magnification (Figure 1). The second generation dendrimer beads are clearly larger in size compared to the parent aminomethylated polystyrene.



Figure 1

The high loading efficiency of these dendrimeric resins was further evaluated by the synthesis of 2-naphthalenesulphonamide 10 and N-phenylacetyl-L-phenylalaninamide 13. Thus, resin 6 was subjected to Fmoc deprotection, followed by reaction with 2-naphthalenesulphonyl chloride in the presence of DIPEA to give the derivatised resin 9. 2-Naphthalenesulphonamide 10 was then selectively cleaved from the resin by treatment with 90% TFA-water in 84% yield, leaving the dendrimer core intact (Scheme 3). The identity and purity of 10 was confirmed by ES-MS, ¹H and ¹³C NMR.

For synthesis of compound 13, the resin 6 was treated with 20% piperidine–DMF followed by acylation with FmocPheOH using standard coupling conditions. The resin 11 showed a loading of 0.9 mmol/g. Fmoc-removal followed by derivatisation with phenylacetic acid afforded resin 12. Compound 13 was then released as before in 87% yield.

Moreover, the above sequences were repeated on the 2^{nd} generation amines and the compounds 10 and 13 were obtained in near quantitative yields on cleavage.



Scheme 3 Reagents and conditions : (i) 20% piperidine/DMF; (ii) 2-naphthalenesulphonyl chloride, DIPEA, DMF; (iii) 90% TFA/water; (iv) FmocPheOH, HATU, DIPEA, DMF; (v) PhCH₂CO₂H, HOBt, TBTU, DIPEA.

In summary, we have demonstrated the construction of a new class of resin-attached dendrimers by using our novel *tris*-Boc tri-amino acid as a building block. These dendrimers provide a rapid and efficient method of increasing resin loading. Furthermore, we envisaged that *via* orthogonal protection on the three amine functionalities, it is possible to devise a multi-release system for generation of one-bead-one-compound libraries.

Abbreviations Boc, *tert*-butoxycarbonyl; DIPCDI, *N,N'*-diisopropylcarbodiimide; DIPEA, diisopropylethylamine; DCM, dichloromethane; DMF, *N,N'*-dimethylformamide; FAB, fast-atom-bombardment; Fmoc, 9-fluorenylmethoxycarbonyl; HOAt, 1-hydroxyazabenzotriazole; HOBt, 1-hydroxybenzotriazole; TBTU, *O*-(1*H*-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetra-fluoroborate; TES, triethylsilane; TFA, trifluoroacetic acid; TNBS, 2,4,6-trinitrobenzenesulfonic acid; Z, benzyloxycarbonyl.

Acknowledgements AM would like to thank the Commonwealth Scholarship Commission, U.K. for financial support.

References

- [1] Newkome GR, Moorefield CN, Vogtle F. Dendritic Molecules: Concepts, Synthesis, Perspectives; VCH; Wenheim, 1996.
- [2] Swali V, Wells NJ, Langley GJ, Bradley M. J. Org. Chem. 1997;62,4902-4903.
- [3] Wells NJ, Davies M, Bradley M. J. Org. Chem. 1998;63,6430-6431.
- [4] Chow HF, Mong TKK, Nongrum MF, Wan CW. Tetrahedron 1998;54:8543-8660.
- [5] Tomalia DA, Baker H, Dewald JR, Hall M, Kallos G, Martin S, Roeck J, Ryder J, Smith P. Polym. J. 1985;17:117-132.
- [6] Newkome GR, Nayak A, Behera RK, Moorefield CN, Baker GR. J. Org. Chem. 1992;57:358-362.
- [7] Moors R, Vogtle F. Chem. Ber. 1993;126:2133-2135.
- [8] Hawker CJ, Frechet JMJ. J. Am. Chem. Soc. 1990;112:7638-7647.
- [9] Wooley KL, Hawker CJ, Frechet JMJ. J. Chem. Soc., Perkin Trans. 1 1991;1059-1076.
- [10] Zeng F, Zimmerman SC. J. Am. Chem. Soc. 1996;118:5326-5327.
- [11] Tam JP. Proc. Natl. Acad. Sci. U.S.A. 1988;85:5409-5413.
- [12] Chhabra SR, Mahajan A, Chan WC. Tetrahedron Lett. 1999;40: preceding letter
- [13] Rink H. Tetrahedron Lett. 1987;28:3787-3790.
- [14] Hancock WS, Battersby JE. Anal. Biochem. 1976;71:260-264.