

This article was downloaded by: [University of Winnipeg]

On: 15 September 2014, At: 06:27

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

Olefinic Peptide Nucleic Acid (OPA)

Daniel Gautschi^a & Christian J. Leumann^{a b}

^a Department of Chemistry and Biochemistry, University of Bern, Bern, Switzerland

^b Department of Chemistry and Biochemistry, University of Bern, Frelestrasse 3, CH-3012, Bern, Switzerland

Published online: 31 Aug 2006.

To cite this article: Daniel Gautschi & Christian J. Leumann (2003) Olefinic Peptide Nucleic Acid (OPA), Nucleosides, Nucleotides and Nucleic Acids, 22:5-8, 1211-1213, DOI: [10.1081/NCN-120022838](https://doi.org/10.1081/NCN-120022838)

To link to this article: <http://dx.doi.org/10.1081/NCN-120022838>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Olefinic Peptide Nucleic Acid (OPA)

Daniel Gautschi and Christian J. Leumann*

Department of Chemistry and Biochemistry, University of Bern,
Bern, Switzerland

ABSTRACT

The olefinic peptide nucleic acid analogues (OPA) monomers containing the bases thymine and adenine were synthesised in 11 steps. Fully modified oligomers containing these units were prepared and their pairing properties assessed by means of UV-melting experiments.

Polyamide or peptide nucleic acids (PNAs), first described in 1991,^[1] are DNA analogues based entirely on an achiral polyamide backbone. A key structural feature of PNA monomers **1** is the central amide linker between the base and the backbone. This tertiary amide functionality is conformationally labile and occurs in both the *E*- and *Z*-rotameric forms in uncomplexed PNA. In PNA/PNA, PNA/DNA and PNA/RNA duplexes, the carbonyl oxygen of the amide linker unit points uniformly to the carboxy terminus of the strand. OPA^[2] was designed to remove this ambiguity. The central amide functionality is replaced by an isostructural, configurationally stable C=C double bond in either the *E* (**2**) or the *Z* (**3**) configuration.

*Correspondence: Christian J. Leumann, Department of Chemistry and Biochemistry, University of Bern, Frelestrasse 3, CH-3012 Bern, Switzerland; E-mail: leumann@ioc.unibe.ch.







Figure 1. Chemical structure of the monomeric units of the different peptide nucleic acid analogues.



Scheme 1. a) *n*-BuLi, THF, -78°C , 10 min, then $(\text{CHO})_n$, r.t., 1.5 h (83%); b) Red-Al, THF, addition 0°C -r.t., then NIS, -78°C , 10 min (90%); c) *t*BuPh₂SiCl, imidazole, THF, r.t., 16 h (97%); d) Reformatsky reagent: BrCH₂CO₂Et, Zn, CH₂(OCH₃)₂, reflux, 30 min; coupling: Pd(PPh₃)₄, DMPU, 65°C , 2 h (88%); e) PPTSA, EtOH, 45°C , 2.5 h (80%); f) PPh₃, LiN₃, CBr₄, DMF, 0°C , 16 h (68%); g) HF·Py/Py 1:9, CH₃CN, r.t., 2.5 h; h) DIAD, PPh₃, N³-benzoylthymine, THF, r.t., 2 h, (74% over two steps); i) DIAD, PPh₃, N⁶-benzoyladenine, DMF, 0°C , 16 h, (50% over two steps); j) LiOH(aq) 1 M, Dioxane, 0°C , 16 h, (74%); k) Lindlar-catalyst, H₂, MeOH, r.t., 16 h; l) MMT-Cl, Et₃N, DMSO, 16 h, (65%).

Table 1. Mass spectrometry data and T_m values [$^{\circ}\text{C}$] (UV-melting-curves, 260 nm) of PNA, All-*E*-OPA and All-*Z*-OPA with parallel and antiparallel DNA ($c = 4 \mu\text{M}$ in 100 mM NaCl, 10 mM, Na_2HPO_4 , pH 7.0).

Sequence	m/z calcd	m/z found (ESI ⁺ TOF)	T_m (antiparallel DNA) ^a	T_m (parallel DNA) ^b
20 dC(ttttaata)-Gly-NH ₂ ^c	3060.9	3060.9	29.4	13.1
21 dC(t ^E t ^E t ^E t ^E a ^E a ^E t ^E a ^E t ^E a ^E)-Gly-NH ₂ ^c	2891.0	2891.3	< 0	16.0
22 Lys-t ^Z t ^Z t ^Z t ^Z a ^Z a ^Z t ^Z a ^Z t ^Z a ^Z -Gly-NH ₂	2731.03	2730.75	not detected	21.5

^ad(TATATTAAAA).

^bd(AAAATTATAT).

^cData from 2.

The MMT-*Z*- OPA monomer **15** and **19**, containing the bases thymine and adenine, were now synthesised in 11 steps, starting from THP-protected-3-Butyn-1-ol. The synthesis is outlined in Sch. 1.

In order to study the pairing properties, oligomers **20–22** were prepared using the MMT-/Acyl-strategy,^[3] and the stability of the duplexes formed with anti-parallel and parallel DNA was assessed by means of UV-melting curves (Table 1). All-*Z*-OPA binds preferentially to parallel DNA ($T_m = 21^{\circ}\text{C}$) rather than antiparallel (no pairing detected). Thus it shows opposite strand alignment behavior compared to PNA. This might be a consequence of the structural preorganisation of the monomeric unit, which mimics the rotameric form of PNA that is not observed in complexes. The overall lower affinity to DNA compared to PNA seems to be a consequence of differential solvation or dipole moment, rather than being of conformational origin.

REFERENCES

1. Nielsen, P.E.; Egholm, M.; Berg, R.H.; Burchardt, O. *Science* **1991**, *254*, 1497–1500.
2. Schütz, R.; Cantin, M.; Roberts, C.; Greiner, B.; Uhlmann, E.; Leumann, C. *Angew. Chem. Int. Ed.* **2000**, *39*, 1250–1253.
3. Will, D.W.; Breipohl, G.; Langner, D.; Knolle, J.; Uhlmann, E. *Tetrahedron* **1995**, *51*, 12,069–12,082.



