

54. Nucleosides

Part LX¹⁾

Synthesis and Characterization of Monomeric Cordycepin-Vitamin and Cordycepin-Lipid Conjugates Model Substances for Biodegradable Ester and Carbonate Linkages in Conjugates and Potential Inhibitors of HIV-1 Replication

by Marita Wasner^{a)}, Robert J. Suhadolnik^{b)d)}, Susan E. Horvath^{b)}, Martin E. Adelson^{d)}, Ning Kon^{b)},
Ming-Xu Guan^{c)}, Earl E. Henderson^{c)d)}, and Wolfgang Pfeleiderer^{a)*}

^{a)} Fakultät für Chemie, Universität Konstanz, Postfach 5560, D-78434 Konstanz

^{b)} Department of Biochemistry, ^{c)} Department of Microbiology and Immunology,
and ^{d)} Fels Institute for Cancer Research and Molecular Biology,
Temple University School of Medicine, Philadelphia, PA 19140, USA

Dedicated to Prof. Dr. Dr. h.c. *Hans-Jürgen Bestmann* on the occasion of his 70th birthday
and in admiration to his interesting contribution to organic chemistry

(2.1.96)

Monomeric 3'-deoxyadenosine (cordycepin) was modified at the 2'-*O*- (13–18) and 5'-*O*-position (25–29) by the vitamins E, D₂, and A and by the two lipids 1,2-di-*O*-palmitoylglycerol and 1,2-di-*O*-hexadecylglycerol *via* succinate or carbonate linkages. These base-labile conjugates afforded protection groups like the 2-(4-nitrophenyl)ethoxycarbonyl (npeoc) and monomethoxytrityl group (MeOTr) that are cleavable without harming the ester and carbonate bonds, respectively. Monomeric conjugates of cordycepin and vitamin E, vitamin D₂, 1,2-di-*O*-palmitoylglycerol, and 1,2-di-*O*-hexadecylglycerol (see 13, 14, 17, 18, 25, 26, 28, and 29) inhibited HIV-1-induced syncytia formation 1.7 to 6.2 fold compared to 1.5-fold for cordycepin (see Table); IC₅₀ values for 25 and 28 were 257 and 267 μM, respectively. In addition, the monomeric cordycepin-vitamin and -lipid conjugates inhibited HIV-1 RT activity 28–49% which compares with a 13% inhibition of HIV-1 RT observed for cordycepin. The minimal inhibition of HIV-1-induced syncytia formation and HIV-1 RT activity did not proceed by the activation of RNase L. The monomeric conjugates tested (13, 14) increased PKR expression.


1. Introduction. – The 3'-deoxyadenosine (cordycepin) was discovered by isolation from *Cordyceps militaris* [2]. It shows some inhibitory activity against *Bacillus subtilis*, avian tubercle bacillus, and Ehrlich ascites tumor cells [3]. Recently, cordycepin trimer core d^{3'}(A2'p5'A2'p5'A) [4] was found to be an inhibitor of HIV-1 reverse-transcriptase (RT) [5] [6] by interfering in the primer complex formation of RT to rRNA^{Lys,3} *via* the anticodon region.

However, the polarity of nucleosides and oligonucleotides is a limiting factor for penetration through cell membranes and, therefore, many attempts of conjugate formation are reported in literature to overcome this problem [7–9]. The attachment of cholesterol to the 2'-*O*- and 5'-*O*-position of the sugar moiety of monomeric and trimeric


¹⁾ Part LIX: [1].

cordycepin was already shown to facilitate membrane crossing [10]. So other lipophilic conjugates of monomeric cordycepin were prepared as model substances for testing reaction conditions to synthesize the trimeric analogues. It is the aim to link the lipophilic residues to the cordycepin moiety by ester and carbonate functions, respectively, since this kind of binding will guarantee the anticipated degradation of the pro-drug in the cell after membrane crossing.






2. Syntheses. – The syntheses of cordycepin [11] and its protected derivatives **1** [12], **2** [13], and **19** [12] were already described in the literature. Starting material for the 2'-*O*-conjugates were compounds **1** and **2**, respectively. Reaction of freshly prepared 2-*ambo*- α -tocopheryl chloroformate with compound **1** in abs. CH₂Cl₂ in the presence of 1-methyl-1*H*-



R ¹	R ¹	R ²	n	R ¹	R ²	n	R ²	n
1 MeOTr	3 MeOTr	VitE	0	9 H	VitE	0	13 VitE	0
2 npeoc	4 MeOTr	VitE	1	10 H	VitE	1	14 VitE	1
	5 npeoc	VitD ₂	1	11 H	PL	1	15 VitD ₂	1
	6 npeoc	VitA	1	12 H	EL	1	16 VitA	1
	7 MeOTr	PL	1				17 PL	1
	8 MeOTr	EL	1				18 EL	1



19	R ³	R ³
20	VitE	25 VitE
21	VitD ₂	26 VitD ₂
22	VitA	27 VitA
23	PL	28 PL
24	EL	29 EL



imidazole and 4-(dimethylamino)pyridine (DMAP) gave the fully protected vitamin E conjugate **3** in 65 % yield bearing a carbonate moiety. The structural analogue carrying a succinyl spacer between cordycepin and vitamin E (= 2-*ambo-α*-tocopherol) moieties was synthesized by a one-pot reaction starting from the 2'-OH component **1** which was first treated with succinic anhydride and DMAP and second with vitamin E in the presence of *N*-[3-(dimethylamino)propyl]-*N*'-ethylcarbodiimide hydrochloride (EDC) to give the corresponding conjugate **4** in 74 % yield. Detritylation of the conjugates **3** and **4** by 2 % TsOH in CH₂Cl₂/MeOH 4:1 afforded the intermediates **9** and **10**, respectively, in excellent yields. Because of the acid lability of the vitamins D₂ and A, the monomethoxytrityl protecting group had to be replaced in the starting material by the base-labile β-eliminating 2-(4-nitrophenyl)ethoxycarbonyl (npeoc) group [14]. Compound **2** [13] was reacted with succinic anhydride and DMAP to give the corresponding 3'-deoxy-*N*⁶,5'-*O*-bis[2-(4-nitrophenyl)ethoxycarbonyl]-2'-*O*-succinyladenosine which was isolated for subsequent esterification by the carbodiimide method (EDC) with vitamin D₂ (ergocalciferol) in the absence of O₂ and light, to give **5** in 56 % yield. Conjugate formation of **2** with A (retinol) took place in a similar manner, but without isolation of the succinyl derivative to give **6** in 74 % yield. In the case of the succinyl ester conjugates with 1,2-di-*O*-palmitoylglycerol [15] [13] and 1,2-di-*O*-hexadecylglycerol [16] [13], the fully protected conjugates **7** and **8** were detritylated by 2 % TsOH in CH₂Cl₂/MeOH 4:1 to give **11** (78 %) and **12** (56 %). The final deblocking was achieved by β-elimination with 0.5M 1,8-diazabicyclo[5.3.0]undec-7-ene (DBU) in abs. pyridine to result in the compounds **13–18**.

The syntheses of the 5'-*O*-conjugates were performed by one-pot reactions starting from *N*⁶,2'-bis[2-(4-nitrophenyl)ethoxycarbonyl]-3'-deoxyadenosine (**19**) [13] which was treated first with succinic anhydride and DMAP, followed by carbodiimide-activated esterification with the different lipophilic alcohols to give **20–24**. β-Elimination of the npeoc groups with 0.5M DBU in abs. pyridine resulted in the 5'-*O*-conjugates **25–29**. It was noticed that the deprotected vitamin A conjugates **16** and **27** are extremely instable to light and O₂ in comparison to their precursors **6** and **22**, respectively. Correct elemental analyses could, therefore, not be obtained for **16** and **27**.

3. Biological Application. – The 2'-*O*- and 5'-*O*-monomeric cordycepin-vitamin and -lipid conjugates were evaluated with respect to give biological activities: *i*) activation of RNase L, *ii*) HIV-1-induced syncytia formation, *iii*) HIV-1 reverse-transcriptase activity, *iv*) PKR expression, and *v*) PKR activity (*Table*). Cordycepin inhibited HIV-1-induced syncytia formation 1.5-fold, compared with a 1.7- to 6.2-fold inhibition observed with the monomeric cordycepin-vitamin and -lipid conjugates **13**, **14**, **17**, **18**, **25**, **26**, **28**, and **29**. IC₅₀ Values for monomeric conjugates **25** and **28** were 254 and 267 μM, respectively. The monomeric conjugates (300 μM) inhibited HIV-1 RT in the range 28–49 %. None of the monomeric cordycepin-vitamin and -lipid conjugates activated recombinant, human GST-RNase L. In view of the inhibition of syncytia formation and HIV-1 RT activity observed in the absence of RNase L activation, the effect of the (2'–5')A conjugates on PKR expression was measured. The two monomeric conjugates **13** and **14** tested resulted in an increase in PKR expression of 65 and 92 %, respectively. Conjugate **14** with a succinyl linker showed a greater increase in PKR expression than did conjugate **13** with a carbonyl linker. The monomeric cordycepin conjugates were used as model substances

Table. Inhibition of HIV-1 Replication and Biological Activities of Monomeric Cordycepin-Vitamin and -Lipid Conjugates 13, 14, 17, 18, 25, 26, 28, and 29^{a)}

		Inhibition of syncytia formation ^{b)}	Inhibition of HIV-1 RT activity [%] ^{c)}	Activation of RNase L ^{d)}	PKR Expression ^{e)} ([%] change)
13	Cordycepin 2'-(vitamin E carbonate) ^{f)}	1.9	49	0	+65
14	Cordycepin 2'-(vitamin E succinate) ^{f)}	1.8	46	0	+92
17	Cordycepin 2'-(2,3-di- <i>O</i> -palmitoyl- glyceryl succinate) ^{f)}	3.9	36	0	n.t. ^{g)}
18	Cordycepin 2'-(2,3-di- <i>O</i> -hexadecyl- glyceryl succinate) ^{f)}	3.7	28	0	n.t. ^{g)}
25	Cordycepin 5'-(vitamin E succinate) ^{f)}	6.2	44	0	n.t. ^{g)}
26	Cordycepin 5'-(vitamin D ₂ succinate) ^{f)}	4.2	30	0	n.t. ^{g)}
28	Cordycepin 5'-(2,3-di- <i>O</i> -palmitoyl- glyceryl succinate) ^{f)}	2.6	31	0	n.t. ^{g)}
29	Cordycepin 5'-(2,3-di- <i>O</i> -hexadecyl- glyceryl succinate) ^{f)}	1.7	43	0	n.t. ^{g)}
	Vitamin D ₂	n.t. ^{g)}	21	n.t. ^{g)}	n.t. ^{g)}
	Cordycepin	1.5	13	n.t. ^{g)}	-27
	Adenine	1.3	9.8	n.t. ^{g)}	n.t. ^{g)}

^{a)} Compounds were tested at 300 μ M.

^{b)} Inhibition of HIV-1 replication was determined by syncytia formation (fold reduction in infection). The mean of triplicate determinations is shown; variance did not exceed 5–10%.

^{c)} Percent inhibition of HIV-1 reverse transcriptase (HIV-1 RT) activity. Control values for HIV-1 RT activity ranged from 24000 to 33000 cpm. The mean of duplicate determinations is shown; variance did not exceed 5–10%.

^{d)} Activation of recombinant RNase L was measured as the percent hydrolysis of [³²P]poly(U) in the presence of cordycepin 2'-*O*- and 5'-*O*-vitamin and -lipid conjugates (10 μ M). The mean of duplicate determinations is shown; variance did not exceed 5–10%.

^{e)} PKR activity was measured as described in the *Exper. Part*. At 20 h post-infection, the quantity of PKR observed varied by < 1%, irrespective of HIV-1 infection.

^{f)} Test compounds were dissolved in MeOH; final concentration of MeOH in the assays was 10%. Data were normalized to a 10% MeOH control.

^{g)} n.t. = not tested.

for the synthesis and characterization of trimeric cordycepin-vitamin and -lipid conjugates.

This research was supported in part by research grants from the U.S. Public Health Service (R01-A1 34765 (R. J. S.), P30-CA-12227, and T32-CA9214 (M. E. A.)) and federal work study awards (S. H.). Our thanks go to Harald Sigmund [13] for giving the lipid derivatives used in this work and Nicolas F. Muto for technical assistance with the HIV-1 RT assays.

Experimental Part

General. TLC: Precoated silica gel TLC sheets *Fl500 LS 254* from Schleicher & Schüll. Prep. TLC: silica gel 60 *PF₂₅₄* (Merck). Prep. column flash chromatography (FC): silica gel for flash chromatography (Baker). HPLC: Merck-Hitachi *L-6200*, *L-3000* photo diode array detector; column *RP18*, 125 $\times$ 4 mm, 5 μ m, Merck; flow rate 1 ml/min. UV/VIS: Perkin-Elmer, *Lambda 5*; $\lambda_{\max}$ in nm (log ϵ). ¹H-NMR: Bruker *AC 250*, δ in ppm rel. to DMSO. EL = 'Ether lipid'.

Bioassay. Recombinant, human RNase L was expressed in *E. coli* (DH5 α) as a fusion protein glutathione-S-transferase(GST)-RNase L as described [17]. The activation of recombinant, human GST-RNase L was measured as the percent of poly(U)[³²P]pCp hydrolysed in the presence of authentic p₃A₃ (10⁻¹⁰–10⁻⁶) or (2'-5')A derivative

as previously described [17]. The infected-centers assay was used to measure the ability of (2'–5')A derivatives (300 μ M) to inhibit HIV-1-induced syncytia formation in human cells as previously described [10]. The effect of (2'–5')A derivatives (300 μ M) on HIV-1 reverse transcriptase (RT) activity was measured as previously described [18]. (2'–5')A derivatives or controls (in 25 μ l) were added to 50 μ l of supernatant from HIV-1 infected Molt4 IIIB cells containing RT activity and incubated for 2 h at 37° in the presence of a cocktail containing 50 mM Tris (pH 8.0), 20 mM DTT (dithiothreitol), 10 mM MgCl₂, 60 mM NaCl, 0.05 % NP-40, 5 μ g/ml of oligodeoxythymidylic acid, 10 μ g/ml of polyriboadenylic acid, 10 mM TTP, and 1 μ Ci of [α -³²P]TTP. 50 μ l of the cocktail were then spotted onto DEAE paper, dried, washed in 2X SSC soln. (3 $\times$, 10 min each), dried, and exposed to radiographic film for 18 to 24 h at –80°. The filters were cut, and final quantification (cpm) was determined by scintillation spectrometry. Test compounds were added at 100 μ M 2 h prior to infection of 1 $\cdot$ 10⁶SupT1 cells with HIV-1 IIIB at a *m.o.i.* of 0.1. NP-40 extracts were prepared 20 h post-infection, and PKR expression was measured as described [19].

1. 3'-Deoxy-N⁶,5'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]adenosine (**2**) [13]. To a soln. of 3'-deoxy-N⁶-[2-(4-nitrophenyl)ethoxycarbonyl]adenosine [12] (1 g, 2.25 mmol) in abs. pyridine (8 ml) at –30° was added dropwise 2-(4-nitrophenyl)ethyl chloroformate [14] (1.5 g, 6.5 mmol) in abs. CH₂Cl₂ (4 ml) within 30 min. Then the mixture was stirred at –10° for 30 min. More 2-(4-nitrophenyl)ethyl chloroformate [14] (800 mg, 3.5 mmol) in abs. CH₂Cl₂ (4 ml) was added at –10°. After 70 min, the mixture was diluted with CHCl₃ (100 ml) and washed with sat. NaHCO₃ soln. (3 $\times$ 50 ml), the aq. phase re-extracted with CHCl₃ (2 $\times$ 50 ml), the combined org. layer dried (MgSO₄), evaporated, and co-evaporated with toluene, and the residue purified by FC (silica gel, 15 $\times$ 3.5 cm, toluene→toluene/AcOEt 4:1→1:1→2:3→1:1 + 1 % MeOH→1:1 + 4 % MeOH→1:1 + 25 % MeOH): 1.05 g (73 %) of **2**. Amorphous solid. UV (MeOH): 266 (4.55). ¹H-NMR ((D₆)DMSO): 10.58 (s, NH); 8.62, 8.49 (2s, H–C(8), H–C(2)); 8.13 (2d, 4 H *o* to NO₂); 7.55 (2d, 4 H *m* to NO₂); 5.99 (d, H–C(1')); 5.79 (d, OH–C(2')); 4.70 (m, H–C(2')); 4.51 (m, H–C(4')); 4.41–4.21 (m, 6 H, 2 H–C(5'), OCH₂CH₂); 3.12–3.00 (2t, 4 H, OCH₂CH₂); 2.25 (m, H–C(3')); 2.03 (m, H–C(3')). Anal. calc. for C₂₈H₂₇N₇O₁₁ (637.6): C 52.75, H 4.27, N 15.38; found: C 52.66, H 4.29, N 15.34.

2. 3'-Deoxy-5'-O-(monomethoxytrityl)-N⁶-[2-(4-nitrophenyl)ethoxycarbonyl]-2'-O-(2-ambo- α -tocopheryl)-oxycarbonyl]adenosine (**3**). The 2-ambo- α -tocopheryl chloroformate was synthesized by adding dropwise 2-ambo- α -tocopherol (431 mg, 1 mmol) and Et₃N (0.14 ml, 1 mmol) in anh. THF to an ice-cooled soln. of trichloromethyl chloroformate (Fluka; 0.181 ml, 1.5 mmol) in anh. THF (4 ml). After stirring (30 min) at 0°, small amounts of charcoal were added, and the mixture was kept for further 15 min. Then the precipitate was filtered twice, and the filtrate was evaporated to a yellow oil (480 mg, 97 %).

To an ice-cooled soln. of 2-ambo- α -tocopheryl chloroformate (247 mg, 0.5 mmol) and 1-methyl-1H-imidazole (41 mg, 0.5 mmol) in abs. CH₂Cl₂ (2 ml) were added some pearls of molecular sieve (4 Å), **1** [12] (179 mg, 0.25 mmol), and catalytic amounts of DMAP. The mixture was kept at r.t. for 2 d, then 2-ambo- α -tocopheryl chloroformate (123 mg, 0.25 mmol) and 1-methyl-1H-imidazole (21 mg, 0.25 mmol) were added, and the mixture was kept at r.t. for another 16 h. FC (silica gel, 16.5 $\times$ 2 cm, toluene→toluene/AcOEt 2:1→3:2→1:1) gave 230 mg of a colourless foam which was dissolved in CH₂Cl₂ (2 ml) and precipitated from petroleum ether (100 ml): 190 mg (65 %) of **3**. Colourless powder. UV (CH₂Cl₂): 273 (sh, 4.47), 267 (4.51), 236 (sh, 4.47). ¹H-NMR ((D₆)DMSO): 10.64 (s, NH); 8.62, 8.59 (2s, H–C(8), H–C(2)); 8.14 (d, 2 H *o* to NO₂); 7.59 (d, 2 H *m* to NO₂); 7.30–7.13 (m, 12 H, MeOTr); 6.78 (d, 2 H *o* to MeO); 6.37 (d, H–C(1')); 5.91 (m, H–C(2')); 4.57 (m, H–C(4')); 4.37 (t, OCH₂CH₂); 3.69 (s, MeO); 3.21 (m, 2 H–C(5')); 3.08 (t, OCH₂CH₂); 2.88–0.7 (m, 2 H–C(3'), tocopheryl). Anal. calc. for C₆₉H₈₅N₆O₁₁ (1174.5): C 70.63, H 7.22, N 7.16; found: C 70.48, H 7.18, N 7.16.

3. 3'-Deoxy-5'-O-(monomethoxytrityl)-N⁶-[2-(4-nitrophenyl)ethoxycarbonyl]-2'-O-{[2-(2-ambo- α -tocopheryloxycarbonyl)ethyl]carbonyl]adenosine (**4**). A mixture of **1** [12] (717 mg, 1 mmol) succinic anhydride (120 mg, 1.2 mmol), and DMAP (159 mg, 1.3 mmol) in abs. CH₂Cl₂ (5 ml) was kept at r.t. for 4.5 h. Then *N*-[3-(dimethylamino)propyl]-*N'*-ethylcarbodiimide hydrochloride (EDC; 250 mg, 1.4 mmol) and 2-ambo- α -tocopherol (603 mg, 1.4 mmol) were added. The mixture was kept at r.t. in the dark for 2 h, then diluted with CHCl₃ (120 ml), washed with 5 % citric acid (60 ml) and sat. NaHCO₃ soln. (60 ml), the aq. phase re-extracted with CHCl₃, the combined org. layer dried (MgSO₄) and evaporated, and the crude product purified by FC (silica gel, 16.5 $\times$ 3 cm, toluene→toluene/AcOEt 1:1→1:1 + 2 % MeOH→1:1 + 3 % MeOH): 911 mg (74 %) of **4**. Amorphous solid. UV (CH₂Cl₂): 273 (sh, 4.47), 267 (4.52), 236 (sh, 4.41). ¹H-NMR ((D₆)DMSO): 10.64 (s, NH); 8.57, 8.56 (2s, H–C(8), H–C(2)); 8.15 (d, 2 H *o* to NO₂); 7.60 (d, 2 H *m* to NO₂); 7.28–7.12 (m, MeOTr); 6.76 (d, 2 H *o* to MeO); 6.24 (s', H–C(1')); 5.84 (m, H–C(2')); 4.46 (m, H–C(4')); 4.38 (t, OCH₂CH₂); 3.68 (s, MeO); 3.2–3.06, 3.0–2.7 (m, 2 H–C(5'), OCH₂CH₂, C(O)CH₂CH₂C(O)); 2.2–0.75 (m, 2 H–C(3'), tocopheryl). Anal. calc. for C₇₂H₈₈N₆O₁₂ (1229.5): C 70.34, H 7.21, N 6.84; found: C 70.24, H 7.31, N 6.82.

4. 3'-Deoxy-2'-O- $\{[2-(\text{ergocalciferoyloxycarbonyl})\text{ethyl}]\text{carbonyl}\}$ -N⁶,5'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]adenosine (**5**). First, 3'-deoxy-N⁶,5'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]-2'-O-succinyladenosine was synthesized by stirring **2** [13] (512 mg, 0.8 mmol), succinic anhydride (480 mg, 4.8 mmol), and DMAP (635 mg, 5.2 mmol) in abs. CH₂Cl₂/DMF 4:1 (15 ml) at r.t. for 2 d. Then MeOH was added, the mixture kept at r.t. for 2 h and evaporated, and the residue purified by FC (silica gel (30 g), $d = 3$ cm, CHCl₃→CHCl₃ + 5% MeOH→CHCl₃ + 10% MeOH) to give a colourless oil which was lyophilized in dry dioxane: 529 mg (90%). Amorphous solid. UV (MeOH): 272 (sh, 4.50), 267 (4.53). ¹H-NMR ((D₆)DMSO): 12.25 (br. COOH); 10.62 (br. NH); 8.62, 8.53 (2s, H-C(8), H-C(2)); 8.15 (m, 4 H *o* to NO₂); 7.55 (2d, 4 H *m* to NO₂); 6.20 (d, H-C(1')); 5.73 (d, H-C(2')); 4.55 (m, H-C(4')); 4.4–4.2 (m, 2 OCH₂CH₂, 2 H-C(5')); 3.10, 3.03 (2t, 2 OCH₂CH₂); 2.7–2.2 (m, C(O)CH₂CH₂C(O), 2 H-C(3')). Anal. calc. for C₃₂H₃₁N₇O₁₄·1/2 dioxane (781.1): C 52.24, H 4.51, N 12.54; found: C 51.66, H 4.47, N 12.13.

A mixture of 3'-deoxy-N⁶,5'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]-2'-O-succinyladenosine (295 mg, 0.4 mmol), EDC (92 mg, 0.48 mmol), DMAP (59 mg, 0.48 mmol), and vitamin D₂ (190 mg, 0.48 mmol) in abs. CH₂Cl₂ (8 ml) was kept at r.t. under N₂ at darkness for 3 h. Then the mixture was diluted with CH₂Cl₂ (80 ml) and washed with sat. NaHCO₃ soln. (2 × 30 ml), the aq. phase re-extracted with CHCl₃, the combined org. layer dried (MgSO₄) and evaporated, and the residue purified by FC (silica gel, 9 × 2 cm, toluene→toluene/AcOEt 1:1 + 2% MeOH→1:1 + 3% MeOH): 250 mg (56%) of **5**. Amorphous solid. UV (MeOH): 272 (sh, 4.68), 266 (4.70). ¹H-NMR (CDCl₃): 8.77, 8.66 (2s, H-C(8), H-C(2)); 8.14–8.07 (m, 4 H *o* to NO₂, NH); 7.41–7.33 (m, 4 H *m* to NO₂); 6.10 (d, H-C(1')); 6.16–6.00 (dd, H-C(6)(VitD₂), H-C(7)(VitD₂)); 5.66 (m, H-C(2')); 5.15 (m, H-C(22)(VitD₂), H-C(23)(VitD₂)); 5.0, 4.75 (2m, 2 H-C(19)(VitD₂)); 4.82 (m, H-C(3)(VitD₂)); 4.55 (m, H-C(4')); 4.5–4.2 (m, 2t, 2 OCH₂CH₂, 2 H-C(5')); 3.12, 3.06 (2t, 2 OCH₂CH₂); 2.8–0.5 (m, C(O)CH₂CH₂C(O), 2 H-C(3')), 36 H (VitD₂)). Anal. calc. for C₆₀H₇₃N₇O₁₄ (1116.3): C 64.56, H 6.59, N 8.78; found: C 64.19, H 6.63, N 8.76.

5. 3'-Deoxy-N⁶,5'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]-2'-O- $\{[2-(\text{retinoyloxycarbonyl})\text{ethyl}]\text{carbonyl}\}$ -adenosine (**6**). As described in *Exper. 3*, with **2** [12] (287 mg, 0.45 mmol), succinic anhydride (54 mg, 0.54 mmol), and DMAP (71 mg, 0.59 mmol) in abs. CH₂Cl₂ (3 ml; 18 h). Then EDC (113 mg, 0.59 mmol) and vitamin A (169 mg, 0.59 mmol; 5 h, N₂, darkness). Workup with CH₂Cl₂ (80 ml) and sat. NaHCO₃ soln. (30 ml). FC (silica gel, 13.5 × 2 cm, toluene/AcOEt 1:1→1:1 + 2% MeOH→1:1 + 3% MeOH→1:1 + 4% MeOH): 337 mg (74%) of **6**. Amorphous solid. UV (CH₂Cl₂): 330 (4.67), 318 (sh, 4.62), 272 (sh, 4.61), 267 (4.63). ¹H-NMR (CDCl₃): 8.71, 8.18–8.10 (s, br. m, H-C(8), H-C(2), 4 H *o* to NO₂, NH); 7.45–7.36 (m, 4 H *m* to NO₂); 6.63, 6.27–6.04 (m, H-C(1'), 5 CH=C(retinyl)); 5.68 (m, H-C(2')); 5.57 (t, H-C(14)(retinyl)); 4.73 (d, 2 H-C(15)); 4.63 (m, H-C(4')); 4.53, 4.44 (2t, 2 OCH₂CH₂); 4.45–4.26 (m, 2 H-C(5')); 3.15, 3.06 (2t, 2 OCH₂CH₂); 2.67 (s, m, H-C(3'), C(O)CH₂CH₂C(O)); 2.25 (dd, H-C(3')); 2.1–1.5 (m, 3s, 2 H-C(4)(retinyl), 2 H-C(2)(retinyl), 2 H-C(3)(retinyl), Me-C(9)(retinyl), Me-C(13)(retinyl), Me-C(5)(retinyl)); 1.00 (s, 2 Me-C(1)(retinyl)). Anal. calc. for C₅₂H₅₉N₇O₁₄ (1006.1): C 62.08, H 5.91, N 9.75; found: C 62.06, H 6.06, N 9.63.

6. 3'-Deoxy-N⁶-[2-(4-nitrophenyl)ethoxycarbonyl]-2'-O-(2-ambo- α -tocopheryloxycarbonyl)adenosine (**9**). Compound **3** (704 mg, 0.6 mmol) was stirred at r.t. in CH₂Cl₂/MeOH 4:1 (12 ml) containing 2% of TsOH·H₂O for 10 min. Then the mixture was diluted with CHCl₃ (150 ml) and washed with sat. NaHCO₃ soln. (2 × 70 ml), the aq. phase re-extracted with CHCl₃, and the combined org. layer dried (MgSO₄) and evaporated. FC (silica gel, 16.5 × 3 cm, CHCl₃→CHCl₃ + 5% MeOH) gave 484 mg of **9** which was dissolved in CH₂Cl₂ (2 ml) and precipitated in petroleum ether (100 ml) and co-evaporated with CH₂Cl₂: 428 mg (79%) of **9**. Amorphous solid. UV (CH₂Cl₂): 273 (sh, 4.38), 267 (4.43). ¹H-NMR ((D₆)DMSO): 10.62 (s, NH); 8.68, 8.62 (2s, d, H-C(8), H-C(2)); 8.14 (d, 2 H *o* to NO₂); 7.60 (d, 2 H *m* to NO₂); 6.32 (d, H-C(1')); 5.72 (m, H-C(2')); 5.11 (t, OH-C(5')); 4.42 (m, H-C(4')); 4.38 (t, OCH₂CH₂); 3.7, 3.55 (2m, 2 H-C(5')); 3.09 (t, OCH₂CH₂); 2.65, 2.3 (2m, 2 H-C(3')); 2.0–0.75 (m, 49 H (tocopheryl)). Anal. calc. for C₄₈H₆₉N₆O₁₀ (902.1): C 65.31, H 7.61, N 9.33; found: C 65.29, H 7.49, N 9.23.

7. 3'-Deoxy-N⁶-[2-(4-nitrophenyl)ethoxycarbonyl]-2'-O- $\{[2-(2\text{-ambo-}\alpha\text{-tocopheryloxycarbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**10**). As described in *Exper. 6*, with **4** (700 mg, 0.57 mmol) and CH₂Cl₂/MeOH 4:1 (11 ml) containing 2% of TsOH·H₂O (20 min). Workup with CHCl₃ (120 ml) and sat. NaHCO₃ soln. (2 × 60 ml). FC (silica gel, 15 × 2 cm, CHCl₃→CHCl₃ + 4% MeOH): 509 mg (93%) of **10**. Amorphous solid. UV (CH₂Cl₂): 273 (sh, 4.39), 267 (4.44). ¹H-NMR ((D₆)DMSO): 10.62 (s, NH); 8.65, 8.60 (2s, H-C(8), H-C(2)); 8.15 (d, 2 H *o* to NO₂); 7.61 (d, 2 H *m* to NO₂); 6.19 ('s', H-C(1')); 5.86 (m, H-C(2')); 5.06 (t, OH-C(5')); 4.38 (m, t, H-C(4'), OCH₂CH₂); 3.65, 3.55 (2m, 2 H-C(5')); 3.10 (t, OCH₂CH₂); 2.9, 2.75 (m, C(O)CH₂CH₂C(O)); 2.55, 2.1 (2m, 2 H-C(3')); 2.0–0.75 (m, 49 H (tocopheryl)). Anal. calc. for C₅₂H₇₂N₆O₁₁ (957.2): C 65.25, H 7.58, N 8.78; found: C 65.12, H 7.62, N 8.83.

8. 3'-Deoxy-2'-O- $\{[2-(2,3\text{-di-}O\text{-palmitoylglycer-1-yloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ -N⁶-[2-(4-nitrophenyl)-ethoxycarbonyl]adenosine (**11**). A mixture of **1** [12] (335 mg, 0.466 mmol), succinic anhydride (56 mg, 0.56 mmol), and DMAP (74 mg, 0.61 mmol) in abs. CH₂Cl₂ (2 ml) was kept at r.t. for 18 h. Then EDC (116 mg, 0.61 mmol) and 1,2-di-*O*-palmitoylglycerol [13] [15] (344 mg, 0.61 mmol) in abs. CH₂Cl₂ (2 ml) were added. The mixture was kept at r.t. for 5 h, then diluted with CHCl₃ (100 ml), and washed with sat. NaHCO₃ soln. (60 ml) and 10% citric acid (50 ml), the aq. phase re-extracted with CHCl₃, the combined org. layer dried (MgSO₄) and evaporated, and the crude product purified by FC (silica gel, 11 × 2 cm, toluene/AcOEt 1:1): **7**, contaminated with some 1,2-di-*O*-palmitoylglycerol. The crude **7** was then treated with CH₂Cl₂/MeOH 4:1 (14 ml) containing 2% of TsOH · H₂O for 25 min. Then the mixture was diluted with CH₂Cl₂ (100 ml) and washed with sat. NaHCO₃ soln. (2 × 50 ml), the aq. phase re-extracted with CH₂Cl₂, the combined org. layer dried (MgSO₄) and evaporated and the residue purified by FC (silica gel, 9 × 2 cm, toluene/AcOEt 1:1 → 1:1 + 7% MeOH): 400 mg (78%) of **11**. Amorphous solid. UV (CH₂Cl₂): 272 (sh, 4.37), 267 (4.42). ¹H-NMR (CDCl₃, 2 rotamers): 8.74, 8.33, 8.19, 8.07 (*s*, 3d, H-C(8), H-C(2), 2 H *o* to NO₂, NH); 7.45 (*d*, 2 H *m* to NO₂); 5.98 (*m*, H-C(1')); 5.65 (*m*, H-C(2')); 5.25 (*q*, H-C(2)(Glyc)); 5.05 (*m*, OH-C(5')); 4.65–4.5 (*m*, H-C(4'), OCH₂CH₂); 4.35–4.05 (*m*, 2 H-C(1)(Glyc), 2 H-C(3)(Glyc), H-C(5')); 3.65 (*m*, H-C(5')); 3.17 (*t*, OCH₂CH₂); 2.93 (*m*, H-C(3')); 2.65 (*m*, C(O)CH₂CH₂C(O)); 2.4–2.25 (*m*, 2 CH₂(α)(Palm), H-C(3')); 1.65 (2 CH₂(β)(Palm)); 1.25 (*m*, 48 H(Palm)); 0.88 (*t*, 2 Me(Palm)). Anal. calc. for C₅₈H₉₀N₆O₁₄ (1095.4): C 63.60, H 8.28, N 7.67; found: C 63.14, H 8.16, N 7.58.

9. 3'-Deoxy-2'-O- $\{[2-(2,3\text{-di-}O\text{-hexadecylglycer-1-yloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ -N⁶-[2-(4-nitrophenyl)-ethoxycarbonyl]adenosine (**12**). As described in *Exper. 8*, with **1** [12] (501 mg, 0.7 mmol), succinic anhydride (84 mg, 0.84 mmol), and DMAP (111 mg, 0.91 mmol) in abs. CH₂Cl₂ (5 ml; 18 h). Then EDC (174 mg, 0.91 mmol) and 1,2-di-*O*-hexadecylglycerol [13] [16] (493 mg, 0.91 mmol) in abs. CH₂Cl₂ (5 ml; 5 h). Workup with CHCl₃ (120 ml), sat. NaHCO₃ soln. (60 ml), and 10% citric acid (80 ml). Purification by FC (silica gel, 5 × 3.5 cm, toluene/AcOEt 1:1 → 1:1 + 2% MeOH → 1:1 + 3% MeOH): **8**, contaminated with some 1,2-di-*O*-hexadecylglycerol. The crude **8** was then treated with CH₂Cl₂/MeOH 4:1 (14 ml) containing 2% of TsOH · H₂O for 15 min. Workup with CH₂Cl₂ (150 ml) and sat. NaHCO₃ (2 × 100 ml). The residue was purified by FC (silica gel, 9 × 2 cm, toluene/AcOEt 1:1 → 1:1 + 2% MeOH → 1:1 + 4% MeOH): 421 mg of **12** (56%). Amorphous solid. UV (CH₂Cl₂): 272 (sh, 4.38), 267 (4.44). ¹H-NMR (CDCl₃): 8.73, 8.21–8.07 (2*s*, *m*, H-C(8), H-C(2), 2 H *o* to NO₂, NH); 7.45 (*d*, 2 H *m* to NO₂); 5.99 (*m*, H-C(1')); 5.62 (*m*, H-C(2')); 5.00 (*m*, OH-C(5')); 4.62–4.50 (*m*, H-C(4'), OCH₂CH₂); 4.35–4.05 (*m*, 2 H-C(5'), H-C(2)(Glyc)); 3.7–3.35, 3.15 (*m*, OCH₂CH₂, 2 H-C(1)(Glyc), 2 H-C(3)(Glyc), 2 CH₂(α)(EL)); 2.95, 2.25 (2*m*, 2 H-C(3')); 2.65 (*m*, C(O)CH₂CH₂C(O)); 1.7, 1.55, 1.3 (3*m*, 56 H(EL)); 0.85 (*t*, 2 Me(EL)). Anal. calc. for C₅₈H₉₄N₆O₁₂ (1067.4): C 65.26, H 8.88, N 7.87; found: C 65.06, H 8.76, N 7.65.

10. 3'-Deoxy-2'-O-(2-ambo-α-tocopheryloxy-carbonyl)adenosine (**13**). Compound **9** (318 mg, 0.35 mmol) was co-evaporated twice with abs. pyridine and then dissolved in abs. pyridine (3.5 ml). DBU (268 mg, 1.76 mmol) was added, the mixture kept at r.t. for 18 h, then AcOH (106 mg, 1.76 mmol) added, and the soln. evaporated. The residue was diluted with CHCl₃ (60 ml) and washed with sat. NaCl soln. (2 × 30 ml), the aq. phase re-extracted with CHCl₃, the combined org. layer dried (MgSO₄), evaporated, and co-evaporated with toluene, and the residue purified by FC (silica gel, 16 × 2 cm, CHCl₃ → CHCl₃ + 5% MeOH): 200 mg (94%) of **13**. Amorphous solid. UV (CH₂Cl₂): 275 (sh, 3.81), 260 (4.12). ¹H-NMR ((D₆)DMSO): 8.36, 8.13 (2*s*, H-C(8), H-C(2)); 7.34 (*s*, NH₂); 6.20 (*d*, H-C(1')); 5.69 (*m*, H-C(2')); 5.20 (*t*, OH-C(5')); 4.40 (*m*, H-C(4')); 3.69, 3.54 (2*m*, 2 H-C(5')); 2.69, 2.40 (2*m*, 2 H-C(3')); 2.10–0.75 (*m*, 49 H(tocopheryl)). Anal. calc. for C₄₀H₆₁N₅O₆ (708.0): C 67.86, H 8.68, N 9.89; found: C 67.76, H 8.59, N 9.74.

11. 3'-Deoxy-2'-O- $\{[2-(2\text{-ambo-}\alpha\text{-tocopheryloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**14**). As described in *Exper. 10*, with **10** (248 mg, 0.26 mmol), and DBU (197 mg, 1.3 mmol) in abs. pyridine (3 ml; 18 h), then AcOH (78 mg, 1.3 mmol). Workup with CHCl₃ (50 ml) and sat. NaCl soln. (2 × 20 ml). Purification by FC (silica gel, 16 × 2 cm, CHCl₃ → CHCl₃ + 3% MeOH → CHCl₃ + 5% MeOH): 143 mg (72%) of **14**. Amorphous solid. UV (CH₂Cl₂): 275 (sh, 3.80), 259 (4.13). ¹H-NMR ((D₆)DMSO): 8.32, 8.12 (2*s*, H-C(8), H-C(2)); 7.32 (*s*, NH₂); 6.08 (*d*, H-C(1')); 5.64 (*m*, H-C(2')); 5.12 (*t*, OH-C(5')); 4.31 (*m*, H-C(4')); 3.61, 3.50 (2*m*, 2 H-C(5')); 2.91, 2.74 (*m*, C(O)CH₂CH₂C(O)); 2.65–0.75 (*m*, 2 H-C(3'), 49 H(tocopheryl)). Anal. calc. for C₄₃H₆₅N₅O₇ (764.0): C 67.60, H 8.58, N 9.17; found: C 67.55, H 8.44, N 8.96.

12. 3'-Deoxy-2'-O- $\{[2-(\text{ergocalciferlyloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**15**). As described in *Exper. 10*, with **5** (180 mg, 0.16 mmol) and DBU (491 mg, 3.22 mmol) in abs. pyridine (6.5 ml; 17 h, N₂, darkness), then AcOH (193 mg, 3.22 mmol). Workup with CH₂Cl₂ (80 ml) and sat. NaHCO₃ soln. (2 × 40 ml). Purification by FC (silica gel, 5.5 × 2 cm, toluene/AcOEt 1:1 → 1:1 + 4% MeOH → 1:1 + 6% MeOH): 104 mg (88%) of **15**. Amorphous solid. UV (MeOH): 262 (4.39). ¹H-NMR ((D₆)DMSO): 8.31, 8.12 (2*s*, H-C(8), H-C(2)); 7.31 (br. NH₂); 6.06 (*d*,

H–C(1''); 6.20–5.95 (*dd*, H–C(6)(VitD₂), H–C(7)(VitD₂)); 5.60 (*m*, H–C(2'')); 5.20 (*m*, H–C(22)(VitD₂), H–C(23)(VitD₂), OH–C(5'')); 5.06, 4.7 (*2m*, 2 H–C(19)(VitD₂)); 4.8 (*m*, H–C(3)(VitD₂)); 4.31 (*m*, H–C(4'')); 3.65–3.45 (*m*, 2 H–C(5'')); 2.85–0.5 (*m*, C(O)CH₂CH₂C(O), 2 H–C(3'), 36 H(VitD₂)). Anal. calc. for C₄₂H₅₉N₅O₆ (730.0): C 69.11, H 8.15, N 9.59; found: C 68.57, H 8.16, N 9.41.

13. 3'-Deoxy-2'-O- $\{[2-(retinylloxycarbonyl)ethyl]carbonyl\}$ adenosine (**16**). As described in *Exper. 10*, with **6** (151 mg, 0.15 mmol) and 0.5M DBU in abs. MeCN (4.56 ml; 18 h), then 2M AcOH (1.14 ml). Workup with CHCl₃ (80 ml) and H₂O (2 $\times$ 30 ml). Purification by FC (silica gel, 10 $\times$ 1 cm, CHCl₃ $\rightarrow$ CHCl₃ + 2% MeOH $\rightarrow$ CHCl₃ + 5% MeOH $\rightarrow$ CHCl₃ + 10% MeOH): 83 mg (89%) of **16**. Amorphous solid. UV (CH₂Cl₂): 330 (4.60), 320 (*sh*, 4.56), 259 (4.28). ¹H-NMR (CDCl₃): 8.25, 7.89 (2*s*, H–C(8), H–C(2)); 6.65–5.5 (*m*, H–C(1'), H–C(2'), 5 CH=C(*retinyl*), OH–C(5'), H–C(14)(*retinyl*)); 4.75 (*d*, 2 H–C(15)); 4.55 (*m*, H–C(4'')); 4.60, 3.65 (*m*, 2 H–C(5'')); 2.65 (*br*, C(O)CH₂CH₂C(O)); 2.95, 2.15 (*2m*, 2 H–C(3'')); 2.05–1.5 (*m*, 2*s*, 2 H–C(4)(*retinyl*), 2 H–C(2)(*retinyl*), Me–C(3)(*retinyl*), Me–C(9)(*retinyl*), Me–C(13)(*retinyl*), Me–C(5)(*retinyl*)); 1.00 (*s*, 2 Me–C(1)(*retinyl*)). Anal. calc. for C₃₄H₄₅N₅O₆ $\cdot$ 1/8 CHCl₃ (634.7): C 64.58, H 7.12, N 11.03; found: C 64.21, H 7.23, N 10.59; a correct C, H, N anal. could not be obtained, due to some instability of **16** against light and oxidation.

14. 3'-Deoxy-2'-O- $\{[2-(2,3-di-O-palmitoylglycer-1-yloxycarbonyl)ethyl]carbonyl\}$ adenosine (**17**). As described in *Exper. 10*, with **11** (340 mg, 0.31 mmol) and DBU (236 mg, 1.55 mmol) in abs. pyridine (3 ml; 18 h), then AcOH (93 mg, 1.55 mmol). Workup with CHCl₃ (100 ml) and sat. NaCl soln. (2 $\times$ 50 ml). Purification by FC (silica gel, 10.5 $\times$ 2 cm, CHCl₃ $\rightarrow$ CHCl₃ + 6% MeOH): 256 mg (92%) of **17**. Amorphous solid. UV (CH₂Cl₂): 259 (4.11). ¹H-NMR (CDCl₃, 2 rotamers): 8.27, 7.84 (2*s*, H–C(8), H–C(2)); 5.90 (*m*, H–C(1'')); 5.82 (*br*, NH₂); 5.62 (*m*, H–C(2'')); 5.24 (*q*, H–C(2)(Glyc)); 4.54 (*t*, OH–C(5'')); 4.29–4.05 (*m*, H–C(4'), 2 H–C(1)(Glyc), 2 H–C(3)(Glyc), H–C(5'')); 3.65 (*m*, H–C(5'')); 2.90 (*m*, H–C(3'')); 2.61 (*m*, C(O)CH₂CH₂C(O)); 2.28 (*m*, 2 CH₂(α)(*Pal*)), H–C(3'')); 1.57 (*m*, 2 CH₂(β)(*Pal*)); 1.22 (*m*, 48 H(*Pal*)); 0.85 (*t*, 2 Me(*Pal*)). Anal. calc. for C₄₉H₈₃N₅O₁₀ (902.2): C 65.23, H 9.27, N 7.76; found: C 65.58, H 9.12, N 7.56.

15. 3'-Deoxy-2'-O- $\{[2-(2,3-di-O-hexadecylglycer-1-yloxycarbonyl)ethyl]carbonyl\}$ adenosine (**18**). As described in *Exper. 10*, with **12** (300 mg, 0.28 mmol) and DBU (320 mg, 2.1 mmol) in abs. MeCN/pyridine 2:1 (4.2 ml; 18 h), then AcOH (126 mg, 2.1 mmol). Workup with CHCl₃ (100 ml) and 10% citric acid soln. (50 ml). Purification by FC (silica gel, 13.5 $\times$ 2 cm, CHCl₃ $\rightarrow$ CHCl₃ + 5% MeOH): 202 mg (82%) of **18**. Amorphous solid. UV (CH₂Cl₂): 259 (4.14). ¹H-NMR (CDCl₃): 8.31, 7.87 (2*s*, H–C(8), H–C(2)); 6.0–5.8 (*m*, H–C(1'), OH–C(5'), NH₂); 5.64 (*m*, H–C(2'')); 4.57 (*m*, H–C(4'')); 4.2–4.0 (*m*, 2 H–C(5'), H–C(2)(Glyc)); 3.65–3.3 (*m*, 2 H–C(1)(Glyc), 2 H–C(3)(Glyc), 2 CH₂(α)(EL)); 2.95, 2.15 (*2m*, 2 H–C(3'')); 2.65 (*m*, C(O)CH₂CH₂C(O)); 1.7–1.2 (*m*, 56 H(EL)); 0.88 (*t*, 2 Me(EL)). Anal. calc. for C₄₉H₈₇N₅O₈ (874.3): C 67.32, H 10.03, N 8.01; found: C 67.33, H 9.85, N 8.09.

16. 3'-Deoxy-N⁶,2'-O-bis $\{[2-(4-nitrophenyl)ethoxycarbonyl]-5'-O-[2-(2-ambo-\alpha-tocopheryloxycarbonyl)ethyl]carbonyl\}$ adenosine (**20**). As described in *Exper. 3*, with **19** [12] (192 mg, 0.3 mmol), succinic anhydride (36 mg, 0.36 mmol), and DMAP (48 mg, 0.39 mmol) in abs. CH₂Cl₂ (3 ml; 18 h). Then EDC (75 mg, 0.39 mmol) and vitamin E (181 mg, 0.44 mmol; 5 h), more EDC (29 mg, 0.15 mmol), vitamin E (65 mg, 0.15 mmol), and DMAP (18 mg, 0.15 mmol; 18 h). Workup with CHCl₃ (70 ml) and 10% citric acid soln. (40 ml). FC (silica gel, 13.5 $\times$ 2 cm, toluene/AcOEt 1:1 $\rightarrow$ 1:1 + 4% MeOH $\rightarrow$ 1:1 + 5% MeOH): 260 mg (75%) of **20**. Amorphous solid. UV (CH₂Cl₂): 272 (*sh*, 4.51), 266 (4.54). ¹H-NMR (CDCl₃): 8.73, 8.31, 8.20–8.13 (2*s*, *m*, H–C(8), H–C(2), 4 H *o* to NO₂, NH); 7.45–7.38 (*m*, 4 H *m* to NO₂); 6.10 (*s*'), H–C(1'')); 5.67 (*m*, H–C(2'')); 4.65 (*m*, H–C(4'')); 4.5–4.25 (*m*, 2 OCH₂CH₂, 2 H–C(5'')); 3.2–2.5 (*m*, 2 OCH₂CH₂, C(O)CH₂CH₂C(O)); 2.25 (*m*, H–C(3'')); 2.05–0.85 (*m*, H–C(3'), 49 H(*tocopheryl*)). Anal. calc. for C₆₁H₇₉N₇O₁₅ (1150.3): C 63.69, H 6.92, N 8.52; found: C 63.60, H 7.00, N 8.17.

17. 3'-Deoxy-5'-O- $\{[2-(ergocalciferilyloxycarbonyl)ethyl]carbonyl\}$ -N⁶,2'-O-bis $\{[2-(4-nitrophenyl)ethoxycarbonyl]adenosine$ (**21**). As described in *Exper. 3*, with **19** [12] (255 mg, 0.4 mmol), succinic anhydride (48 mg, 0.48 mmol), and DMAP (64 mg, 0.52 mmol) in abs. CH₂Cl₂ (8 ml; 18 h). Then EDC (100 mg, 0.52 mmol) and vitamin D₂ (206 mg, 0.52 mmol; 3 h, darkness). Workup with CHCl₃ (80 ml) and sat. NaHCO₃ soln. (40 ml). Purification by FC (silica gel, toluene/AcOEt 1:1 $\rightarrow$ 1:1 + 4% MeOH $\rightarrow$ 1:1 + 5% MeOH): 354 mg (79%) of **21**. Amorphous solid. UV (CH₂Cl₂): 272 (*sh*, 4.66), 267 (4.68). ¹H-NMR (CDCl₃): 8.74, 8.21–8.14 (*s*, *m*, H–C(8), H–C(2), 4 H *o* to NO₂, NH); 7.46–7.38 (*m*, 4 H *m* to NO₂); 6.12 (*d*, H–C(1'')); 6.20, 6.00 (*2d*, H–C(6)(VitD₂), H–C(7)(VitD₂)); 5.70 (*m*, H–C(2'')); 5.19 (*m*, H–C(22)(VitD₂), H–C(23)(VitD₂)); 5.04, 4.82 (*2m*, 2 H–C(Vit19)(D₂)); 4.92 (*m*, H–C(3)(VitD₂)); 4.66–4.25 (*m*, H–C(4'), 2 OCH₂CH₂, 2 H–C(5'')); 3.25–3.1 (*m*, 2 OCH₂CH₂); 2.9–0.5 (*m*, C(O)CH₂CH₂C(O), 2 H–C(3'), 36 H(VitD₂)). Anal. calc. for C₆₀H₇₃N₇O₁₄ (1116.28): C 64.56, H 6.59, N 8.78; found: C 64.18, H 6.55, N 8.89.

18. 3'-Deoxy-5'-O- $\{[2-(\text{retinylloxycarbonyl})\text{ethyl}]\text{carbonyl}\}$ -N⁶,2'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]-adenosine (**22**). As described in *Exper.* 3, with **19** [12] (287 mg, 0.45 mmol), succinic anhydride (54 mg, 0.54 mmol), and DMAP (71 mg, 0.59 mmol) in abs. CH₂Cl₂ (3 ml; 18 h). Then EDC (113 mg, 0.59 mmol) and vitamin A (169 mg, 0.59 mmol; 5 h, N₂, darkness). Workup with CH₂Cl₂ (80 ml) and sat. NaHCO₃ soln. (30 ml). FC (silica gel, 13.5 × 2 cm, toluene/AcOEt 1:1 → 1:1 + 2% MeOH → 1:1 + 3% MeOH): 293 mg (65%) of **22**. Amorphous solid. UV (CH₂Cl₂): 331 (4.67), 318 (sh, 4.62), 272 (sh, 4.61), 267 (4.63). ¹H-NMR (CDCl₃): 8.72, 8.18–8.13 (*s*, *m*, H–C(8), H–C(2), 4 H *o* to NO₂); 7.43–7.35 (*m*, 4 H *m* to NO₂); 6.10 (*d*, H–C(1')); 5.67 (*m*, H–C(2')); 6.61, 6.15, 6.1–6.0 (*m*, 5 CH=C(retinyl)); 5.55 (*t*, H–C(14)(retinyl)); 4.70 (*d*, 2 H–C(15)); 4.60 (*m*, H–C(4')); 4.51, 4.40 (2*t*, 2 OCH₂CH₂); 4.45–4.25 (*m*, 2 H–C(5')); 3.15–3.0 (*m*, 2 OCH₂CH₂); 2.65–2.5 (*m*, H–C(3')), C(O)CH₂CH₂C(O); 2.2 (*dd*, H–C(3')); 2.0–1.8 (*m*, 2*s*, 2 H–C(4)(retinyl), 2 H–C(2)(retinyl), 2 H–C(3)(retinyl), Me–C(9)(retinyl), Me–C(13)(retinyl), Me–C(5)(retinyl)); 0.99 (*s*, 2 Me–C(1)(retinyl)). Anal. calc. for C₅₂H₅₉N₇O₁₄ (1006.1): C 62.08, H 5.91, N 9.75; found: C 62.08, H 5.90, N 9.46.

19. 3'-Deoxy-N⁶,2'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]-5'-O- $\{[2-(2,3\text{-di-O-palmitoylglycer-1-yloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**23**). As described in *Exper.* 3, with **19** [12] (383 mg, 0.6 mmol), succinic anhydride (72 mg, 0.72 mmol), and DMAP (95 mg, 0.78 mmol) in abs. CH₂Cl₂ (5 ml; 18 h). Then EDC (150 mg, 0.78 mmol) and 1,2-di-O-palmitoylglycerol [13] [15] (478 mg, 0.84 mmol; 2 h). Workup with CHCl₃ (100 ml) and 10% citric acid soln. (50 ml). FC (silica gel, 13 × 2 cm, toluene/AcOEt 1:1 → 1:1 + 5% MeOH): 575 mg (74%) of **23**. Amorphous solid. UV (CH₂Cl₂): 272 (sh, 4.50), 267 (4.54). ¹H-NMR (CDCl₃): 8.74, 8.22–8.15 (*s*, *m*, H–C(8), H–C(2), 4 H *o* to NO₂, NH); 7.47–7.39 (*m*, 4 H *m* to NO₂); 6.13 (*m*, H–C(1')); 5.69 (*m*, H–C(2')); 5.26 (*q*, H–C(2)(Glyc)); 4.65 (*m*, H–C(4')); 4.54–4.10 (*m*, 2*t*, 2 OCH₂CH₂, 2 H–C(1)(Glyc), 2 H–C(3)(Glyc), 2 H–C(5')); 3.19–3.09 (*m*, 2 OCH₂CH₂); 2.7–2.6 (*m*, C(O)CH₂CH₂C(O), H–C(3')); 2.4–2.25 (*m*, 2 CH₂(α)(Palm), H–C(3')); 1.8–1.6 (*m*, 2 CH₂(β)(Palm)); 1.25 (*m*, 48 H(Palm)); 0.88 (*t*, 6 H, Me(Palm)). Anal. calc. for C₆₇H₉₇N₇O₁₈ (1288.5): C 62.45, H 7.59, N 7.61; found: C 62.40, H 7.61, N 7.58.

20. 3'-Deoxy-2'-O- $\{[2-(2,3\text{-di-O-hexadecylglycer-1-yloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ -N⁶,2'-O-bis[2-(4-nitrophenyl)ethoxycarbonyl]adenosine (**24**). As described in *Exper.* 3, with **19** [12] (319 mg, 0.5 mmol), succinic anhydride (60 mg, 0.6 mmol), and DMAP (79 mg, 0.65 mmol) in abs. CH₂Cl₂ (5 ml; 18 h). Then EDC (125 mg, 0.65 mmol) and 1,2-di-O-hexadecylglycerol [13] [16] (352 mg, 0.65 mmol) in abs. CH₂Cl₂ (5 ml; 5 h). Workup with CHCl₃ (120 ml), sat. NaHCO₃ soln. (60 ml) and 10% citric acid soln. (80 ml). FC (silica gel, 13 × 2 cm, toluene/AcOEt 1:1 → 1:1 + 2% MeOH → 1:1 + 4% MeOH): 385 mg (61%) of **24**. Amorphous solid. UV (CH₂Cl₂): 272 (sh, 4.51), 267 (4.55). ¹H-NMR (CDCl₃): 8.74, 8.21–8.15 (*s*, *m*, H–C(8), H–C(2), 4 H *o* to NO₂, NH); 7.45 (2*d*, 4 H *m* to NO₂); 6.13 (*s*', H–C(1')); 5.69 (*m*, H–C(2')); 4.62 (*m*, H–C(4')); 4.54, 4.46 (2*t*, 2 OCH₂CH₂); 4.35–4.0 (*m*, 2 H–C(5'), H–C(2)(Glyc)); 3.65–3.4 (*m*, 2 H–C(1)(Glyc), 2 H–C(3)(Glyc), 2 CH₂(α)(EL)); 3.2–3.05 (*m*, 2 OCH₂CH₂); 2.7–2.55 (*m*, H–C(3'), C(O)CH₂CH₂C(O)); 2.25 (*m*, H–C(3')); 1.75, 1.5, 1.25 (3*m*, 56 H(EL)); 0.88 (*t*, 2 Me(EL)). Anal. calc. for C₆₇H₁₀₁N₇O₁₆ (1260.6): C 63.84, H 8.08, N 7.78; found: C 63.97, H 8.12, N 7.75.

21. 3'-Deoxy-5'-O- $\{[2-(2\text{-ambo-}\alpha\text{-tocopheryloxycarbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**25**). As described in *Exper.* 10, with **20** (400 mg, 0.35 mmol) and DBU (529 mg, 3.5 mmol) in abs. pyridine (7 ml; 17 h), then AcOH (209 mg, 3.5 mmol). Workup with CHCl₃ (100 ml) and sat. NaCl/NaHCO₃ soln. (2 × 40 ml). Purification by FC (silica gel, 10 × 2 cm, CHCl₃ → CHCl₃ + 5% MeOH → CHCl₃ + 7% MeOH): 248 mg (93%) of **25**. Amorphous solid. UV (CH₂Cl₂): 259 (4.11). ¹H-NMR (CDCl₃): 8.31, 8.10 (2*s*, H–C(8), H–C(2)); 6.05 (br. *s*, NH₂); 5.94 (*s*', H–C(1')); 5.6 (br. *s*, OH–C(2')); 4.67 (*m*, H–C(2'), H–C(4')); 4.40 (*m*, 2 H–C(5')); 3.0–0.75 (*m*, C(O)CH₂CH₂C(O), 2 H–C(3'), 49 H (tocopheryl)). Anal. calc. for C₄₃H₆₅N₅O₇ · 1/4H₂O (764.0): C 67.20, H 8.59, N 9.11; found: C 67.01, H 8.36, N 9.20.

22. 3'-Deoxy-5'-O- $\{[2-(\text{ergocalciferlyloxycarbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**26**). As described in *Exper.* 10, with **21** (250 mg, 0.22 mmol) and DBU (682 mg, 4.48 mmol) in abs. pyridine (9 ml; 17 h, N₂, darkness), then AcOH (269 mg, 4.48 mmol). Workup with CH₂Cl₂ (100 ml) and sat. NaHCO₃ soln. (2 × 50 ml). Purification by FC (silica gel, 5 × 2 cm, toluene/AcOEt 1:1 → 1:1 + 6% MeOH): 156 mg (95%) of **26**. Amorphous solid. UV (MeOH): 262 (4.39). ¹H-NMR (CDCl₃): 8.31, 8.13 (2*s*, H–C(8), H–C(2)); 6.20–5.95 (*dd*, H–C(6)(VitD₂), H–C(7)(VitD₂)); 5.95 (*d*, H–C(1')); 5.85 (br. *s*, NH₂); 5.60 (*m*, H–C(2')); 5.25 (*m*, H–C(22)(VitD₂)); H–C(23)(VitD₂); 5.15, 4.8 (2*m*, 2 H–C(19)(VitD₂)); 5.0 (*m*, H–C(3)(VitD₂)); 4.55 (*m*, H–C(4')); 4.1, 3.65 (2*m*, 2 H–C(5')); 3.1–0.6 (*m*, C(O)CH₂CH₂C(O), 2 H–C(3'), 36 H (VitD₂)). Anal. calc. for C₄₂H₅₉N₅O₆ (730.0): C 69.11, H 8.15, N 9.59; found: C 68.57, H 8.16, N 9.41.

23. 3'-Deoxy-5'-O- $\{[2-(\text{retinylloxycarbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**27**). As described in *Exper.* 10, with **22** (73 mg, 72.6 μmol) and DBU (111 mg, 0.726 mmol) in abs. pyridine (1.5 ml; 18 h, N₂, darkness), then AcOH (44 mg, 0.726 mmol). Workup with CHCl₃ (40 ml) and H₂O (15 ml). Purification by FC (silica gel, 9 × 1 cm,

$\text{CHCl}_3 \rightarrow \text{CHCl}_3 + 2\% \text{ MeOH} \rightarrow \text{CHCl}_3 + 5\% \text{ MeOH}$: 38 mg (84%) of **27**. Amorphous solid. UV (CH_2Cl_2): 330 (4.59), 320 (sh, 4.55), 259 (4.28). $^1\text{H-NMR}$ (CDCl_3): 8.29, 8.05 (2s, H-C(8), H-C(2)); 6.62, 6.3–5.98 (m, H-C(1'), H-C(2')), 5 CH=C(retinyl), OH-C(2''); 5.55 (t, H-C(14)(retinyl)); 4.78 (m, 2 H-C(15), H-C(4')); 4.4 (m, 2 H-C(5')); 2.65 (br. s, C(O) $\text{CH}_2\text{CH}_2\text{C(O)}$); 2.25 (m, 2 H-C(3')); 2.05–1.5 (m, 2s, 2 H-C(4)(retinyl), 2 H-C(2)(retinyl), 2 H-C(3)(retinyl), Me-C(9)(retinyl), Me-C(13)(retinyl), Me-C(5)(retinyl)); 1.00 (s, 2 Me-C(1)(retinyl)). A correct C,H,N anal. could not be obtained, due to some instability of **27** against light and oxidation.

24. 3'-Deoxy-5'-O- $\{[2-(2,3\text{-di-O-palmitoylglycer-1-yloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**28**). As described in *Exper. 10*, with **23** (400 mg, 0.31 mmol) and DBU (473 mg, 3.1 mmol) in abs. pyridine (6.2 ml; 18 h), then AcOH (186 mg, 3.1 mmol). Workup with CHCl_3 (100 ml) and sat. NaCl/ NaHCO_3 soln. (2×40 ml). Purification by FC (silica gel, 8.5×2 cm, $\text{CHCl}_3 \rightarrow \text{CHCl}_3 + 5\% \text{ MeOH} \rightarrow \text{CHCl}_3 + 8\% \text{ MeOH} \rightarrow \text{CHCl}_3 + 10\% \text{ MeOH}$): 275 mg (98%) of **28**. Amorphous solid. UV (CH_2Cl_2): 259 (4.12). $^1\text{H-NMR}$ (CDCl_3): 8.32, 8.08 (2s, H-C(8), H-C(2)); 5.92–5.83 (m, H-C(1'), NH_2); 5.27 (q, H-C(2)(Glyc)); 4.77 (m, H-C(2'), H-C(4')); 4.45–4.1 (m, 2 H-C(1)(Glyc), 2 H-C(3)(Glyc), 2 H-C(5')); 2.63 (m, C(O) $\text{CH}_2\text{CH}_2\text{C(O)}$); 2.35 (m, 2 $\text{CH}_2(\alpha)$ (Palm), 2 H-C(3')); 1.7–1.5 (m, 2 $\text{CH}_2(\beta)$ (Palm)); 1.25 (m, 48 H(Palm)); 0.88 (t, 2 Me(Palm)). Anal. calc. for $\text{C}_{49}\text{H}_{83}\text{N}_5\text{O}_{10}$ (902.2): C 65.23, H 9.27, N 7.76; found: C 65.22, H 9.18, N 7.55.

25. 3'-Deoxy-5'-O- $\{[2-(2,3\text{-di-O-hexadecylglycer-1-yloxy-carbonyl})\text{ethyl}]\text{carbonyl}\}$ adenosine (**29**). As described in *Exper. 10*, with **24** (303 mg, 0.24 mmol) and DBU (365 mg, 2.4 mmol) in abs. pyridine (4.8 ml; 18 h), then AcOH (144 mg, 2.4 mmol). Workup with CH_2Cl_2 (80 ml) and sat. NaHCO_3 soln. (2×40 ml). Purification by FC (silica gel, 9×2 cm, $\text{CHCl}_3 \rightarrow \text{CHCl}_3 + 1\% \text{ MeOH} \rightarrow \text{CHCl}_3 + 2\% \text{ MeOH} \rightarrow \text{CHCl}_3 + 5\% \text{ MeOH}$): 165 mg (79%) of **29**. Amorphous solid. UV (CH_2Cl_2): 259 (4.14). $^1\text{H-NMR}$ (CDCl_3): 8.32, 8.1 (2s, H-C(8), H-C(2)); 6.22 (br. s, NH_2); 6.03 (m, H-C(1')); 4.95 (m, H-C(2'), OH-C(2'')); 4.45–4.15 (m, H-C(4'), 2 H-C(5'), H-C(2)(Glyc)); 3.7–3.35 (m, 2 H-C(1)(Glyc), 2 H-C(3)(Glyc), 2 $\text{CH}_2(\alpha)$ (EL)); 2.67 (m, C(O) $\text{CH}_2\text{CH}_2\text{C(O)}$); 2.25 (m, 2 H-C(3')); 1.6–1.2 (m, 56 H(EL)); 0.88 ('t', 2 Me(EL)). Anal. calc. for $\text{C}_{49}\text{H}_{87}\text{N}_5\text{O}_8 \cdot 1/4\text{H}_2\text{O}$ (883.3): C 66.97, H 10.04, N 7.97; found: C 66.53, H 9.74, N 8.22.

REFERENCES

- [1] M. Pfister, H. Schirmeister, M. Mohr, S. Farkas, K. P. Stengele, T. Reiner, M. Dunkel, S. Gokhale, R. Charubala, W. Pfeleiderer, *Helv. Chim. Acta* **1995**, *78*, 1705.
- [2] K. G. Cunningham, S. A. Hutchinson, W. Manson, F. S. Spring, *J. Chem. Soc.* **1951**, 2299.
- [3] R. J. Suhadolnik, in 'Nucleoside Antibiotics', John Wiley & Sons, Inc., New York, 1970, p. 50.
- [4] H. Sawai, J. Imai, K. Lesiak, M. I. Johnston, P. F. Torrence, *J. Biol. Chem.* **1983**, *258*, 1671.
- [5] W. E. G. Müller, B. E. Weiler, R. Charubala, W. Pfeleiderer, L. Leserman, R. W. Sobol, R. J. Suhadolnik, H. C. Schröder, *Biochemistry* **1991**, *30*, 2027.
- [6] H. C. Schröder, R. J. Suhadolnik, W. Pfeleiderer, R. Charubala, W. E. G. Müller, *Int. J. Biochem.* **1992**, *24*, 55.
- [7] S. L. Beaucage, R. P. Iyer, *Tetrahedron* **1993**, *49*, 1925.
- [8] M. Manoharan, C. J. Guinasso, P. D. Cook, *Tetrahedron Lett.* **1991**, *32*, 7171.
- [9] D. W. Will, T. Brown, *Tetrahedron Lett.* **1992**, *33*, 2729.
- [10] M. Wasner, E. E. Henderson, R. J. Suhadolnik, W. Pfeleiderer, *Helv. Chim. Acta* **1994**, *77*, 1757.
- [11] D. G. Norman, C. B. Reese, *Synthesis* **1983**, 304.
- [12] R. Charubala, E. Uhlmann, F. Himmelsbach, W. Pfeleiderer, *Helv. Chim. Acta* **1987**, *70*, 2028.
- [13] H. Sigmund, Ph. D. Thesis, University of Konstanz, 1992.
- [14] F. Himmelsbach, B. S. Schulz, T. Trichtinger, R. Charubala, W. Pfeleiderer, *Tetrahedron* **1984**, *40*, 59.
- [15] R. J. Howe, T. Malkin, *J. Chem. Soc.* **1951**, 2663.
- [16] A. Hermetter, F. Paltauf, in 'Ether Lipids. Biochemical and Biomedical Aspects', Eds. M. K. Mangold and F. Paltauf, Academic Press, New York, 1983, p. 389.
- [17] R. W. Sobol, E. E. Henderson, N. Kon, J. Shao, E. Mordechai, N. L. Reichenbach, R. Charubala, H. Schirmeister, W. Pfeleiderer, R. J. Suhadolnik, *J. Biol. Chem.* **1995**, *270*, 5963.
- [18] E. E. Henderson, J.-Y. Yang, R.-D. Zhang, M. Bealer, *Virology* **1991**, *182*, 186.
- [19] E. Mordechai, N. Kon, E. E. Henderson, R. J. Suhadolnik, *Virology* **1995**, *206*, 913.