



S0040-4039(96)00041-X


## A New Approach to 1-Deoxy-Azasugars: Asymmetric Synthesis of Deoxymannojirimycin

Yi-Ming Xu, Wei-Shan Zhou\*

Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences,  
 354 Fenglin Lu, Shanghai 200032, CHINA

**Abstract:** A concise and flexible method based upon the kinetic resolution of the  $\alpha$ -furfurylamine derivative (3) for asymmetric synthesis of 1-deoxy-azasugars has been provided and deoxymannojirimycin (2) has been synthesized.

Naturally occurring polyhydroxylated piperidine alkaloids such as deoxynojirimycin (1) and deoxymannojirimycin (2) which can be regarded as 1-deoxy-azasugars have received much attention in recent years. Due to their biological activity as inhibitors of glycosidase enzyme from a variety of sources<sup>1</sup> and their potential value as therapeutic agents,<sup>2</sup> much effort has been directed toward the stereoselective synthesis of these alkaloids.<sup>3</sup> The majority of these syntheses lack flexibility and general applicability. Recently two flexible synthetic methods have been published using the dihydro-pyridone system as the building block but proceeding through a relative long route.<sup>2b,4</sup> Consequently development of a concise and flexible method for construction of these 1-deoxy-azasugars continues to be important in order to probe structure-activity correlations.




Our group has previously developed an efficient method for kinetic resolution of racemic  $\alpha$ -furfurylamine derivatives.<sup>5</sup> This reaction has been employed for syntheses of several types of alkaloids.<sup>6</sup> In this paper we wish to report the application of this kinetic resolution to synthesis of 1-deoxy-azasugars.

Starting from 2-furaldehyde, the  $\alpha$ -furfurylamine derivative (3) could be prepared in 40% overall yield in three steps.<sup>5b</sup> Kinetic resolution of (3) in the reported procedure<sup>5</sup> yielded (2R, 6R)-(4) and (-)-(3) which could be converted to enantiomer of (4) by treatment with m-CPBA. Both (4) and its enantiomer can be applied for the synthesis of 1-deoxy-azasugars. Evidently our strategy has a potential to generate all isomers of 1-deoxy-azasugars. Here we describe the synthesis of deoxymannojirimycin (2) as an example of this strategy.

As depicted in **Scheme 1**, treatment of (4) with triethyl orthoformate generated (5). Reduction of (5) with NaBH<sub>4</sub> and CeCl<sub>3</sub>·7H<sub>2</sub>O afforded solely the alcohol (6) in which the configuration of hydroxyl group was assigned by 2D-NOESY spectroscopic analysis and this configuration was found to be opposite to that desired for the target molecule. Inversion of this configuration was successfully achieved by employing Mitsunobu reaction. Removal of ethoxy group of (7) by NaBH<sub>4</sub>, then Sharpless asymmetric dihydroxylation<sup>7</sup> led exclusively to the diol (9) which was protected to give diacetate (10). Attempts to selectively remove

methyl group in (10) with  $(\text{CH}_3)_3\text{SiI}$  failed, whereas demethylation proceeded smoothly by using  $\text{BBr}_3$ . Finally deprotection of (11) by sodium naphthalide and chromatography on column of Dowex-50 ( $\text{H}^+$ ) gave deoxymannojirimycin (2). m.p.  $185^\circ\text{C}$ ;  $[\alpha]_{\text{D}}^{20} -27^\circ$  (c 0.1 in MeOH); [lit.<sup>8</sup> m.p.  $185\text{--}187^\circ\text{C}$ ,  $[\alpha]_{\text{D}}^{20} -26.7^\circ$  (0.12 in MeOH)]. The proton NMR spectrum and the mass spectrum of (2) were identical with those of authentic sample.<sup>9</sup>



**Scheme 1. Reagents and conditions:** a)  $\text{Ti}(\text{O}i\text{Pr})_4$ ,  $L\text{-(+)-DIPT}$ , TBHP, silica gel,  $\text{CaH}_2$ ,  $\text{CH}_2\text{Cl}_2$ ,  $25^\circ\text{C}$ , 3days; b)  $\text{HC}(\text{OEt})_3$ ,  $\text{BF}_3\cdot\text{OEt}_2$ , THF,  $0^\circ\text{C}$  (76.5%); c)  $\text{NaBH}_4$ ,  $\text{CeCl}_3\cdot 7\text{H}_2\text{O}$ ,  $-30^\circ\text{C}$  (72.3%); d) DEAD-TPP,  $\text{PhCOOH}$ , THF, r.t. (91.7%); e)  $\text{NaBH}_4$ ,  $\text{HCOOH}$ ,  $0^\circ\text{C}$  (86.7%); f)  $(\text{DHQ})_2\text{-PHAL}$ ,  $\text{OsO}_4$ ,  $\text{K}_3\text{Fe}(\text{CN})_6$ ,  $\text{K}_2\text{CO}_3$ ,  $t\text{-BuOH}$ , r.t., 2days (84.8%); g)  $\text{Ac}_2\text{O}$ , Pyridine, DMAP, r.t. (100%); h)  $\text{BBr}_3$ ,  $\text{CH}_2\text{Cl}_2$ ,  $-78^\circ\text{C}$  (71.6%); i)  $\text{Na/Naphthalene}$ , DME,  $-60^\circ\text{C}$  (50.7%)

In summary, a new strategy for enantioselective synthesis of 1-deoxy-azasugars based upon kinetic resolution of the  $\alpha$ -furfurylamine derivative (3) has been proposed.<sup>10</sup> By this strategy, work on deoxynojirimycin (1) and other analogues is in progress.

**Acknowledgment:** We thank the National Natural Science Foundation of China and the State Key Laboratory of Bio-Organic & Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences for support of this work. We also thank Dr. Wang Zhi-Min for his constructive discussion.

## References and Notes

- a) Fuhrmann, U.; Bause, E.; Legler, G.; Hloegh, H. *Nature* **1994**, 307, 755 b) Fleet, G. W. J.; Fellows, L. E.; Smith, P. W. *Tetrahedron* **1987**, 43, 979 and references cited therein.
- a) Look, G. C.; Fotsch, C. H.; Wong, C. H. *Acc. Chem. Res.* **1993**, 26, 182 b) Cook, G. R.; Beholz, L. G.; Stille, J. R. *J. Org. Chem.* **1994**, 59, 3575 and references cited therein.
- a) Hughes, A. B.; Rudge, A. J. *Natural Product Reports* **1994**, 135 b) Furneaux, R. H.; Tyler, P. C.; Whitehouse, L. A. *Tetrahedron Lett.* **1993**, 34, 3613 and several references cited therein.
- Altenbach, H. j.; Himmeldirk, K. B. *Tetrahedron Asymmetry* **1995**, 6, 1077
- a) Zhou, W. S.; Lu, Z. H.; Wang, Z. M. *Tetrahedron Lett.* **1991**, 32, 1467 b) Zhou, W. S. Lu, Z. H.; Wang, Z. M. *Tetrahedron* **1993**, 49, 2641
- a) Lu, Z. H.; Zhou, W. S. *J. Chem. Soc. Perkin Trans. I* **1993**, 593 b) Lu, Z. H.; Zhou, W. S. *Tetrahedron* **1993**, 49, 4659 c) Zhou, W. S.; Xie, W. G.; Lu, Z. H.; Pan, X. F. *Tetrahedron Lett.* **1995**, 36, 1291
- Sharpless, K. B.; Amberg, W.; Bennani, Y. L.; Crispino, G. A.; Hartung, J.; Jeong, K.-S.; Kwong, H.-L.; Morikawa, K.; Wang, Z.-M.; Xu, D.; Zhang, X.-L. *J. Org. Chem.* **1992**, 57, 2768
- Fleet, G. W. J.; Smith, P. W. *Tetrahedron Lett.* **1985**, 26, 1469
- Park, K. H.; Yoon, Y. J.; Lee, S. G. *J. Chem. Soc. Perkin Trans. I* **1994**, 2621
- As we completed the synthesis of (2), H. J. Altenbach and R. Wischnat reported the synthesis of protected racemic mannojirimycin from a related furfurylamine derivative: *Tetrahedron Lett.* **1995**, 36, 4983.