

Article

Continuous Flow Alcoholysis of Dialkyl *H*-Phosphonates with Aliphatic Alcohols

Erika Bálint * , Ádám Tajti , Nóra Tóth and György Keglevich * 

Department of Organic Chemistry and Technology, Budapest University of Technology and Economics, 1521 Budapest, Hungary; tajti.adam@mail.bme.hu (Á.T.); toth.nora@mail.bme.hu (N.T.)

* Correspondence: ebalint@mail.bme.hu (E.B.); gkeglevich@mail.bme.hu (G.K.); Tel.: +36-1-463-3653 (E.B.)

Received: 14 June 2018; Accepted: 1 July 2018; Published: 3 July 2018



Abstract: The continuous flow alcoholysis of dialkyl *H*-phosphonates by aliphatic alcohols in the absence of a catalyst was elaborated using a microwave (MW) reactor equipped with a flow cell. By the precise control of the reaction conditions, the synthesis could be fine-tuned towards dialkyl *H*-phosphonates with two different and with two identical alkyl groups. In contrast to the “traditional” batch alcoholysis, flow approaches required shorter reaction times, and the products became available at a larger scale.

Keywords: dialkyl *H*-phosphonates; alcoholysis; transesterification; microwave; continuous flow reactor

1. Introduction

Flow chemistry induced an inevitable revolution in the field of chemical transformations [1]. In modern chemistry, reactions have to be strictly controlled, which requires more and more precise synthetic techniques. In contrast to the “traditional” batch reactions, flow approaches create a significantly different processing environment, which enables a more efficient control of the reaction conditions (such as the reaction time or temperature). Due to this, flow protocols may be significantly better with respect to purity, selectivity and yields, as compared to batch approaches [2].


In microwave (MW) chemistry, the continuous flow technique represents a special importance. Although applications of the MW technique proved to be useful in many chemical transformations, the scale-up of MW-assisted reactions means a challenge due to the limited geometry of the MW devices [3–6]. One possibility to solve this problem is the use of continuous flow MW reactors, where the reaction mixture flows through an irradiated flow cell. The MW unit can be of a similar size used in batch mode. During the last decade, the advantages of the applications of continuous flow MW reactors were reported in certain transformations; however, in most cases, the usefulness of the occasional devices may be questionable, as non-professional MW reactors do not allow reproductions, and what is more important, the temperatures were not reported [7,8].

Esters represent a fundamental family among organic compounds. Esters of carboxylic acids may be important intermediates, solvents or products in organic chemistry [9].

The most common preparations of simple carboxylic esters comprise the acid-catalyzed reaction of a carboxylic acid with alcohol (Fischer esterification) and the reaction of an ester with alcohol (alcoholysis) [10]. Continuous flow Fischer esterifications may be carried out in systems containing a packed catalyst bed [11–20] or a heated coil [21,22], as well as in microreactors [23]. In a few cases, the esterifications were performed in continuous flow MW systems based on professional MW reactors [24–26] or in household MW ovens [27,28]. The continuous flow alcoholysis of carboxylic esters may also be performed in reactors equipped with a catalyst bed [29] or a heated coil [21,30]. The alcoholysis is also of great importance in biodiesel production, the area of which was recently summarized by Lee and co-workers [31].


Organophosphorus esters are also of great importance [32]. Dialkyl esters of phosphorous acid (dialkyl *H*-phosphonates) are widely applied building blocks in syntheses [33]. They are important starting materials of the Kabachnik–Fields condensations and the aza-Pudovik reactions [34] (towards α -aminophosphonates), the Pudovik reactions [35] (resulting in the formation of α -hydroxyphosphonates) and further organophosphorus transformations, such as the Hirao reaction [36] and the phospho-Michael addition [37]. Dialkyl *H*-phosphonates bearing different alkoxy groups on the phosphorus atom are valuable intermediates for P-chiral organophosphorus derivatives [38,39].

The industrial synthesis of dialkyl *H*-phosphonates is comprised of the reaction of phosphorus trichloride with alcohols (Scheme 1) [40–45]. Although this transformation is convenient, efficient and may be scaled up, it requires a solvent, and the liberating HCl decreases the atom efficiency. It is also a disadvantage that the synthesis of derivatives with different alkyl groups is not possible.




Scheme 1. Synthesis of dialkyl *H*-phosphonates from phosphorus trichloride.

Another possibility for the preparation of dialkyl *H*-phosphonates is alcoholysis (Scheme 2), which provides the products in good yields [46–54]. The only by-product is the leaving alcohol. The relatively high temperature and the need for the high excess of the alcohol mean disadvantages.



Scheme 2. Synthesis of dialkyl *H*-phosphonates by alcoholysis.

Besides the two main routes mentioned above, there are a few less important methods for the synthesis of dialkyl *H*-phosphonates. Such protocols are the oxidative reaction of elemental phosphorus with alcohols (Scheme 3a) [55–60] and the Ni-catalyzed oxidation of hypophosphorous esters (Scheme 3b) [61].



Scheme 3. Miscellaneous synthesis of dialkyl *H*-phosphonates.

The synthesis of dialkyl-*H*-phosphonates bearing different alkyl groups is not easy and requires special methods (Table 1).

Table 1. Synthetic methods for the preparation of dialkyl *H*-phosphonates bearing two different alkyl groups.

Entry	Reaction	Average Yield (%)	Flow Compatible	Ref.
1	$\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{OR}^1 \end{array} \xrightleftharpoons{\text{R}^2\text{OH}} \begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{OR}^2 \end{array}$	30–50	+	[46,50,52]
2	$\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{OR}^1 \end{array} \xrightarrow[\text{X} = \text{OH}, \text{HSO}_4]{\text{Bu}_4\text{N}^{\oplus} \text{X}^{\ominus}} \left[\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{O}^{\ominus} \oplus \text{NBu}_4 \end{array} \right] \xrightarrow{\text{R}^2\text{I}} \begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{OR}^2 \end{array}$	50–70	–	[62,63]
3	$\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{OR}^1 \end{array} \xrightarrow{\text{NH}_4\text{OH}} \left[\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{O}^{\ominus} \oplus \text{NH}_4 \end{array} \right] \xrightarrow[\text{pyridine}]{\text{R}^2\text{OH}, \text{tBuC(O)Cl}} \begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{OR}^2 \end{array}$	50–85	–	[64]
4	$\begin{array}{c} \text{OR}^1 \\ \\ \text{Cl}-\text{P}-\text{Cl} \\ \\ \text{Cl} \end{array} \xrightarrow[\text{- 2 HCl}]{\text{2 R}^2\text{OH}, \text{pyridine}} \left[\begin{array}{c} \text{OR}^1 \\ \\ \text{R}^2\text{O}-\text{P}-\text{OR}^2 \\ \\ \text{OR}^2 \end{array} \right] \xrightarrow[\text{- R}^2\text{Cl}]{\text{HCl}} \begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{P}-\text{OR}^1 \\ \\ \text{OR}^2 \end{array}$	30–60	–	[65]

Phosphites with different alkyl groups may be synthesized by the partial alcoholysis of dialkyl *H*-phosphonates (Table 1, Entry 1). The reaction conditions must be controlled strictly to avoid complete transesterification.

Dialkyl *H*-phosphonates with two different alkyl groups may be obtained by the controlled O–C cleavage of dialkyl *H*-phosphonates (Table 2, Entries 2 and 3). In the first step of the first method, a phosphite-ammonium salt is prepared, which is then alkylated with alkyl iodides (Table 2, Entry 2), or with alcohols in the presence of pivaloyl chloride and pyridine (Table 2 entry 3) to furnish the target compounds selectively. A serious drawback of these transformations is the low atom efficiency.

Table 2. Alcoholysis of dialkyl *H*-phosphonates in a batch microwave (MW) reactor [33].

$$\begin{array}{c}
 \text{R}^1\text{O}-\text{P}(=\text{O})(\text{H})-\text{OR}^1 \\
 \mathbf{A}
 \end{array}
 + \text{R}^2\text{OH}
 \xrightarrow[\text{- R}^1\text{OH}]{\text{Batch MW, T, t}}
 \begin{array}{c}
 \text{R}^2\text{O}-\text{P}(=\text{O})(\text{H})-\text{OR}^1 \\
 \mathbf{B}
 \end{array}
 +
 \begin{array}{c}
 \text{R}^2\text{O}-\text{P}(=\text{O})(\text{H})-\text{OR}^2 \\
 \mathbf{C}
 \end{array}$$

$\text{R}^2 = \text{Me, Et, } ^i\text{Pr, } ^n\text{Bu, } ^n\text{Pent}$
 $\text{R}^1 = \text{Me, Et}$

Entry	R ¹	R ²	R ² OH (equiv)	T (°C)	t (min)	Composition (%) ^a		
						A	B	C
1	Me	Et	25	100	120	22	56	22
2	Me	Et	50	175	40	0	4	96
3	Me	ⁿ Bu	25	125	60	0	40	60
4	Me	ⁿ Bu	50	150	60	0	0	96
5	Et	Me	25	125	120	49	38	13
6	Et	Me	50	175	40	0	21	79
7	Et	ⁱ Pr	25	125	60	26	57	17
8	Et	ⁱ Pr	50	175	40	0	6	94
9	Et	ⁿ Bu	25	125	60	25	54	21
10	Et	ⁿ Bu	50	175	40	0	2	98
11	Et	ⁿ Pent	25	125	60	3	52	45
12	Et	ⁿ Pent	50	175	40	0	8	92

^a Analyzed by GC. Bold numbers indicate the target compounds in the reactions.

Dialkyl *H*-phosphonates bearing different alkoxy groups can also be obtained in the reaction of alkyl dichlorophosphites and alcohols. The primarily formed trialkyl phosphite is cleaved by a part of the hydrochloric acid formed (Table 2, Entry 4).

From these synthetic methods, the alcoholysis may be flow compatible. This solvolytic reaction may be carried out without any catalyst by fine-tuning the reaction conditions, to afford mixed or fully-transesterified dialkyl phosphites as the predominant component.

The alcoholysis of dialkyl *H*-phosphonates in a batch MW reactor was investigated by us previously (Table 2) [52]. The reaction of dialkyl *H*-phosphonates (**A**) with aliphatic alcohols under mild conditions (lower alcohol excess and lower temperature) afforded the derivatives with different alkyl groups (**B**) as the main products, while in the case of applying higher excess of alcohol and higher temperature, the fully-transesterified dialkyl *H*-phosphonates (**C**) predominated in the mixture.

In this paper, we aimed at elaborating an MW-assisted continuous flow method for the alcoholysis of dialkyl *H*-phosphonates. Our purpose was to find the optimum conditions for the formation of dialkyl *H*-phosphonates bearing different alkyl groups and for the fully-transesterified phosphites in a flow MW reactor. Furthermore, we wished to prepare new dialkyl *H*-phosphonates with two different alkyl groups, which may be valuable building blocks of chiral organophosphorus compounds.

2. Results and Discussion

In the first case, the alcoholysis of dimethyl *H*-phosphonate (DMP) (**1**) with *n*-butanol was investigated in the absence of any catalyst, in a CEM[®] (Matthews, NC, USA) MW reactor equipped with a commercially available CEM[®] continuous flow cell (Figures 1 and 2). The mixture of DMP (**1**) and a 25-fold excess of *n*-butanol was fed into the reactor by an HPLC pump at a flow rate of 0.1–1.4 mL/min (corresponding to residence times of 60–5 min, respectively). The temperature was monitored and controlled by an IR sensor. The mixture leaving the reactor was cooled down using a spiral-like cooler and was passed through a back pressure regulator operating at 250 psi. Consecutive fractions of the leaving mixture were analyzed by GC measurements in order to determine the composition and to identify the stationary operation.




Figure 1. Schematic drawing of the continuous flow system developed.




Figure 2. Sketch of the continuous flow cell.

The reaction conditions and the results are summarized in Table 3. To find the optimum conditions for the formation of *n*-butyl methyl *H*-phosphonate (nBMP) (**2a**) and *n*-dibutyl *H*-phosphonate (DnBP) (**3a**), the alcoholysis was carried out applying different temperatures and residence times. Performing the reaction at 100 °C, at a flow rate of 1.4 mL/min (with a residence time of 5 min), the conversion was only 26%, and the proportion of nBMP (**2a**) was 25%, along with 1% of DnBP (**3a**) (Table 3, Entry 1). Using a flow rate of 0.45 mL/min (with a residence time of 15 min), the starting DMP (**1**) was still the main component of the leaving mixture; however, the proportion of nBMP (**2a**) was 44% (Table 3, Entry 2). Increasing the residence time to 30 min (by decreasing the flow rate to 0.25 mL/min), the ratio of mixed *H*-phosphonate (**2a**) was 9% higher (53%) (Table 3, Entry 3). Applying a longer residence time of 45 or 60 min (flow rate of 0.15 or 0.10 mL/min), the proportion of phosphite **2a** was somewhat lower (50% or 47%), and the ratio of DnBP (**3a**) increased to 26% or 35%, respectively (Table 3, Entries 4 and 5). The maximum proportion of nBMP (**2a**) (53%) was obtained applying a reaction time of 30 min at 100 °C (Table 3, Entry 3). Increasing or decreasing the residence time at the same temperature resulted in lower ratio of product **2a** (Table 3, Entries 1, 2 and 4, 5, respectively). In a comparative thermal experiment at 100 °C at a residence time of 30 min, the conversion was only 15% (Table 3, Entry 6). Next, the effect of the reaction temperature was investigated at a residence time of 30 min (Table 3, Entries 6–8) (Figure 3). At 125 °C, the mixture was comprised of 7% of DMP (**1**), 43% of nBMP (**2a**) and

50% of DnBP (**3a**) (Table 3, Entry 7). Increasing the temperature to 150 °C, the conversion was complete, and the fully-transesterified product (**3a**) predominated in a proportion of 78% (Table 3, Entry 8). Performing the alcoholysis at 175 °C, the ratio of DnBP (**3a**) was 91% (Table 3, Entry 9). Carrying out the reaction at a longer residence time (45 min) and/or higher excess of the ⁿBuOH (50 equiv), the composition did not change. Applying common heating at 175 °C, the reaction was not complete, and the ratio of product **3a** was 8% lower, than under MW conditions (Table 3, Entry 10). It can be observed that the difference between MW-assisted and conventionally-heated experiments was significant at 100 °C, while significantly smaller at 175 °C. After column chromatography, nBMP (**2a**) was obtained in a yield of 48%, while DnBP (**3a**) was isolated in a yield of 88% (Table 3, Entries 3 and 9).

Table 3. Continuous flow alcoholysis of DMP with ⁿBuOH.

$$\text{MeO}-\text{P}(\text{O})(\text{MeO})-\text{H} + {}^n\text{BuOH} \xrightarrow[\text{- MeOH}]{\text{MW or } \Delta, \text{ T, } \tau} \text{}^n\text{BuO}-\text{P}(\text{O})(\text{MeO})-\text{H} + \text{}^n\text{BuO}-\text{P}(\text{O})(\text{}^n\text{BuO})-\text{H}$$

1 **2a** **3a**

Entry	Mode of Heating	Power (W)	T (°C) ^a	Flow Rate (mL/min) ^b	τ (min)	Conversion (%)	Composition (%) ^c			Yield (%) ^d
							1	2a	3a	
1	MW	22	100	1.4	5	26	74	25	1	-
2	MW	14	100	0.45	15	52	48	44	8	-
3	MW	10	100	0.25	30	65	35	53	12	48 (2a)
4	MW	8	100	0.15	45	76	24	50	26	-
5	MW	5	100	0.10	60	83	18	47	35	-
6	Δ	-	100	0.25	30	15	85	15	0	-
7	MW	18	125	0.25	30	93	7	43	50	-
8	MW	38	150	0.25	30	100	0	22	78	-
9	MW	59	175	0.25	30	100	0	9	91 ^{e,f}	88 (3a)
10	Δ	-	175	0.25	30	98	2	15	83	-

^a The pressure was 17 bar; ^b Based on the HPLC pump; ^c the mixtures from the stationary operation were analyzed by GC; ^d after column chromatography; ^e No change on longer residence time (45 min); ^f No change on higher excess (50 equiv) of ⁿBuOH. Bold numbers indicate the target compounds in the reactions.




Figure 3. Composition of the reaction mixtures of the MW-assisted alcoholysis of DMP with ⁿBuOH at different temperatures at a residence time of 30 min (Table 3 entries 3 and 7–9).

Comparative batch experiments were also carried out in the DMP (**1**)—ⁿBuOH model reaction at 100 °C. The alcohol excess (25 equiv) and the reaction time range studied (5–60 min) were the same

as in the flow approaches. The batch results are listed in Table 4. Similarly to the flow experiments, the composition was highly reaction time-dependent.

Table 4. Comparative experiments for the alcoholysis of dimethyl *H*-phosphonate (DMP) with ⁿBuOH in a batch MW reactor.

$$\begin{array}{c} \text{MeO} \\ \diagup \\ \text{P}=\text{O} \\ \diagdown \\ \text{MeO} \end{array} \text{H} + \text{}^n\text{BuOH} \xrightarrow[\text{- MeOH}]{\text{MW, } 100\text{ }^\circ\text{C, } t} \begin{array}{c} \text{}^n\text{BuO} \\ \diagup \\ \text{P}=\text{O} \\ \diagdown \\ \text{MeO} \end{array} \text{H} + \begin{array}{c} \text{}^n\text{BuO} \\ \diagup \\ \text{P}=\text{O} \\ \diagdown \\ \text{}^n\text{BuO} \end{array} \text{H}$$

Entry	t (min)	Conversion (%)	Composition (%) ^a		
			1	2a	3a
1	5	24	76	20	4
2	15	46	54	37	9
3	30	59	41	47	12
4	45	67	33	49	18
5	60	73	26	45	29

^a Based on GC.

Comparing the conversions of the flow and batch processes, the two plots show a similar shape; however, after a reaction time of 15 min, the batch conversions were 5–10% lower than the flow results (Figure 4).




Figure 4. Reaction/residence time-dependence of the conversion of the alcoholysis of DMP with ⁿBuOH at 100 °C.

A more significant difference may be observed if the ratio of product 2a is examined separately (Figure 5). Maximum points are present on both curves: at 30 min and 53% in the flow experiments and at 45 min 49% in the case of the batch approaches. Based on the comparison, the flow reaction reaches the optimum point of phosphite 2a under a shorter time, with a higher ratio.

The next model was the continuous flow alcoholysis of diethyl *H*-phosphonate (DEP) (**4**) (Table 6). Based on the experiences from the transesterification of DMP (**1**), the reactions were carried out using 25 equivalents of the alcohols at a flow rate of 0.25 mL/min (or at a residence time of 30 min). First, the alcoholysis of DEP (**4**) by *n*BuOH was investigated. Performing the reaction at 100 °C, the unreacted DEP (**4**) was the main component along with 34% of *n*-butyl ethyl *H*-phosphonate (nBEP) (**5a**) and 8% of DBP (**3a**) (Table 6, Entry 1). At a higher temperature of 125 °C, the ratio of nBEP (**5a**) was 42%, and after column chromatography, compound **5a** could be isolated in a yield of 38% (Table 6, Entry 2). Increasing the temperature to 150 °C and 175 °C, the proportion of **5a** was decreased, while that of DBP (**3a**) increased (Table 6, Entries 3 and 4). In the latter case, the alcoholysis was complete, and the DBP (**3a**) was obtained in a yield of 85% (Table 6, Entry 4). The transesterification of DEP (**4**) by *n*PrOH at 125 °C took place similarly as the alcoholysis by *n*BuOH, and the *n*-propyl ethyl *H*-phosphonate (PrEP) (**5b**) was obtained in a yield of 36% (Table 6, Entries 2 and 5). At 175 °C, the conversion was not complete, and the resulting mixture contained only 74% of DPrP (**3b**) besides 4% of unreacted DEP (**4**) and 22% of PrEP (**5b**) (Table 6, Entry 6). To obtain a higher proportion of product **3b**, the temperature was increased to 200 °C. Hence, phosphite **3b** was formed in 84% (Table 6, Entry 7). In the case of *i*BuOH, the tendency was similar, and the highest amount of mixed phosphonate (**5c**) was formed at 125 °C, while at 175 and 200 °C, the fully-transesterified product (**3c**) predominated (Table 6, Entries 8–10). Using PentOH at 125 °C, the pentyl ethyl *H*-phosphonate (PeEP) (**5d**) was the main component in the departing mixture, from which it was isolated in a yield of 40% (Table 6, Entry 11). For the preparation of dipentyl *H*-phosphonate (DPeP) (**3d**), the application of 175 °C was enough, and compound **3d** was obtained in a yield of 80% (Table 6, Entry 12).

Table 6. Continuous flow alcoholysis of DEP with alcohols.

Entry	R	Power (W)	T (°C) ^a	Composition (%) ^b			Yield (%) ^c
				4	5	3	
1	<i>n</i> Bu	10	100	58	34	8	-
2		17	125	44	42	14	38 (5a)
3		38	150	7	32	61	-
4		57	175	0	11	89 ^{d,e}	85 (3a)
5	<i>n</i> Pr	18	125	51	41	8	36 (5b)
6		54	175	4	22	74	-
7		83	200	2	14	84 ^{d,e}	78 (3b)
8	<i>i</i> Bu	21	125	21	40	39	36 (5c)
9		63	175	3	26	71	-
10		105	200	0	16	84 ^{d,e}	77 (3c)
11	<i>n</i> Pent	10	125	31	44	25	40 (5d)
12		39	175	2	13	85 ^{d,e}	80 (3d)

^a The pressure was 17 bar; ^b the mixtures from the stationary operation were analyzed by GC; ^c after column chromatography; ^d no change upon longer residence time (45 min); ^e no change upon higher excess (50 equiv) of *n*BuOH. Bold numbers indicate the target compounds in the reactions.

The catalyst-free alcoholysis of dialkyl *H*-phosphonates with various aliphatic alcohols was carried out efficiently in a continuous flow MW reactor. Applying a residence time of 30 min at different temperatures, the alcoholysis could be fine-tuned towards the dialkyl *H*-phosphonates with two

different or with two identical alkyl groups. In the case of the reaction studied, the dimethyl phosphite proved to be the more reactive starting material. After almost complete conversions, the $(RO)_2P(O)H$ species formed by transesterification were obtained in yields above 85%. The ratio of $(RO)(R'O)P(O)H$ derivatives with two different alkyl groups was around 50% in all mixtures, affording the target compounds in a yield range of ca. 40–50%. By the method developed, a series of mixed dialkyl *H*-phosphonates, which are valuable building blocks for the preparation of chiral organophosphorus derivatives, was synthesized.

3. Materials and Methods

3.1. General

GC measurements were performed on an HP5890 Series 2 GC-FID chromatograph, using a $15\text{ m} \times 0.18\text{ mm}$ Restek, Rtx-5 column with a film layer of $0.20\text{ }\mu\text{m}$. The temperature of the column was initially held at $40\text{ }^\circ\text{C}$ for 1 min, followed by programming at $25\text{ }^\circ\text{C}/\text{min}$ up to $300\text{ }^\circ\text{C}$ and a final period at $300\text{ }^\circ\text{C}$ (isothermal) for 10 min. The temperature of the injector was $290\text{ }^\circ\text{C}$ and of the FID detector was $300\text{ }^\circ\text{C}$. The carrier gas was N_2 .

GC-MS measurements were performed on an Agilent 6890 N-GC-5973 N-MSD chromatograph, using a $30\text{ m} \times 0.25\text{ mm}$ Restek, Rtx-5SILMS column with a film layer of $0.25\text{ }\mu\text{m}$. The initial temperature of the column was $45\text{ }^\circ\text{C}$ for 1 min, followed by programming at $10\text{ }^\circ\text{C}/\text{min}$. up to $310\text{ }^\circ\text{C}$ and a final period at $310\text{ }^\circ\text{C}$ (isothermal) for 17 min. The temperature of the injector was $250\text{ }^\circ\text{C}$. The carrier gas was He, and the operation mode was splitless.

High resolution mass spectrometric measurements were performed using a TripleTOF 5600+ mass spectrometer in positive electrospray mode.

The ^{13}C - and ^1H -NMR spectra were obtained in CDCl_3 solution on a Bruker DRX-500 spectrometer operating at 125.7 and 500.1 MHz, respectively. The ^{13}C and ^1H chemical shifts are referred to TMS. ^{31}P -NMR spectra were obtained on a Bruker AV-300 spectrometer at 121.5 MHz. Chemical shifts are downfield relative to 85% H_3PO_4 .

3.2. Equipment

The continuous flow reactions were performed in a self-developed continuous flow system comprising a 300-W CEM[®] Discover-focused microwave reactor equipped with a CEM[®] 10-mL Flow Cell Accessory continuous flow unit (irradiated volume 7 mL), a Gilson 305 HPLC pump, an HPLC backpressure regulator with a 250-psi (17.2 bar) cartridge and a cooler. Teflon[®] (Wilmington, DE, USA) PFA tubes with an outside diameter of 0.125" (3.175 mm) and an inside diameter of 0.064" (1.575 mm) were used. The exact lengths and volumes of each tube part are shown in Figure 6. All of the tubes, screws and ferrules applied were fully compatible with a regular HPLC system.




Figure 6. Design parameters of the continuous flow system.

3.3. General Procedure for the Continuous Flow Alcoholysis of Dialkyl H-Phosphonates

A mixture of 25.0 mmol of the dialkyl H-phosphonate (2.7 mL of dimethyl H-phosphonate, 3.9 mL of diethyl H-phosphonate) and 25 equivalents, 0.625 mol of alcohol (47 mL of *n*-propanol, 57 mL of *n*-butanol, 58 mL of *i*-butanol, 68 mL of *n*-pentanol) was homogenized by stirring for 5 min at 25 °C. The reactor was flushed with 20 mL of the mixture with a flow rate of 10 mL/min at 25 °C and 17 bar. Next, under the same pressure, the flow rate was set to the desired value, and the vessel was irradiated with a power of 45–300 W for 3–5 min, until the temperature reached the desired value, then the power was controlled automatically by the software of the MW reactor. The flow rate and the necessary power at stationary operation can be seen in Tables 3, 5 and 6. The operation was regarded as the steady state on the basis of the results of the GC measurements. The products were separated by column chromatography using silica gel as the absorbent and ethyl acetate as the eluent. Yields were calculated on the basis of the weights obtained after separation and evaporation taking into consideration the quantity of the dialkyl H-phosphonate fed in during a given time. The ³¹P-NMR and mass data for the dialkyl H-phosphonates are listed in Table 7.

3.4. General Procedure for the Comparative Batch Experiments

A mixture of 0.5 mmol (0.05 mL) of dimethyl H-phosphonate and 25 equivalents, 12.5 mmol (1.1 mL) of *n*-butanol was heated at 100 °C in a vial in the MW reactor for 5–60 min as shown in Table 4. The volatile components were removed in vacuum, and the residual oil was analyzed by GC.

Methyl propyl H-phosphonate (2b): Yield: 42%; ³¹P-NMR and HRMS (see Table 7); ¹³C-NMR (CDCl₃) δ 10.0 (CH₃CH₂), 23.8 (*J* = 6.2, CH₂), 52.0 (*J* = 5.8, CH₃O), 67.1 (*J* = 6.0, CH₂O); ¹H-NMR (CDCl₃) δ 0.98 (t, *J*_{HH} = 7.4, 3H, CH₃CH₂), 1.65–1.82 (m, 2H, CH₂), 3.78, (d, *J*_{PH} = 11.9, 3H, CH₃O) 3.97–4.12 (m, 2H, CH₂O), 6.79 (d, *J*_{PH} = 695.4, 1H, PH).

i-Butyl methyl H-phosphonate (2c): Yield: 46%; ³¹P-NMR and HRMS (see Table 7); ¹³C-NMR (CDCl₃) δ 18.6 (*J* = 3.1, CH₃CH), 29.2 (*J* = 6.4, CH), 51.9 (*J* = 5.7, CH₃O), 71.8 (*J* = 6.3, CH₂O); ¹H-NMR (CDCl₃) δ 0.89 (d, *J*_{HH} = 6.8, 6H, CH₃CH), 1.84–1.96 (m, 1H, CH), 3.71 (d, *J*_{PH} = 11.9, 3H, CH₃O) 3.74–3.83 (m, 2H, CH₂O), 6.72 (d, *J*_{PH} = 695.8, 1H, PH).

Methyl pentyl H-phosphonate (2d): Yield: 50%; ³¹P-NMR and HRMS (see Table 7); ¹³C-NMR (CDCl₃) δ 13.9 (CH₃CH₂), 22.2 (CH₃CH₂), 27.6 (CH₃CH₂CH₂), 30.1 (*J* = 6.2, CH₂CH₂O), 51.9 (*J* = 5.7, CH₃O), 67.0 (*J* = 6.1, CH₂O); ¹H-NMR (CDCl₃) δ 0.91 (t, *J* = 6.9, 3H, CH₃CH₂), 1.29–1.44 (m, 4H, CH₂CH₂CH₃), 1.63–1.78 (m, 2H, CH₂CH₂O), 3.78, (d, *J*_{PH} = 11.9, 3H, CH₃O) 3.99–4.17 (m, 2H, CH₂O), 6.79 (d, *J*_{PH} = 695.2, 1H, PH).

Ethyl propyl H-phosphonate (5b): Yield: 30%; ³¹P-NMR and HRMS (see Table 7); ¹³C-NMR (CDCl₃) δ 10.0 (CH₃CH₂CH₂), 16.3 (*J* = 6.1, CH₃CH₂O), 23.8 (*J* = 6.4, CH₃CH₂CH₂), 61.8 (*J* = 5.8, CH₂CH₂O), 67.3 (*J* = 6.0, CH₃CH₂O); ¹H NMR (CDCl₃) δ 0.96 (t, *J*_{HH} = 7.4, 3H, CH₃CH₂CH₂), 1.35 (t, *J*_{HH} = 7.1, 3H, CH₃CH₂O), 1.63–1.78 (m, 2H, CH₃CH₂CH₂), 3.94–4.07, (m, 2H, CH₂CH₂O) 4.07–4.21 (m, 2H, CH₃CH₂O), 6.80 (d, *J*_{PH} = 692.4, 1H, PH).

i-Butyl ethyl H-phosphonate (5c): Yield: 36%; ³¹P-NMR and HRMS (see Table 7); ¹³C-NMR (CDCl₃) δ 16.3 (*J* = 6.2, CH₃CH₂), 18.7 (*J* = 1.7, CH₃CH), 29.1 (*J* = 6.5, CH), 61.8 (*J* = 5.7, CH₃CH₂O), 71.6 (*J* = 6.2, CHCH₂O); ¹H NMR (CDCl₃) δ 0.96 (d, *J*_{HH} = 6.7, 6H, CH₃CH), 1.37 (t, *J*_{HH} = 7.1, 3H, CH₃CH₂), 1.89–2.06 (m, 1H, CH), 3.77–3.91, (m, 2H, CHCH₂O) 4.09–4.22 (m, 2H, CH₃CH₂O), 6.82 (d, *J*_{PH} = 692.7, 1H, PH).

Table 7. ^{31}P -NMR and mass data for the dialkyl *H*-phosphonates prepared.

Compound	δ_{P} (CDCl_3)	δ_{P} [Ref]	$[\text{M} + \text{H}]^+$ found	$[\text{M} + \text{H}]^+$ requires
2a	9.3 (695.0)	9.3 [52]	153.0684	153.0681
2b	9.3 (695.4)	-	139.0522	139.0523
2c	6.7 (695.7)	-	153.0676	153.0681
2d	6.8 (695.2)	-	167.0833	167.0837
3a	7.8	7.9 [66]	195.1153	195.1150
3b	7.9	7.8 [67]	167.0833	167.0837
3c	8.1	8.0 [67]	195.1145	195.1150
3d	7.8	8.1 [68]	223.1462	223.1463
5a	7.6 (693.0)	7.6 [52]	167.0840	167.0837
5b	8.6 (692.4)	-	153.0673	153.0681
5c	7.8 (692.8)	-	167.0832	167.0837
5d	7.7 (692.1)	7.7 [52]	181.0992	181.0994

4. Conclusions

In summary, a MW-assisted continuous flow method was developed for the catalyst-free alcoholysis of dialkyl *H*-phosphonates. By the optimization of the reaction parameters, the alcoholysis was fine-tuned towards dialkyl *H*-phosphonates with two different or with two identical alkyl groups. The selectivity of the reaction was similar in the flow and batch approaches; however, the continuous protocol offered several advantages. The dialkyl *H*-phosphonates with two different alkyl groups can be obtained at a shorter reaction time and in a higher ratio. Furthermore, a somewhat higher productivity (0.8–1.0 g/h) can be attained by the continuous flow method. Five new dialkyl *H*-phosphonates with different alkyl groups were isolated and characterized, which may be valuable building blocks in the synthesis of chiral organophosphorus compounds.

Author Contributions: E.B., Á.T. and G.K. conceived of and designed the experiments. Á.T. and N.T. performed the experiments. E.B. and G.K. contributed reagents/materials/analysis tools. E.B., Á.T. and G.K. wrote the paper.

Funding: The project was supported by the Hungarian Research Development and Innovation Fund (FK123961 and K119202), by the National Research, Development and Innovation Fund of Hungary in the frame of the FIEK_16-1-2016-0007 (Higher Education and Industrial Cooperation Center) project and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences (BO/00278/17/7) (E.B.).

Conflicts of Interest: The authors declare no conflict of interest.

References

- Plutschack, M.B.; Pieber, B.; Gilmore, K.; Seeberger, P.H. The Hitchhiker's guide to flow chemistry. *Chem. Rev.* **2017**, *117*, 11796–11893. [[CrossRef](#)] [[PubMed](#)]
- Glasnov, T. *Continuous-Flow Chemistry in the Research Laboratory*; Springer International Publishing: Basel, Switzerland, 2016; ISBN 978-3-319-32194-3.
- Bálint, E.; Keglevich, G. The Spread of the Application of the Microwave Technique in Organic Synthesis. In *Milestones in Microwave Chemistry*; Keglevich, G., Ed.; Springer: Basel, Switzerland, 2016; pp. 1–10. ISBN 978-3-319-30632-2.
- de la Hoz, A.; Loupy, A. (Eds.) *Microwaves in Organic Synthesis*, 3rd ed.; Wiley: Weinheim, Germany, 2012; ISBN 978-3-527-65131-3.
- Moseley, J.D. Microwave heating as a tool for process chemistry. In *Microwave Heating as a Tool for Sustainable Chemistry*; Leadbeater, N., Ed.; CRC Press: New York, NY, USA, 2010; pp. 105–147. ISBN 9781138111981.
- Kappe, C.O.; Stadler, A.; Dallinger, D. *Microwaves in Organic and Medicinal Chemistry*, 2nd ed.; Wiley: Weinheim, Germany, 2012; Volume 52, ISBN 978-3-527-33185-7.
- Estela, L.; Pouxb, M.; Benamaraa, N.; Polaerta, I. Continuous flow-microwave reactor: Where are we? *Chem. Eng. Process.* **2016**, *113*, 56–64. [[CrossRef](#)]
- Baxendale, I.; Hayward, J.; Ley, S. Microwave reactions under continuous flow conditions. *Comb. Chem. High Throughput Screen.* **2007**, *10*, 802–836. [[CrossRef](#)] [[PubMed](#)]

9. Riemenschneider, W.; Bolt, H.M. Esters, Organic. In *Ullmann's Encyclopedia of Industrial Chemistry*, 7th ed.; Wiley: Weinheim, Germany, 2005; pp. 245–265. ISBN 978-3-527-30673-2.
10. Otera, J.; Nishikido, J. (Eds.) *Esterification: Methods, Reactions, and Applications*, 2nd ed.; Wiley: Weinheim, Germany, 2010; ISBN 978-3-527-32289-3.
11. Woodcock, L.L.; Wiles, C.; Greenway, G.M.; Watts, P.; Wells, A.; Eyley, S. Enzymatic synthesis of a series of alkyl esters using novozyme 435 in a packed-bed, miniaturized, continuous flow reactor. *Biocatal. Biotransform.* **2008**, *26*, 501–507. [[CrossRef](#)]
12. Junior, I.I.; Flores, M.C.; Sutili, F.K.; Leite, S.G.F.; Miranda, L.S.D.M.; Leal, I.C.R.; de Souza, R.O.M.A. Fatty acids residue from palm oil refining process as feedstock for lipase catalyzed monoacylglycerol production under batch and continuous flow conditions. *J. Mol. Catal. B Enzym.* **2012**, *77*, 53–58. [[CrossRef](#)]
13. Sutili, F.K.; Ruela, H.S.; Leite, S.G.F.; Miranda, L.S.D.M.; Leal, I.C.R.; de Souza, R.O.M.A. Lipase-catalyzed esterification of steric hindered fructose derivative by continuous flow and batch conditions. *J. Mol. Catal. B Enzym.* **2013**, *85–86*, 37–42. [[CrossRef](#)]
14. Patil, N.G.; Benaskar, F.; Rebrov, E.V.; Meuldijk, J.; Hulshof, L.A.; Hessel, V.; Schouten, J.C. Scale-up of Microwave Assisted Flow Synthesis by Transient Processing through Monomode Cavities in Series. *Org. Process Res. Dev.* **2014**, *18*, 1400–1407. [[CrossRef](#)]
15. Sutili, F.K.; Ruela, H.S.; Nogueira, D.D.O.; Leal, I.C.R.; Miranda, L.S.D.M.; de Souza, R.O.M.A. Enhanced production of fructose ester by biocatalyzed continuous flow process. *Sustain. Chem. Process.* **2015**, *3*, 6. [[CrossRef](#)]
16. Okuno, Y.; Isomura, S.; Sugamata, A.; Tamahori, K.; Fukuhara, A.; Kashiwagi, M.; Kitagawa, Y.; Kasai, E.; Takeda, K. Convenient and Simple Esterification in Continuous-Flow Systems using g-DMAP. *ChemSusChem* **2015**, *8*, 3587–3589. [[CrossRef](#)] [[PubMed](#)]
17. Koreniuk, A.; Maresz, K.; Odrozek, K.; Jarzebski, A.B.; Mrowiec-Bialon, J. Highly effective continuous-flow monolithic silica microreactors for acid catalyzed processes. *Appl. Catal. A* **2015**, *489*, 203–208. [[CrossRef](#)]
18. Baek, H.; Minakawa, M.; Yamada, Y.M.A.; Han, J.W.; Uozumi, Y. In-Water and Neat Batch and Continuous-Flow Direct Esterification and Transesterification by a Porous Polymeric Acid Catalyst. *Sci. Rep.* **2016**, *6*, 25925. [[CrossRef](#)] [[PubMed](#)]
19. Furuta, A.; Fukuyama, T.; Ryu, I. Efficient Flow Fischer Esterification of Carboxylic Acids with Alcohols Using Sulfonic Acid-Functionalized Silica as Supported Catalyst. *Bull. Chem. Soc. Jpn.* **2017**, *90*, 607–612. [[CrossRef](#)]
20. Iemhoff, A.; Sherwood, J.; McElroy, C.R.; Hunt, A.J. Towards sustainable kinetic resolution, a combination of bio-catalysis, flow chemistry and bio-based solvents. *Green Chem.* **2018**, *20*, 136–140. [[CrossRef](#)]
21. Razzaq, T.; Glasnov, T.N.; Kappe, C.O. Continuous-Flow Microreactor Chemistry under High-Temperature/Pressure Conditions. *Eur. J. Org. Chem.* **2009**, 1321–1325. [[CrossRef](#)]
22. Archambault, C.M.; Leadbeater, N.E. A benchtop NMR spectrometer as a tool for monitoring mesoscale continuous-flow organic synthesis: Equipment interface and assessment in four organic transformations. *RSC Adv.* **2016**, *6*, 101171–101177. [[CrossRef](#)]
23. Gumel, A.M.; Annuar, M.S.M. *Thermomyces lanuginosus* lipase-catalyzed synthesis of natural flavor esters in a continuous flow microreactor. *3 Biotech* **2016**, *6*, 24. [[CrossRef](#)] [[PubMed](#)]
24. Tajti, Á.; Tóth, N.; Bálint, E.; Keglevich, G. Esterification of benzoic acid in a continuous flow microwave reactor. *J. Flow Chem.* **2017**, *8*, 11–19. [[CrossRef](#)]
25. Krull, M.; Moschaeuser, R. Continuous Method for Producing Esters of Aromatic Carboxylic Acids. U.S. Patent 0088918, 12 April 2012.
26. Cablewski, T.; Faux, A.F.; Strauss, C.R. Development and Application of a Continuous Microwave Reactor for Organic Synthesis. *J. Org. Chem.* **1994**, *59*, 3408–3412. [[CrossRef](#)]
27. Chen, S.-T.; Chiou, S.-H.; Wang, K.-T. Preparative scale organic synthesis using a kitchen microwave oven. *J. Chem. Soc. Chem. Commun.* **1990**, 807–809. [[CrossRef](#)]
28. Pipus, G.; Plazl, I.; Koloini, T. Esterification of benzoic acid in microwave tubular flow reactor. *Chem. Eng. J.* **2000**, *76*, 239–245. [[CrossRef](#)]
29. Asadi, M.; Hooper, J.F.; Lupton, D.W. Biodiesel synthesis using integrated acid and base catalysis in continuous flow. *Tetrahedron* **2016**, *72*, 3729–3733. [[CrossRef](#)]

30. Adeyemi, A.; Bergman, J.; Branalt, J.; Savmarker, J.; Larhed, M. Continuous Flow Synthesis under High-Temperature/High-Pressure Conditions Using a Resistively Heated Flow Reactor. *Org. Process Res. Dev.* **2017**, *21*, 947–955. [[CrossRef](#)]
31. Tran, D.-T.; Chang, J.-S.; Lee, D.-J. Recent insights into continuous-flow biodiesel production via catalytic and non-catalytic transesterification processes. *Appl. Energy* **2017**, *185*, 376–409. [[CrossRef](#)]
32. Corbridge, D.E.C. *Phosphorus: Chemistry, Biochemistry and Technology*, 6th ed.; CRC Press: New York, NY, USA, 2013; ISBN 978-1-439-84088-7.
33. Troev, K.D. *Chemistry and Application of H-Phosphonates*; Elsevier: Amsterdam, The Netherlands, 2006; ISBN 978-0-080-47649-0.
34. Bálint, E.; Tajti, Á.; Tripolszky, A. Synthesis of α -aminophosphonates by the Kabachnik–Fields reaction and by the Pudovik reaction. In *Organophosphorus Chemistry*; Keglevich, G., Ed.; Walter de Gruyter GmbH: Berlin, Germany, 2018; pp. 108–147. ISBN 978-3-11-053453-5.
35. Rádai, Z.; Kiss, N.Z.; Keglevich, G. Synthesis of α -hydroxyphosphonates, an important class of bioactive compounds. In *Organophosphorus Chemistry*; Keglevich, G., Ed.; Walter de Gruyter GmbH: Berlin, Germany, 2018; pp. 91–107. ISBN 978-3-11-053453-5.
36. Henyecz, R.; Keglevich, G. P-C couplings by the Hiaro reaction. In *Organophosphorus Chemistry*; Keglevich, G., Ed.; Walter de Gruyter GmbH: Berlin, Germany, 2018; pp. 158–178. ISBN 978-3-11-053453-5.
37. Enders, D.; Saint-Dizier, A.; Lannou, M.-I.; Lenzen, A. The phospho-Michael addition in organic synthesis. *Eur. J. Org. Chem.* **2006**, 29–49. [[CrossRef](#)]
38. Tajti, Á.; Bálint, E.; Keglevich, G. Synthesis of ethyl octyl α -aminophosphonate derivatives. *Curr. Org. Synth.* **2016**, *13*, 638–675. [[CrossRef](#)]
39. Bálint, E.; Tajti, Á.; Kalocsai, D.; Mátravölgyi, B.; Konstantin, K.; Czugler, M.; Keglevich, G. Synthesis and utilization of optically active α -aminophosphonate derivatives by Kabachnik-Fields reaction. *Tetrahedron* **2017**, *73*, 5659–5667. [[CrossRef](#)]
40. Gerrard, W. The interaction of n-butyl alcohol and the chlorides and oxychloride of phosphorus in the absence and in the presence of pyridine. *J. Chem. Soc.* **1940**, 1466–1469. [[CrossRef](#)]
41. Hardy, E.E.; Anniston, A.; Kosolapoff, G.M. Halogenated Compounds and Process for Making Same. U.S. Patent 2409039, 8 October 1946.
42. Foss, O. Di-O-alkylmonothiophosphates and the corresponding pseudohalogens. *Acta Chem. Scand.* **1947**, *1*, 8–31. [[CrossRef](#)] [[PubMed](#)]
43. Campbell, C.H.; Chadwick, D.H.; Kaufman, S. Continuous process for preparing dialkyl phosphites. *Ind. Eng. Chem. Res.* **1957**, *49*, 1871–1873. [[CrossRef](#)]
44. Mitschke, K.-H. Method for the Combined Production of Diethyl Phosphite and Ethyl Chloride. Application Number WO200424742 A1, 25 March 2004.
45. Kendall, A.J.; Salazar, C.A.; Martino, P.F.; Tyler, D.R. Direct conversion of phosphonates to phosphine oxides: An improved synthetic route to phosphines including the first synthesis of methyl JohnPhos. *Organometallics* **2014**, *33*, 6171–6178. [[CrossRef](#)]
46. Kosolapoff, G.M. Preparation of some mixed dialkyl phosphites. *J. Am. Chem. Soc.* **1951**, *73*, 4989. [[CrossRef](#)]
47. Fields, E. The synthesis of esters of substituted amino phosphonic acids. *J. Am. Chem. Soc.* **1952**, *74*, 1528–1531. [[CrossRef](#)]
48. Oswald, A.A. Synthesis of cyclic phosphorus acid esters by transesterification. *Can. J. Chem.* **1959**, *37*, 1499–1504. [[CrossRef](#)]
49. Kuskov, V.K.; Gradis, G.K. Reaction of diethyl phosphite with sodium alcoholates. *Dokl. Akad. Nauk SSSR* **1953**, *92*, 323–324.
50. Froneman, M.; Modro, T.A. The titanium-mediated transesterification of phosphorus esters. *Tetrahedron Lett.* **1988**, *27*, 3327–3330. [[CrossRef](#)]
51. Aitken, R.A.; Collett, C.J.; Mesher, S.T.E. Convenient preparation of long-chain dialkyl phosphates: Synthesis of dialkyl phosphates. *Synthesis* **2012**, *44*, 2515–2518. [[CrossRef](#)]
52. Bálint, E.; Tajti, Á.; Drahos, L.; Ilia, G.; Keglevich, G. Alcoholysis of dialkyl phosphites under microwave conditions. *Curr. Org. Chem.* **2013**, *17*, 555–562. [[CrossRef](#)]
53. Zarrougui, R.; Raouafi, N.; Lemordant, D. New series of green cyclic ammonium-based room temperature ionic liquids with alkylphosphite-containing anion: Synthesis and physicochemical characterization. *J. Chem. Eng. Data* **2014**, *59*, 1193–1201. [[CrossRef](#)]

54. Salin, A.V.; Il'in, A.V.; Shamsutdinova, F.G.; Fatkhutdinov, A.R.; Galkin, V.I.; Islamov, D.R.; Kataeva, O.N. Phosphine-catalyzed addition of P(O)-H compounds to ethyl phenylpropiolate. *Tetrahedron Lett.* **2015**, *56*, 6282–6286. [[CrossRef](#)]
55. Ernsberger, M.L.; Hill, J.W. Preparation of Organic Phosphorus Compounds, and in Particular, of Dialkyl Phosphites. U.S. Patent 2661364, 1 December 1953.
56. Budnikova, Y.G.; Kargin, Y.M. Electrosynthesis of aliphatic esters of phosphorus acids from white phosphorus in alcohol solutions, involving radical cations of phenothiazine and triarylamine. *Russ. J. Gen. Chem.* **1995**, *65*, 504–507.
57. Abdreimova, R.R.; Akbayeva, D.N.; Polimbetova, G.S.; Caminade, A.-M.; Majoral, J.-P. Chlorine free synthesis of organophosphorus compounds based on the functionalization of white phosphorus (P₄). *Phosphorus Sulfur Silicon Relat. Elem.* **2000**, *156*, 239–254. [[CrossRef](#)]
58. Trofimov, B.; Timokhin, B.; Gusarova, N.; Kazantseva, M. Golubin, Cu-catalyzed oxidative phosphorylation of alkanols with white phosphorus and H₂O₂. *Phosphorus Sulfur Silicon Relat. Elem.* **2002**, *177*, 2385–2390. [[CrossRef](#)]
59. Budnikova, Y.H.; Kafiyatullina, A.G.; Sinyashin, O.G.; Abdreimova, R.R. Electrochemical synthesis of phosphorus esters from white phosphorus in the presence of copper complexes and ethanol. *Russ. Chem. Bull.* **2003**, *52*, 929–938. [[CrossRef](#)]
60. Budnikova, Y.H.; Yakhvarov, D.G.; Sinyashin, O.G. Electrocatalytic eco-efficient functionalization of white phosphorus. *J. Organomet. Chem.* **2005**, *690*, 2416–2425. [[CrossRef](#)]
61. Fisher, H.C.; Prost, L.; Montchamp, J.L. Organophosphorus chemistry without PCl₃: A bridge from hypophosphorous acid to H-phosphonate diesters. *Eur. J. Org. Chem.* **2013**, 7973–7978. [[CrossRef](#)]
62. Kluba, M.; Zwierzak, A. Alkylation of tetra-n-butylammonium alkyl hydrogen phosphites. A new route to mixed dialkyl phosphites. *Synthesis* **1978**, *2*, 134–137. [[CrossRef](#)]
63. Ilia, G.; Kurunczi, L. Synthesis of mixed alkylphosphites and alkylphosphates. *Phosphorus Sulfur Silicon Relat. Elem.* **2003**, *178*, 1513–1519. [[CrossRef](#)]
64. Berchel, M.; Haddad, J.; Le Corre, S.S.; Haelters, J.-P.; Jaffrès, P.-A. Synthesis of lipid-based unsymmetrical O,O-dialkylphosphites. *Tetrahedron Lett.* **2015**, *56*, 2345–2348. [[CrossRef](#)]
65. Guin, J.; Wang, Q.; van Gemmeren, M.; List, B. The catalytic asymmetric Abramov reaction. *Angew. Chem. Int. Ed.* **2014**, *53*, 1–5. [[CrossRef](#)] [[PubMed](#)]
66. Peng, W.; Schreeve, J.M. Rapid and high yield oxidation of phosphine, phosphite and phosphinite compounds to phosphine oxides, phosphates and phosphinates using hypofluorous acid–acetonitrile complex. *J. Fluorine Chem.* **2005**, *126*, 1054–1056. [[CrossRef](#)]
67. Santschi, N.; Togni, A. Electrophilic Trifluoromethylation of S-Hydrogen Phosphorothioates. *J. Org. Chem.* **2011**, *76*, 4189–4193. [[CrossRef](#)] [[PubMed](#)]
68. Kers, A.; Kers, I.; Stawinski, J.; Sobkowski, M.; Kraszewski, A. Studies on Aryl H-Phosphonates; Part 2: A General Method for the Preparation of Alkyl H-Phosphonate Monoesters. *Synthesis* **1995**, 427–430. [[CrossRef](#)]

Sample Availability: Samples of the compounds **2a–d**, **3a–d**, **5a–d** are available from the authors.



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).