This article was downloaded by: [Florida Atlantic University] On: 28 July 2013, At: 18:19 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

Synthesis of 2-(3'-Azido- and 3'-Amino-3'-deoxy-β-Dribofuranosyl)thiazole-4-carboxamide

Cheng Wu Liang ^a , Myong Jung Kim ^a , Lak Shin Jeong ^b & Moon Woo Chun ^{a c}

^a College of Pharmacy, Seoul National University, Seoul, Korea

^b College of Pharmacy, Ewha Womans University, Seoul, Korea

^c College of Pharmacy, Seoul National University, Seoul, 151-742, Korea Published online: 31 Aug 2006.

To cite this article: Cheng Wu Liang , Myong Jung Kim , Lak Shin Jeong & Moon Woo Chun (2003) Synthesis of 2-(3'-Azido- and 3'-Amino-3'-deoxy-β-D-ribofuranosyl)thiazole-4-carboxamide, Nucleosides, Nucleotides and Nucleic Acids, 22:11, 2039-2048, DOI: <u>10.1081/NCN-120026405</u>

To link to this article: <u>http://dx.doi.org/10.1081/NCN-120026405</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

Synthesis of 2-(3'-Azido- and 3'-Amino-3'-deoxy-β-Dribofuranosyl)thiazole-4-carboxamide

Cheng Wu Liang,¹ Myong Jung Kim,¹ Lak Shin Jeong,² and Moon Woo Chun^{1,*}

¹College of Pharmacy, Seoul National University, Seoul, Korea ²College of Pharmacy, Ewha Womans University, Seoul, Korea

ABSTRACT

In view of biological activities of tiazofurin and azido or aminosugar nucleosides, novel azido- and amino-substituted tiazofurin derivatives (1 and 2) were efficiently synthesized starting from 1,2;5,6-di-*O*-isopropylidene-D-glucose.

Key Words: IMPDH inhibitor; Amino sugar; Anticancer nucleosides.

INTRODUCTION

Inosine 5'-monophosphate dehydrogenase (IMPDH) is a critical enzyme in the regulation of cell proliferation and differentiation. This enzyme catalyzes the NAD⁺-dependent oxidation of inosine 5'-monophosphate (IMP) to xanthine 5'-monophosphate (XMP), the rate limiting step in de novo biosynthesis of guanine nucleotides. Therefore, the biochemical effect of IMPDH inhibition in sensitive cell types is decrease in intracellular guanine nucleotide levels,^[1] and the decrease in cellular GTP and deoxy GTP pool levels blocks DNA and RNA synthesis in rapidly

2039

DOI: 10.1081/NCN-120026405 Copyright © 2003 by Marcel Dekker, Inc.

Downloaded by [Florida Atlantic University] at 18:19 28 July 2013

1525-7770 (Print); 1532-2335 (Online) www.dekker.com

^{*}Correspondence: Moon Woo Chun, College of Pharmacy, Seoul National University, Seoul 151-742, Korea; Fax: 82-2-878-9682; E-mail: mjnuh@hanmail.net.

proliferating tumor cells. Because of its critical role in purine biosynthesis, IMPDH is a drug design target for anticancer, antiviral, immunosuppressive and antimicrobial chemotherapy.^[2]

Several compounds have been described as IMPDH inhibitors and among them, tiazofurin, 2- β -D-ribofuranosylthiazole-4-carboxamide, is a C-nucleoside with potent inhibitory activity against IMPDH^[3] currently undergoing clinical trials as an antitumor agent.^[4] This oncolytic C-nucleoside is converted to the nicotinamide adenine dinucleotide (NAD) analogue, thiazole-4-carboxamide adenine dinucleotide (TAD), which binds to the NAD cofactor-binding site of the enzyme and inhibits IMPDH activity.^[5] Recently, several analogues of tiazofurin have been synthesized and are shown in Fig. 1.

It was reported that aminosugar nucleosides possess antiviral and anticancer activities^[6,7] and 3'-azido-3'-deoxythymidine was converted to 3'-amino-3'-deoxy-thymidine in some cells.^[8] One of the most important examples is puromycin,^[9] a derivative of 3'-amino-3'-deoxyadenosine.

Based upon these findings, it was of interest to put an azido or amino group at the C-3' position of tiazofurin in order to compare biological activities with the normal ribonucleoside. An amino group serves as a bioisostere of a hydroxyl group and 2'-hydroxyl substituted 3'-aminonucleoside apparently have a higher population of N-type nucleoside conformations.^[10] Here, we report the synthesis of the azido- and amino-substituted tiazofurin derivatives (1 and 2), starting from 1,2;5,6-di-*O*-isopropylidene-D-glucose.

RESULTS AND DISCUSSION

Our synthetic strategy to the target nucleosides is to synthesize ribofuranosyl cyanide as a key intermediate from 1,2;5,6-di-O-isopropylidene-D-glucose (3) then to cyclocondense with L-cysteine ethyl ester hydrochloride.





Thiazole-4-carboxamide Nucleosides

Starting from commercially available diacetone-D-glucose (3), the hydroxyl group of 3 was converted to a triflate using trifluoromethanesulfonic anhydride. Triflate 4 was used in next step without purification and reacted with sodium azide in DMF to afford the desired azido sugar 5a (44%) and the eliminated enose derivative 5b (32%), respectively. The ratio of 5a and 5b was dependent on the reagent, the reaction temperature and the solvent used.^[11] The reaction mixture could not be heated due to an undesired elimination reaction and when the tosylate of 3 was reacted with sodium azide in presence of 18-crown-6 in DMF, the yield was very poor. The 5,6-O-isopropylidene of 5a was selectively removed using 75% acetic acid to give the diol 6. Oxidative cleavage of 6 with sodium periodate followed by reduction of the resulting aldehyde with sodium borohydride yielded the ribofuranose 7. which was treated with benzoyl chloride to afford the benzoate 8. It was reported that the reaction of 1,2,3,5-tetra-O-acetyl-β-D-ribofuranose with cyanotrimethylsilane in the presence of BF3 OEt2 gave 3,5-di-O-acetyl-1,2-O-(1-endo-cvanoethvlidene)-α-D-ribofuranose.^[12] Therefore, to synthesize 1-O-acetyl-2-O-benzoyl derivative, methylglycoside 9 was prepared from the benzoate 8 by acid-catalyzed methano-1 lysis^[13] of 8. Benzovlation of 9 afforded the dibenzoate 10, which was treated with Ac₂O/AcOH/H₂SO₄ to give 1-O-acetyl-3-azido-2,5-di-O-benzoylribofuranose (11). 3-Azido-2,5-di-O-benzoyl- β -D-ribofuranosyl cyanide (12), a key intermediate, was prepared by reaction of the corresponding acetate 11 with cyanotrimethylsilane and stannic chloride in dichloromethane^[14] (Sch. 1).

Thiazoline **13** was obtained from cyanide **12** by treatment with L-cysteine ethyl ester hydrochloride in presence of triethylamine according to a previous report^[15] and dehydrogenated by treatment with bromotrichloromethane in combination with DBU^[16] at 0°C to give the thiazole **14**. ¹H NMR spectrum of **14** showed the presence of vinyl proton of the thiazole ring. Ester aminolysis and debenzoylation using methanolic ammonia^[17] afforded 3'-azido-substituted tiazofurin derivative **1** in high yield. Catalytic hydrogenation of **1** gave 3'-amino-3'-deoxytiazofurin (**2**) (Sch. 2).

The final nucleosides 1 and 2 were assayed against several viruses such as HIV-1, HSV-1, HSV-2 and HBV. These compounds were found to be inactive against all viruses tested up to $100 \,\mu\text{g}/\text{M}\ell$. Compound 1 and 2 have been evaluated in vitro for their ability to inhibit the growth of Human 143B Osteosarcoma tumor cells, but found to be completely inactive.

In summary, we have successively synthesized the novel 3'-azido- and 3'-amino-3'-deoxytiazofurin (1 and 2) by cyclocondensation of glycosyl cyanide and L-cysteine ethyl ester hydrochloride, followed by dehydrogenation, but the final nucleosides did not exhibit any significant biological activity.

EXPERIMENTAL

General Methods. NMR spectra were recorded in a 300 MHz apparatus using tetramethylsilane (TMS) as an internal standard, and the chemical shifts are reported in ppm (δ). Coupling constants are reported in hertz (Hz). Infrared spectra were recorded in a Perkin-Elmer 1710 FTIR spectrophotometer. Mass spectra recorded by FAB (Fast atom bombardment) on a VG Tro-2, GC-MS. TLC were carried

2041



Scheme 1. Reagents and conditions: (a) Tf₂O, pyridine, CH₂Cl₂, 0°C, 1 h; (b) NaN₃, DMF, RT, 16 h, 5a(44%), 5b(32%); (c) 75% AcOH, 55°C, 1.5 h, 90%; (d) 1: NaIO₄, EtOH, 0°C, 0.5 h; 2: NaBH₄, 0°C, 2 h, 84%; (e) BzCl, pyridine, RT, 3 h, 95%; (f) 1% HCl in MeOH, RT, 2 h, 80%; (g) BzCl, pyridine, RT, 3 h, 97%; (h) AcOH, Ac₂O, c-H₂SO₄, RT, 30 min, 98%; (i) TMSCN, SnCl₄, CH₂Cl₂, 70°C, 3 h, 63%.

out on Merck silica gel 60 F_{254} precoated plates, and silica gel column chromatography was performed on silica gel 60, $230 \sim 400$ mesh, Merck. All anhydrous solvents were distilled over CaH₂ or Na/benzophenone prior to use.

3-Azido-3-deoxy-1,2;5,6-di-*O*-isopropylidene- α -D-allofuranose (5a) and 3-Deoxy-1,2;5,6-di-*O*-isopropylidene- α -D-erythro-hex-3-enofuranose (5b). Trifluoro-methanesulfonic anhydride (2.9 mL, 17.24 mmol) was added dropwise to a solution of 1,2;5,6-di-*O*-isopropylidene- α -D-glucose (3g, 11.53 mmol) and pyridine (2.8 mL, 34.62 mmol) in dichloromethane (30 mL) at 0°C. The reaction mixture was stirred for 1 h at 0°C and extracted with dichloromethane and water. The organic layer was washed with brine, dried (MgSO₄), filtered and concentrated to give **4**.

The triflate **4** was dissolved in anhydrous DMF (20 mL) and treated with sodium azide (2.25 g, 34.61 mmol). The reaction mixture was stirred for 16 h at room temperature. Ethyl acetate and water were added and aqueous phase was extracted with

Downloaded by [Florida Atlantic University] at 18:19 28 July 2013



Scheme 2. Reagents and conditions: (a) L-cysteine ethyl ester hydrochloride, Et_3N , MeOH, RT, 2h, 75%; (b) DBU, BrCCl₃, CH₂Cl₂, 0°C, 16h, 88%; (c) NH₃/MeOH, RT, 24h, 92%; (d) H₂.Pd/C, EtOH, RT, 14h, 90%.

ethyl acetate (×3). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 4/1) to give **5a** (1.45 g, 5.08 mmol, 44%) as an oil and **5b** (0.89 g, 3.67 mmol, 32%) as a solid.

5a: $R_f 0.33$ (hexane/EtOAc = 4/1); IR (KBr): 2987, 2109, 1377, 1259, 1214, 1164, 1065, 1024 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 5.76 (d, 1H, *J* = 3.7 Hz, H-1), 4.70 (t, 1H, *J* = 4.1 Hz, H-2), 3.95–4.21 (m, 4H, H-4, H-5, H-6), 3.48 (dd, 1H, *J* = 4.9, 9.0 Hz, H-3), 1.56, 1.46, 1.36, 1.34 (4 s, 12H, 4 × CH₃); FAB-MS *m/z*: 286 (M + H)⁺.

5b: $R_f 0.52$ (hexane/EtOAc = 4/1); IR(KBr): 2988, 2938, 1667, 1376, 1321, 1214, 1157, 1073 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 6.05 (d, 1H, J = 5.1 Hz, H-1), 5.26 (m, 1H, H-3), 5.21 (m, 1H, H-2), 4.54 (dd, 1H, J = 5.9, 6.6 Hz, H-5), 4.10 (dd, 1H, J = 6.8, 8.3 Hz, H-6a), 3.92 (dd, 1H, J = 5.9, 8.5 Hz, H-6b), 1.44, 1.42, 1.37 (3 s, 12H, 4 × CH₃); FAB-MS m/z: 243 (M + H)⁺.

3-Azido-3-deoxy-1,2-*O***-isopropylidene-** α **-D-allofuranose (6). 5a** (2.15 g, 7.54 mmol) was dissolved in 75% acetic acid (30 mL). The reaction mixture was stirred for 2 h at 55°C, evaporated to dryness and coevaporated with toluene. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 1/2) to give 6 (1.66 g, 6.77 mmol, 90%) as an oil. ¹H NMR (300 MHz, CDCl₃): δ 5.81 (d, 1H, J = 3.7 Hz, H-1), 4.75 (dd, 1H, J = 3.7, 4.8 Hz, H-2), 4.07 (dd, 1H, J = 4.2, 9.3 Hz, H-4), 3.75–4.01 (m, 3H, H-5, H-6), 3.59 (dd, 1H, J = 4.9, 9.4 Hz, H-3), 2.94 (brs, 1H, OH),

2.51 (brs, 1H, OH), 1.59, 1.38 (2 s, 6H, $2 \times CH_3$); FAB-MS m/z: 268 (M + Na)⁺, 246 (M + H)⁺.

3-Azido-3-deoxy-1,2-*O***-isopropylidene-***a***-D-ribofuranose** (7). A solution of sodium periodate (1.57 g, 7.34 mmol) in water (15 mL) was added dropwise to a stirred solution of **6** (1.66 g, 6.77 mmol) in EtOH (30 mL) at 0°C. After 30 min, sodium borohydride (597 mg, 15.8 mmol) was added. The reaction mixture was stirred for 2 h at 0°C, filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 1/1) to give 7 (1.217 g, 5.65 mmol, 84%) as an oil. IR (KBr): 3437, 2988, 2108, 1258, 1110, 1020 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 5.79 (d, 1H, J = 3.4 Hz, H-1), 4.72 (t, 1H, J = 4.0 Hz, H-2), 4.09 (m, 1H, H-4), 3.96 (dd, 1H, J = 2.4, 12.5 Hz, H-5a), 3.66 (dd, 1H, J = 2.9, 12.6 Hz, H-5b), 3.56 (dd, 1H, J = 4.6, 9.5 Hz, H-3), 1.56, 1.35 (2 s, 6H, 2 × CH₃); FAB-MS *m/z*: 238 (M + Na)⁺, 216 (M + H)⁺.

3-Azido-5-*O***-benzoyl-3-deoxy-1,2-***O***-isopropylidene-α-D-ribofuranose (8).** To a stirred solution of 7 (1.217 g, 5.65 mmol) in pyridine (20 mL) was added benzoyl chloride (0.98 mL, 8.44 mmol) at 0°C. The mixture was stirred for 3 h at room temperature and evaporated to dryness. The residue was dissolved in EtOAc and extracted with water. The organic layer was washed with saturated NaHCO₃ and brine, dried (MgSO4), filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 4/1) to give **8** (1.72 g, 5.39 mmol, 95%) as an oil. IR (KBr): 2108, 1724, 1275, 1213, 1111, 1021, 710 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.60–8.03 (m, 5H, Ph), 5.85 (d, 1H, J = 3,7 Hz, H-1), 4.78 (dd, 1H, J = 3.7, 4.6 Hz, H-2), 4.66 (dd, 1H, J = 3.1, 12.3 Hz, H-5a), 4.46 (dd, 1H, J = 4.6, 12.2 Hz, H-5b), 4.37 (m, 1H, H-4), 3.43 (dd, 1H, J = 4.7, 9.7 Hz, H-3), 1.52, 1.38 (2 s, 6H, 2 × CH₃); FAB-MS *m/z*: 342 (M + Na)⁺.

3-Azido-5-O-benzoyl-1-O-methyl-3-deoxy-D-ribofuranose (9). To a stirred solution of **8** (1.72 g, 5.39 mmol) in MeOH (15 mL) was added acetyl chloride (0.2 mL) at room temperature. After stirring for 3 h at room temperature, the reaction mixture was neutralized with pyridine and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 2/1) to give **9** (1.257 g, 4.3 mmol, 80%) as an oil. IR (KBr): 3462, 2938, 2109, 1722, 1274, 1120, 1070, 713 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.43–8.10 (m, 5H, Ph), 4.95 (d, 1H, J = 4.4 Hz, α isomer H-1), 4.89 (s, 1H, β isomer H-1), 4.33–4.61 (m, 6H, α and β isomer H-2, H-4, H-5a), 4.13–4.28 (m, 2H, H-5b, β isomer H-3), 3.95 (dd, 1H, J = 4.0, 7.6 Hz, α isomer H-3), 3.45 (s, 3H, α isomer OCH₃); 5AB-MS *m*/*z*: 294 (M + H)⁺, 262 (M-OCH₃)⁺.

3-Azido-2,5-di-O-benzoyl-1-O-methyl-3-deoxy-D-ribofuranose (10). To a stirred solution of **9** (1.257 g, 4.3 mmol) in pyridine (10 mL) was added benzoyl chloride (0.75 mL, 6.46 mmol) dropwise at room temperature and the reaction mixture was stirred for 3 h at room temperature. The reaction mixture was evaporated and the residue was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried (MgSO₄), filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 4/1) to

Thiazole-4-carboxamide Nucleosides

give **10** (1.645 g, 4.14 mmol, 97%) as an oil. IR (KBr): 2111, 1725, 1268, 1111, 1070, 710 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.45–8.18 (m, 10H, 2 × Ph), 5.51 (d, 1H, J = 4.6 Hz, H-2), 5.05 (s, 1H, H-1), 4.63 (m, 1H, H-4), 4.45–4.51 (m, 2H, H-5), 4.32 (dd, 1H, J = 4.6, 7.8 Hz, H-3), 3.38 (s, 3H, OCH₃); FAB-MS *m*/*z*: 398 (M + H)⁺, 366 (M-OCH₃)⁺.

1-O-Acetyl-3-azido-2,5-di-O-benzoyl-3-deoxy-D-ribofuranose (11). To a stirred solution of **10** (1.645 g, 4.14 mmol) in acetic acid (16 mL) were added acetic anhydride (4 mL) and c-H₂SO₄ (1.12 mL) at room temperature. The reaction mixture was stirred for 30 min at room temperature and poured into saturated NaHCO₃ solution. The mixture was extracted with dichloromethane (×3). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 3/1) to give **11** (1.73 g, 4.07 mmol, 98%) as an oil. IR (KBr): 2952, 2113, 1726, 1268, 1111, 1022, 7111 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.35–8.02 (m, 10H, 2 × Ph), 6.52 (d, 1H, J = 4.4 Hz, α isomer H-1), 6.22 (s, 1H, β isomer H-1), 5.52 (d, 1H, J = 4.6 Hz, β isomer H-2), 5.39 (dd, 1H, J = 4.4, 7.5 Hz, α isomer H-2), 4.37–4.78 (m, 6H, α and β isomer H-4, H-5), 4.26 (dd, 1H, J = 4.6, 7.9 Hz, β isomer H-3), 4.20 (dd, 1H, J = 3.6, 7.5 Hz, α isomer H-3), 2.05 (s, 3H, α isomer OAc); FAB-MS *m/z*: 448 (M + Na)⁺, 366 (M-OAc)⁺.

3-Azido-2,5-di-*O***-benzoyl-1-cyano-3-deoxy-β-D-ribofuranose (12).** To a stirred solution of **11** (1.73 g, 4.07 mmol) and TMSCN (2.17 mL, 16.27 mmol) in anhydrous dichloromethane (20 mL) was added 1M solution SnCl₄ in dichloromethane (0.8 mL) dropwise at room temperature. The reaction mixture was refluxed for 3 h and poured into saturated NaHCO₃ solution. The mixture was extracted with dichloromethane (×3). The combined organic layers were washed with brine, dried (MgSO₄), filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 3/1) to give **12** (1 g, 2.55 mmol, 63%) as an oil. IR (KBr): 2116, 1725, 1601, 1452, 1267, 1093, 711 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.46–8.15 (m, 10H, 2 × Ph), 5.85 (dd, 1H, J = 2.9, 4.6 Hz, H-2), 4.88 (d, 1H, J = 3.0 Hz, H-1), 4.65 (dd, 1H, J = 3.2, 12.7 Hz, H-5a), 4.54 (dd, 1H, J = 3.4, 12.4 Hz, H-5b), 4.39–4.49 (m, 2H, H-3, H-4); FAB-MS *m/z*: 415 (M + Na)⁺, 393 (M + H)⁺.

Ethyl 2-(3-Azido-2,5-di-*O*-benzoyl-3-deoxy-β-D-ribofuranosyl)-thiazoline-4-carboxylate (13). To a stirred solution of 3-azido-2,5-di-*O*-benzoyl-1-cyano-β-D-ribofuranose 12 (763 mg, 1.94 mmol) in dry MeOH (25 mL) was added L-cysteine ethyl ester hydrochloride (542 mg, 2.92 mmol) followed by TEA (0.4 mL, 2.88 mmol) at room temperature. The reaction mixture was stirred for 2 h and evaporated to dryness. The residue was dissolved in CH₂Cl₂ and washed with water, saturated NaHCO₃ solution and brine. The organic layer was dried (MgSO₄), filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 3/1) to give 13 (758 mg, 1.45 mmol, 75%) as an oil IR (KBr): 2984, 2098, 1724, 1272, 1122, 1064, 1025, 713 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.44–8.13 (m, 10H, 2 × Ph), 5.84 (m, 1H, H-2'), 5.03 (m, 2H, H-1', CHCO₂Et), 4.69 (m, 1H, H-4'), 4.51 (m, 1H, H-5'a), 4.13 (m, 4H, H-3', H-5'b, SCH₂), 3.37 (m, 2H, OCH₂CH₃), 1.24 (m, 3H, OCH₂CH₃); FAB-MS m/z: 419 (M-OBz)⁺; Anal. calcd for C₂₅H₂₄N₄O₇S: C, 57.24; H, 4.61; N, 10.68; S, 6.11. Found: C, 57.02; H, 4.51; N, 10.58; S, 6.21.

Ethyl 2-(3-Azido-2,5-di-*O*-benzoyl-3-deoxy-β-D-ribofuranosyl)-thiazole-4-carboxylate (14). To a stirred solution of 13 (758 mg, 1.45 mmol) in anhydrous dichloromethane was added 1,8-diazabicyclo[5.4.0]-undec-7-ene (0.43 mL, 2.88 mmol). The solution was cooled to 0°C and BrCCl₃ (0.17 mL, 1.73 mmol) was added dropwise. The reaction mixture was stirred for 16h at 0°C and evaporated to dryness. The residue was dissolved in EtOAc and washed with saturated aqueous NH₄Cl (×3). The organic layer was dried (MgSO₄), filtered and evaporated to dryness. The residue was purified by silica gel flash column chromatography (hexane/EtOAc = 3/1) to give 14 (666 mg, 1.27 mmol, 88%) as an oil. IR (KBr): 2983, 2113, 1725, 1267, 1096, 711 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 7.42–8.24 (m, 11H, SCH, 2 × Ph), 5.96 (dd, 1H, J = 2.9, 4.9 Hz, H-2'), 5.58 (d, 1H, J = 2.9 Hz, H-1'), 4.80 (dd, 1H, J = 3.0, 12.3 Hz, H-5'a), 4.36 (m, 5H, H-3', H-4', H-5'b, OCH₂CH₃), 1.37 (t, 3H, J = 7.06 Hz, OCH₂CH₃); FAB-MS m/z: 545 (M + Na)⁺, 523 (M + H)⁺; Anal. calcd for C₂₅H₂₂N₄O₇S: C, 57.46; H, 4.24; N, 10.72; S, 6.14. Found: C, 57.66; H, 4.20; N, 10.62; S, 6.24.

2-(3-Azido-2,5-di-*O***-benzoyl-3-deoxy-β-D-ribofuranosyl)thiazole-4-carboxamide** (1). A mixture of **14** (666 mg, 1.27 mmol) in saturated NH₃ in MeOH (20 mL) was stirred for 24 h at room temperature and evaporated to dryness. The residue was purified by silica gel flash column chromatography (CH₂Cl₂/MeOH = 7/1) to give 1 (334 mg, 1.17 mmol, 92%) as a white foam. IR (KBr): 3427, 2925, 2109, 1660, 1260, 1047 cm⁻¹; ¹H NMR (300 MHz, CD₃OD): δ 8.19 (s, 1H, SCH), 5.00 (d, 1H, J = 5.4 Hz, H-1'), 4.45 (t, 1H, J = 5.5 Hz, H-2'), 4.02 (d"d", 1H, J = 3.9, 4.1, 5.4 Hz, H-4'), 3.92 (dd, 1H, J = 5.4, 5.6 Hz, H-3'), 3.71 (dd, 1H, J = 3.9, 12.2 Hz, H-5'a), 3.64 (dd, 1H, J = 4.4, 12.2 Hz, H-5'b); additional signals in ¹H NMR (300 MHz, DMSO-d₆): δ 7.71 (brs, 1H, NH₂), 7.57 (brs, 1H, NH₂), 6.15 (d, 1H, J = 5.6 Hz, OH), 5.01 (t, 1H, J = 5.3 Hz, OH); FAB-MS m/z: 308 (M + Na)⁺, 286 (M + H)⁺; Anal. calcd for C₉H₁₁N₅O₄S: C, 37.89; H, 3.89; N, 24.55; S, 11.24. Found: C, 37.54; H, 4.08; N, 24.25; S, 11.42.

2-(3-Amino-2,5-di-*O***-benzoyl-3-deoxy-β-D-ribofuranosyl)thiazole-4-carboxamide** (2). To a solution of **1** (280 mg, 0.98 mmol) in EtOH (20 mL) was added 5% Pd/C (100 mg) and the reaction mixture was degassed and stirred at one atmosphere of hydrogen for 14 h at room temperature. The catalyst was removed by filtration and the filtrate was evaporated to dryness. The residue was recrystalized from MeOH/EtOAc to give **2** (230 mg, 0.89 mmol, 90%) as a white solid. IR (KBr): 3433, 2918, 1658, 1381, 1043 cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ 8.18 (s, 1H, SCH), 7.64 (br s, 1H, NH₂), 7.55 (br s, 1H, NH₂), 4.99 (d, 1H, J=1.95 Hz, H-1′), 4.01 (dd, 1H, J=1.95, 9.72 Hz, H-2′), 3.63 (m, 2H, H-4′, H-3′), 3.55 (m, 1H, H-5′a), 2.96 (dd, 1H, J=4.9, 8.5 Hz, H-5′b), 1.41 (brs, 2H, NH₂); FAB-MS m/z: 260 [M + H]⁺; Anal. calcd for C₉H₁₃N₃O₄S: C, 41.69; H, 5.05; N, 16.21; S, 12.37. Found: C, 41.52; H, 5.15; N, 16.01; S, 12.29. Downloaded by [Florida Atlantic University] at 18:19 28 July 2013

REFERENCES

- (a) Eugui, E.M.; Almquist, S.T.; Mullwe, C.D.; Allison, A.C. Lymphocyteselective cytostatic and immunosuppressive effects of mycophenolic acid in vitro: Role of deoxyguanosine nucleotide depletion. Scand. J. Immunol. 1991, 33, 161; (b) Franklin, T.J.; Morris, W.P. Pharmacodynamics of the inhibition of GTP synthesis in vivo by mycophenolic acid. Adv. Enzyme Regul. 1994, 34, 107.
- 2. Franchetti, P.; Cappellacci, L.; Grifantini, M. IMP dehydrogenase as a target of antitumor and antiviral chemotherapy. Farmaco **1996**, *51*, 457; and references therein.
- Jayaram, H.N.; Dion, R.L.; Glazer, R.J.; Johns, D.G.; Robins, R.K.; Srivastava, P.C.; Cooney, D.A. Initial studies on the mechanism of action of a new oncolytic thiazole nucleoside, 2-β-D-ribofuranosylthiazole-4-carboxamide (NSC 286193). Biochem. Pharmacol. **1982**, *31*, 2371.
- (a) Tricot, G.J.; Jayaram, H.N.; Lapis, E.; Natsumeda, Y.; Nichols, C.R.; Kneebone, P.; Heerema, N.; Weber, G.; Hoffman, R. Biochemically directed therapy of leukemia with tiazofurin, a selective blocker of inosine 5'-phosphate dehydrogenase activity. Cancer Res. 1989, 49, 3696; (b) Wright, D.G.; Boosalis, M.S.; Waraska, K.; Osrly, L.J.; Weeintraub, L.R.; Vosburgh, E. Tiazofurin effects on IMP-dehydrogenase activity and expression in the leukemia cells of patients with CML blast crisis. Anticancer Res. 1996, 16, 3349.
- Lui, M.S.; Faderam, M.A.; Liepnieks, J.J.; Natsumeda, Y.; Alah, E.; Jayaram, H.N.; Weber, G. Modulation of IMP dehydrogenase activity and guanylate metabolism by tiazofurin (2-β-D-ribofuranosylthiazole-4-carboxamide). J. Biol. Chem. 1984, 259, 5078.
- 6. Suhadolnik, R.J. Nucleoside Antibiotics; Wiley-Interscience: New York, 1970.
- Suhadolnik, R.J. Nucleosides as Biological Probes; Wiley-Interscience: New York, 1979.
- 8. Balzarini, J. Metabolism and mechanism of antiretroviral action of purine and pyrimidine derivatives. Pharm. World Sci. **1994**, *16*, 113.
- 9. Baker, B.R.; Joseph, J.P.; Williams, J.H. Synthetic studies. VII. Partial synthesis of amino acid analogs. J. Am. Chem. Soc. 1955, 77, 1.
- 10. Herdewijn, P.; Balzarin, J.; Masanori, B.; Pauwels, R.; Van Aerschot, A.; Janssen, G.; De Clercq, E. Synthesis and anti-HIV activity of different sugarmodified pyrimidine and purine nucleosides. J. Med. Chem. **1988**, *31*, 2040 and references therein.
- Neufellner, E.; Kapeller, H.; Griengl, H. A novel access to derivatives of 3azido-3-deoxy-4a-carba-α-DL-ribofuranose potential intermediates for the synthesis of carbachryscandin and carbapuromycin. Tetrahedron 1988, 54, 11,043.
- Utimoto, K.; Wakabayashi, Y.; Horiie, T.; Inoue, M.; Shishiyama, Y.; Obayashi, M.; Nozaki, H. Cyanotrimethylsilane as a versatile reagent for introducing cyanide functionality. Tetrahedron 1983, 39, 967.
- Bolon, P.J.; Sells, T.B.; Nuesca, Z.M.; Purdy, D.F.; Nair, V. Novel isomeric dideoxynucleosides as potential antiviral agents. Tetrahedron 1994, 50, 7747.
- 14. Utimoto, K.; Horiie, T. A facile preparation of *O*-acelated β -D-ribofuranosyl cyanide, an important intermediate of c-nucleoside synthesis. Tetrahedron Lett. **1982**, *23*, 237.

- Ramasamy, K.S.; Bandaru, R.; Averett, D. A new synthetic methodology for tiazofurin. J. Org. Chem. 2000, 65, 5849.
- 16. Williams, D.R.; Lowder, P.D.; Gu, Y.G.; Brooks, D.A. Studies of mild dehydrogenations in heterocyclic systems. Tetrahedron Lett. **1997**, *38*, 331.
- 17. Srivastava, P.C.; Pickering, M.V.; Allen, L.B.; Streeter, D.G.; Campbell, M.T.; Witkowski, J.T.; Sidwell, R.W.; Robins, R.K. Synthesis and antiviral activity of certain thiazole C-nucleosides. J. Med. Chem. **1977**, *20*, 236.

Received February 21, 2003 Accepted August 11, 2003

2048