

PII: S0040-4039(96)00845-3

Catalysis and Regioselectivity in the Michael Addition of Azoles. Kinetic vs. Thermodynamic Control

András Horváth

Alkaloida Chemical Company Ltd., H-4440 Tiszavasvári, Pf. 1, Hungary

Abstract: Bicyclic guanidine bases, TBD and MTBD were found to be highly efficient catalysts in the Michael addition of azoles with α , β -unsaturated nitriles and esters. The factors influencing regioselectivity have been elucidated, and some new azole-Michael adducts were synthesized. These were shown to be useful starting compounds for the regioselective *N*-alkylation of the corresponding azoles. Copyright © 1996 Elsevier Science Ltd

Michael adducts are useful starting compounds for obtaining the thermodynamically less stable regioisomers of various N-substituted azoles, compounds with diverse pharmacological uses.¹ Since overall efficiency of that procedure is mainly dependent on the outcome of the N-protective step (*i.e.* the Michael addition) we found it worthy to study the factors influencing the yield and regioselectivity of the Michael addition of azoles.

Though azoles with higher nucleophilicity, such as imidazole, *C*-alkyl imidazoles and 1,2,4-triazole react fairly well with Michael acceptors without any catalyst, reaction of azoles with lower nucleophilicity proceeds very sluggishly, and catalysts such as pyridine,^{2,3} triethylamine,^{4,5} tetraalkylammonium hydroxides,^{2,4,6-8} KOH^{9,10} or MeONa^{11,12} were employed. Our investigations indicated that while triethylamine and pyridine showed low efficiency in most of the cases, other bases lead to considerable amounts of byproducts (e.g. by alcohol addition or polymerization) due to their nucleophilic character.

In the present study, we found the commercially available strong, bicyclic guanidine bases, TBD and MTBD to be excellent catalysts for the reaction of azoles with α , β -unsaturated nitriles and esters (Table 1):

Azole	Acrylonitrile	Solvent	Catalyst	t	τ	Isolated	Lit.
	(eq.)			(°C)	(hours)	yield (%)	
Benzimidazole	2.5	EtOH	PhNMe ₃ OH	40		83	6
	1	acrylonitrile	BnNMe ₃ OH	25	24		13
		acrylonitrile	Et ₃ N	76	90	72	14
	1.2	MeCN	MTBD ^a	25	1	91	14
4-Phenyl-imidazole	3	EtOH	КОН	78	2	54	9
	1.2	MeCN	TBD ^b	25	0.15	95	14
4-Nitro-imidazole	1.2	DMSO	pyridine	140	10	90	3
	1.2	MeCN	MTBD	80	3	92	14

Table 1. Comparative data on the addition of some azoles to acrylonitrile

^a Aldrich no. 35,950-5; ^b Aldrich no. 34,557-1

In the presence of 1-5 mol % TBD or MTBD in acetonitrile solution, the Michael-additions proceeded faster even at lower temperatures, and no side reactions could be detected. The azole-Michael adducts could be isolated in higher yields and purity when compared to literature methods.¹⁵

Systematic studies on regioselectivity of Michael additions of imidazoles and benzimidazoles are scarce, except for reports on 2,4-dialkylimidazoles,¹⁶ 5-nitrobenzimidazoles¹⁷ and 4-nitroimidazoles.³ The present study extends to the Michael additions of the most common α , β -unsaturated nitriles and esters with imidazoles and benzimidazoles bearing various 4-substituents (Table 2):



Scheme 1

<u>**Table 2.**</u> Ratio of 3:4(1,4 to 1,7 adducts) and 6:7(1,4 to 1,5 adducts) in the TBD-catalyzed Michael addition of azoles at room temperature in MeCN as solvent, determined by ¹H NMR of crude products

Azole	Substituent		2a	2ь	2c	2d
	R	Effect	Acrylonitrile	Et-acrylate	Crotononitrile	Me-crotonate
1a ^a	NO ₂	-E, -I	$\sim 100:0^{b}$	> 99 : 1	~ 100 : 0	~ 100 : 0
1b	Me	+I	86:14	81:19	> 99 : 1 ^b	90:10
5a ^{c,d}	NO ₂	-E, -I	$\sim 100:0^b$	> 99 : 1	~ 100 : 0	> 99 : 1
5b	CH=CHCO ₂ Et	-E, -I	$\sim 100:0^{b}$	> 99 : 1	~ 100 : 0	> 99 : 1
5c	Ph	+E, -I	$> 99:1^{b}$	88:12	~ 100 : 0	97:3
5d ^{18,d}	CH=CHCMe ₂ OH	+E, -I	80:20	75:25	93:7	89:11
5e ^d	(CH ₂) ₂ CO ₂ Et	+I	72:28	65:35	88:12	82:18
5f ^e	Me	+I	63:37	63:37	81:19 ^b	81:19

^{*a*} Reactions in 2 as solvent in the presence of 1 eq. of DMAP at 80 °C; ^{*b*} Isolated yields: **3aa**: 88 %, **3bc**: 96 %, **6aa**: 92 %, **6ba**: 92 %, **6ca**: 95 %, **6fc oxalate**: 65 %; ^{*c*} Reaction at 80 °C; ^{*d*} Catalyst: MTBD; ^{*e*} Without catalyst at 80 °C

We found that TBD-catalyzed addition of acrylonitrile and crotononitrile was more selective in most cases than addition of the corresponding α , β -unsaturated esters. To elucidate the origin of this observation, first, we proved (by ¹H NMR in CD₃CN solution) that TBD reacts rapidly with an equivalent of acrylonitrile or ethyl acrylate to form the corresponding Michael addition products (TBD-MAs): 1,3,4,6,7,8-hexahydro-2*H*-pyrimido[1,2-a]pyrimidine-1-propanoic acid derivatives (nitrile and ethyl ester, respectively).

The *in situ* formed TBD-MAs were then reacted with an equivalent of azole (4-phenyl-imidazole or 4methyl-benzimidazole). While reactions of cyanoethyl-TBD at 30 °C proceeded slowly towards equilibrium mixtures containing the azole, azole-MA, TBD and TBD-MA, no reaction was observed with (ethoxycarbonyl)ethyl-TBD in the 30-70 °C temperature range. Moreover, reaction of the cyanoethyl-azoles with an equivalent of TBD proceeded towards the same equilibrium mixtures, while the azole-ethyl acrylate adducts did not react with TBD, demonstrating that TBD-catalyzed reactions of 4-phenyl-imidazole and 4-methylbenzimidazole with acrylonitrile are reversible, while additions of ethyl acrylate are irreversible. Thus, the regioselectivity data reflect *thermodynamic control* for the α,β -unsaturated nitriles and *kinetic control* in the case of the α,β -unsaturated esters. This is true for all azoles studied (except for **5f**, with all four reactions thermodinamically controlled under the conditions stated), as it was checked by reacting the isolated adducts with TBD.

Compounds **3bc**, **6aa** and **6ca** were already shown to be useful substrates for obtaining *N*-substituted derivatives of the parent azoles **1b**, **5a** and **5c**, respectively. Derivatives of **5e** are accessible comfortably from **6ea** obtained by hydrogenation of **6ba** over Pd/C (96 % yield). The most reluctant case,¹ the regioselective alkylation of 4-methyl-imidazole could be solved using conventional separation techniques: the pure 1,4-adduct of 4-methyl-imidazole and crotononitrile (α ,4-dimethyl-1*H*-imidazole-1-propanenitrile, **6fc**) was obtained by fractional crystallization of the adduct mixture as the oxalate salt. Compound **6fc** provides an easy access to both 1-substituted-5- (**8a**: R= benzyl and **8b**: R= pivaloyloxymethyl) and 1-substituted-4-methyl-imidazoles (*e. g.* 1-phenacyl-4-methyl-imidazole **9**):¹⁹





The efficient catalysis of the Michael additions of azoles by the guanidine bases was used in the highyield preparation of the previously unknown cyanoethyl derivatives of 4-nitrobenzimidazole **3aa**, urocanic acid ethyl ester **6ba**, 3-phenyl-pyrazole **10** and 3-phenyl-1,2,4-triazole **11** (Scheme 3) in thermodynamically



Scheme 3

controlled reactions with acrylonitrile. These adducts are useful starting compounds for the preparation of the thermodynamically less stable *N*-substituted derivatives of the parent azoles: quaternization (with allyl bromide or dimethyl sulfate, respectively) followed by deprotection afforded the known 1-methyl-7-nitrobenzimidazole²⁰ (yield: 75 %), *E*-1-(2-propenyl)-1*H*-imidazole-5-propenoic acid ethyl ester (or N^{π} -allyl urocanic acid ethyl ester²¹) (89 %), 1-methyl-5-phenyl-1*H*-pyrazole²² (60 %) and 4-methyl-3-phenyl-4*H*-1,2,4-triazole²³ (64 %), respectively.

<u>Acknowledgments</u>: The author thanks Dr. Zoltán Szabó and Dr. József Jekő for performing NMR and MS measurements, Dr. Zoltán Györgydeák and Dr. Tibor Tímár for discussions on the manuscript and Mrs. Piroska Fekete and Mrs. Magdolna Dancs for technical assistance.

References and Notes:

- 1. Horváth, A. Synthesis 1995, 1183-1189; see also references cited therein.
- 2. Wiley, R. H.; Smith, N. R.; Johnson, D. M.; Moffat, J. J. Am. Chem. Soc. 1954, 76, 4933-4935.
- 3. Rao, A. K. S. B.; Rao, C. G.; Singh, B. B. J. Org. Chem. 1990, 55, 3702-3704.
- 4. Bogatkov, S. V.; Kormanskaya, B. M.; Mochalin, V.; Cherkasova, E. M. *Khim. Geterotsikl. Soedin.* 1971, 7, 662-664; *Chem. Abstr.* 1972, 76, 59525x.
- 5. Dziklinska, H.; Dziergowski, S.; Jezewski, A.; Plenkiewicz, J. Bull. Soc. Chim. Belg. 1989, 98, 277-283.
- 6. Efros, L. S.; Porai-Koshits, P. A. Zhur. Obschchei Khim. 1953, 23, 697-705; Chem. Abstr. 1954, 48, 7603.
- 7. Zelnik, R.; Pesson, M. Bull. Soc. Chim. Fr. 1959, 1667-1669.
- 8. Zajaczkowska, J.; Eckstein, M. Diss. Pharm. Pharmacol. 1968, 20, 497-505.
- Iradyan, M. A.; Torosyan, A. G.; Mirzoyan, R. G.; Badalyants, I. P.; Isaakyan, Z. S.; Manucharyan, D. S.; Dayan, M. K.; Sakanyan, G. S.; Dzhagatspanyan, I. A.; Akonyan, N. E.; Ter-Zaharyan, Y. Z.; Aroyan, A. A. Khim.-Farm. Zh. 1977, 11, 42-48; Chem. Abstr. 1978, 88, 22759y.
- 10. Bany, T.; Dobosz, M. Ann. Univ. Mariae Curie-Sklodowska, Sect. AA: Chem. 1980, 31-32, 271-276.
- 11. Rybar, A.; Stibranyi, L. Collect. Czech. Chem. Commun. 1973, 38, 1571-1577.
- 12. Dallacker, F.; Minn, K. Chem.-Ztg. 1986, 110, 101-108.
- 13. Diamond, J.; Wohl, R. A. Eur. Pat. Appl. 34,116 (1981); Chem. Abstr. 1981, 95, 203961y.
- 14. Horváth, A. Syntheses of Azolyl-Alkyl-Thioureas and Azolyl-Alkyl-Pyrimidines, Lajos Kossuth University, Debrecen, Hungary, 1995.
- 3aa: mp 195-196 °C (acetone). ¹H NMR (200 MHz) (DMSO-d₆): 3.19 (t, 2H), 4.72 (t, 2H), 7.52 (t, 1H), 8.10 (dd, 1H), 8.25 (dd, 1H), 8.63 (s, 1H). MS (El⁺, 70 eV): m/z (%): 216 (M⁺, 100), 186 (68), 146 (33), 118 (94).
 6ba: mp 120-121.5 °C (EtOH). ¹H NMR (CDCl₃): 1.31 (t, 3H), 2.84 (t, 2H), 4.16-4.33 (m, 4H), 6,57 (d, 1H, *J*= 15 Hz), 7.22 (m, 1H), 7.55 (d, 1H, *J*= 15 Hz), 7.59 (m, 1H). MS: 219 (M⁺, 27), 174 (100), 147 (46), 105 (25). 6fc oxalate: mp 104-105.5 °C (EtOH). ¹H NMR (DMSO-d₆): 1.51 (d, 3H, *J*= 6.8 Hz), 2.20 (d, 3H, *J*= 0.9 Hz), 3.17 (d, 2H, *J*= 6.6 Hz), 4.77 (sext, 1H, *J*= 6.7 Hz), 7.38 (m, 1H), 8.51 (d, 1H, *J*= 1.5 Hz). MS: 149 (base M⁺, 48), 109 (100), 81 (43). 10: yield: 95 %, mp 51-53 °C (petr. ether : tolucne 2:1). ¹H NMR (CDCl₃): 3.00 (t, 2H), 4.42 (t, 2H), 6.57 (d, 1H, *J*= 3.0 Hz), 7.30-7.47 (m, 3H), 7.52 (d, 1H, *J*= 3.0 Hz), 7.78 (m, 2H). MS: 197 (M⁺, 43), 157 (79), 144 (9), 130 (24), 77 (100), 63 (23), 51 (78). 11: yield: 86 %, mp 86.5-88 °C (H₂O : EtOH 1:1). ¹H NMR (CDCl₃): 3.04 (t, 2H), 4.46 (t, 2H), 7.37-7.52 (m, 3H), 8.03-8.15 (m, 2H), 8.20 (s, 1H). MS: 198 (M⁺, 100), 158 (8), 145 (7), 131 (42), 104 (34), 77 (19).
- (a) Yamauchi, M.; Masui, M. Chem. Ind. (London) 1976, 31-32; (b) Sawa, N.; Okamura, S. Nippon Kagaku Zasshi 1969, 90, 704-707; Chem. Abstr. 1969, 71, 101773a.
- 17. Efros, A. M. Zhur. Obschchei Khim. 1960, 30, 3565-3569; Chem. Abstr. 1961, 55, 18712d.
- 18. Compound 5d was obtained in 45 % yield by reaction of urocanic acid ethyl ester with excess MeMgI in ether, mp 133-137 °C (EtOAc). ¹H NMR (DMSO-d₆): 1.24 (s, 6H), 4.59 (OH), 6.22 (d, 1H, *J*= 16 Hz), 6.37 (d, 1H, *J*= 16 Hz), 6.98 (m, 1H), 7.57 (m, 1H). MS: 152 (M⁺, 11), 133 (100). 119 (32).
- 8b: yield: 52 %, colourless oil. ¹H NMR (CDCl₃): 1.18 (s, 9H), 2.27 (d, 3H, J= 1.0 Hz), 5.80 (s, 2H), 6,78 (m, 1H), 7.61 (m, 1H). MS: 196 (M⁺, 7), 95 (20), 94 (28), 57 (100). 9: yield 94 %, mp 121-124 °C (petr. ether : EtOAc 2:1). ¹H NMR (CDCl₃): 2.28 (m, 3H), 5.32 (s, 2H), 6.66 (m, 1H), 7.41 (m, 1H), 7.47-7.73 (m, 3H), 7.99 (m, 2H). MS: 200 (M⁺, 13), 105 (100), 77 (38).
- 20. Leandri, G.; Mangini, A.; Montanari, F.; Passerini, R. Gazz. Chim. Ital. 1955, 85, 769-839.
- 21. Lauth-de Viguerie, N.; Sergueeva, N.; Damiot, M.: Mawlawi, H.; Riviere, M.; Lattes, A. Heterocycles 1994, 37, 1561-1578.
- 22. von Auwers, K.; Schmidt, W. Ber. Dtsch. Chem. Ges. 1925, 58, 528-543.
- 23. Kubota, S.; Uda, M.; Nakagawa, T. J. Heterocycl. Chem. 1975, 12, 855-860.

(Received in UK 12 March 1996; revised 29 April 1996; accepted 3 May 1996)