

Palladium(II)-catalyzed oxidation of acrylate esters to acetals in supercritical carbon dioxide

Lanqi Jia, Huanfeng Jiang* and Jinheng Li

Guangzhou Institute of Chemistry, Chinese Academy of Sciences, PO Box 1122, Guangzhou 510650, China.
E-mail: jhf@mail.gic.ac.cn

Received (in Cambridge, UK) 11th March 1999, Accepted 20th April 1999

The development of a new palladium(II)-catalyzed oxidation of methyl acrylate, affording methyl 3,3-dimethoxypropanoate in excellent conversion and selectivity in supercritical CO₂, is presented.

Alkyl 3,3-dialkoxypropanoates are important intermediates in organic synthesis and have been used to synthesize a variety of compounds, including coumarins,¹ porphyrins,² spermine metabolites³ and loganin.⁴ Previously reported methods for the preparation of alkyl 3,3-dialkoxypropanoates are not satisfactory, due to the high cost of the starting materials and the low overall yields of the process. The Pd^{II}-catalyzed oxidation of terminal olefins with water, which is well known as the Wacker reaction, produces methyl ketones.⁵ A similar reaction of terminal olefins bearing electron-withdrawing groups upon treatment of alcohols catalyzed by Pd^{II} affords acetals at the terminal carbon in DME. High yields of methyl 3,3-dimethoxypropanoate (92%), however, were afforded only in the presence of HMPA. In the absence of HMPA the yield was only 46%.⁶

Using supercritical carbon dioxide (scCO₂) as a reaction solvent has been attracting extensive attention in recent years. First, CO₂ is inexpensive, nonflammable, nontoxic⁷ and chemically inert under many conditions. ScCO₂, as an environmentally friendly reaction medium, may be a substitute for volatile and toxic organic solvents.⁸ Secondly, scCO₂ possesses hybrid properties of both liquid and gas, to the advantage of some reactions involving gaseous reagents.⁹ Control of the solvent density by variation of the temperature and pressure enables the solvent properties to be 'tuned' to reactants.¹⁰ Finally, separating of CO₂ from the reaction mixture is energy-efficient and simple.

Organic reactions in scCO₂ have facilitated great progress in organometallic chemistry, particularly in homogeneous and


heterogeneous catalysis. Several recent reports have shown that some interesting results are specifically associated with the use of scCO₂ as a reaction solvent.¹¹ Here we disclose our work on a Pd^{II}-catalyzed terminal olefin oxidation in scCO₂.[‡] Our results show that Pd^{II}-catalyzed oxidation of methyl acrylate could be successfully processed in scCO₂ to give the dimethyl acetal as major product when an excess of MeOH was used (Scheme 1). Our previous experiments show that it is necessary to add an excess of MeOH to promote partial dissolution of the inorganic catalysts in scCO₂, since MeOH is miscible in scCO₂.¹²

The catalytic activity of PdCl₂ and PdCl₂(MeCN)₂ for oxidation of methyl acrylate were investigated. The reaction catalyzed by PdCl₂ were slightly faster than that by PdCl₂(MeCN)₂ (Table 1, entries 2 and 3).

It has been reported that CuCl was superior to CuCl₂ as the cocatalyst in the oxidation of acrylate esters.^{13,6b} Interestingly, in our experiments carried out in scCO₂, 0.4 equiv. of CuCl₂ gave similar or even better results than the same amount of CuCl (Table 1, entries 8 and 9).

Hosokawa reported that the oxidation of methyl acrylate to the acetal was carried out at 50 °C in DME.⁶ We found that the oxidation reaction could be completed at a lower temperature in scCO₂. In our experiment, the reaction afforded very good conversion and selectivity at 40 °C. The reaction was still smoothly accomplished at 27 °C, at which temperature CO₂ is a liquid, although the amount of byproducts increased slightly.

The CO₂ pressure had a great effect on the selectivity. The best results were obtained when the pressure of CO₂ was controlled between 12–13 MPa. As the pressure of CO₂ was reduced, the conversion and selectivity also declined. For example, the reaction gave **2** in a yield of only 75.8% and the



Scheme 1

Table 1 Oxidation reaction of methyl acrylate in scCO₂^a

| Entry | Catalyst | T/°C | P _{O₂} /MPa | P _{CO₂} /MPa | Conv. | Yield ^b (%) | | | Selectivity for 2 ^c (%) |
|-------|---|------|---------------------------------|----------------------------------|-------|------------------------|----------|----------|---|
| | | | | | | 2 | 3 | 4 | |
| 1 | PdCl ₂ (MeCN) ₂ , CuCl ^d | 50 | 0.1 | 13 | 99.0 | 81.1 | 3.5 | 8.8 | 86.8 |
| 2 | PdCl ₂ (MeCN) ₂ , CuCl ^d | 50 | 0.1 | 13 | 49.4 | 36.6 | 5.9 | 2.7 | 81.0 |
| 3 | PdCl ₂ , CuCl ₂ ^e | 50 | 0.1 | 13 | 71.0 | 42.9 | 6.6 | 3.1 | 81.6 |
| 4 | PdCl ₂ , CuCl ₂ | 50 | 0.1 | 13 | 98.5 | 82.7 | 4.0 | 3.5 | 91.7 |
| 5 | PdCl ₂ (MeCN) ₂ , CuCl | 40 | 0.1 | 12 | 93.0 | 82.4 | 2.8 | 3.0 | 93.4 |
| 6 | PdCl ₂ , CuCl ₂ | 27 | 0.1 | 12 | 93.0 | 75.0 | 5.1 | 2.9 | 90.3 |
| 7 | PdCl ₂ , CuCl ₂ | 40 | 0.1 | 9 | 92.0 | 75.8 | 2.8 | 3.4 | 92.4 |
| 8 | PdCl ₂ , CuCl ₂ | 40 | 0.5 | 13 | 99.1 | 87.9 | 1.6 | 2.5 | 95.5 |
| 9 | PdCl ₂ , CuCl | 40 | 0.5 | 13 | 99.7 | 86.3 | 1.5 | 4.5 | 93.5 |
| 10 | PdCl ₂ , CuCl ₂ | 40 | 1.0 | 12 | 99.4 | 91.7 | 1.2 | 2.0 | 96.6 |

^a Methyl acrylate (5 mmol), MeOH (1 ml ≈ 24.7 mmol), PdCl₂ or PdCl₂(MeCN)₂ (3 mol%), CuCl or CuCl₂ (0.4 equiv.), 12 h. ^b By GC. ^c Selectivity = [**2**/(**2** + **3** + **4**)] × 100. ^d PdCl₂(MeCN)₂ (1 mol%), CuCl or CuCl₂ (2 mol%). ^e CuCl₂ (2 mol%).

amount of other unidentified products increased when 9 MPa CO₂ was charged in the autoclave.

The oxygen pressure also affected the conversion and selectivity. Lower oxygen pressure (0.1 MPa) gave satisfactory conversion but the selectivity to **2** was never higher than 93.4%. The best selectivity (over 96%) was obtained as the oxygen pressure was elevated to 1.0 MPa. Using ethyl acrylate as the substrate, the reaction affords similar results, and the further experiments are now in progress.

In summary, the oxidation reaction of methyl acrylate afforded the acetal in high conversion and selectivity at lower temperature when supercritical carbon dioxide instead of DME was used as the reaction medium without the presence of HMPA. The pressure of CO₂ and O₂ had a remarkable effect on the conversion and selectivity.

The authors are grateful to the National Natural Science Foundation of China for financial support of this work (29772036) and Dr Fanglü Huang for helpful discussions.

Notes and references

† Typical procedure: Reactions were carried out in a HF-25 autoclave. Pd^{II} catalyst (0.15 mmol, 3 mol%), CuCl₂ (4 mmol), MeOH (1 ml, 24.7 mmol) and acrylate ester (5 mmol) were added into a 25 ml autoclave in sequence. O₂ and then liquid CO₂ were pumped into the autoclave using a cooling pump to give the desired pressure. The autoclave was then put into an oil bath under magnetic stirring for the desired reaction time. After the reaction, the autoclave was allowed to cool to -30 °C. CO₂ was vented and the surplus was extracted with n-hexane. The extract was filtrated and condensed under reduced pressure. The product was analyzed using GC

(quantitative analysis) and GC-MS, ¹H NMR ¹³C NMR and IR analysis (identification of products).

- 1 D. G. Crosby and R. V. Berthold, *J. Org. Chem.*, 1962, **27**, 3083.
- 2 J. P. Collman, A. O. Chong, G. B. Jameson, R. T. Oakley, E. Rose, E. R. Schmittou and J. A. Ibers, *J. Am. Chem. Soc.*, 1981, **103**, 516.
- 3 M. Israel, E. C. Zoll, N. Muhammad and E. J. Modest, *J. Med. Chem.*, 1973, **16**, 1.
- 4 G. Büchi, J. A. Carlson, J. E. Powell, Jr. and L.-F. Titzze, *J. Am. Chem. Soc.*, 1973, **95**, 504.
- 5 For a comprehensive review, see: J. Tsuji, *Synthesis*, 1984, 369.
- 6 (a) T. Hosokawa, S. Aoki and S.-I. Murahashi, *Synthesis*, 1992, 558; (b) T. Hosokawa, T. Ohta, S. Kanayama and S.-I. Murahashi, *Synthesis*, 1992, 558; (b) T. Hosokawa, T. Ohta, S. Kanayama and S.-I. Murahashi, *J. Org. Chem.*, 1987, **52**, 1758.
- 7 For discussions of the advantages of scCO₂, see T. Clifford and K. Bartle, *Chem. Ind.*, 1996, 449; M. Poliakov and S. Howdle, *Chem. Ber.*, 1995, **31**, 118; H. Black, *Environ. Sci. Technol.*, 1996, **30**, 124A.
- 8 D. A. Morgenstern, R. M. LeLacheur, D. K. Morita, S. L. Borokowsky, S. Feng, G. H. Brown, L. Luan, M. F. Gross, M. J. Burk and W. Tumas, *ACS Symp. Ser.*, 1996, **626**, 132.
- 9 J. W. Rathke, R. J. Klinger and T. R. Krause, *Organometallics*, 1991, **10**, 1350; P. G. Jessop, T. Ikariya and R. Noyori, *Nature*, 1994, **368**, 231.
- 10 M. A. Carroll and A. B. Holmes, *Chem. Commun.*, 1998, 1395.
- 11 M. J. Burk, S. Feng, M. F. Gross and W. Tumas, *J. Am. Chem. Soc.*, 1995, **117**, 8277.
- 12 L. Jia, H. Jiang and J. Li, *Green Chem.*, 1999, 91.
- 13 J. Tsuji, H. Nagashima and H. Nemoto, *Org. Synth.*, 1984, **62**, 9.

Communication 9/019351