

Preliminary Note

Synthesis and reactions of *N,N*-dichloroperfluoroalkanesulfonylamides

Shi-zheng Zhu*, Cheng-ming Zhou, Ai-wen Li and Bin Xu

Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Lu, Shanghai 200032 (China)





(Received June 4, 1993; accepted August 29, 1993)

Abstract

Reactions of *N,N*-dichloroperfluoroalkanesulfonylamides with alkenes, benzene, dimethyl sulfide and dimethyl sulfoxide in the presence of zinc dust gave addition or insertion products via the fluoroalkanesulfonylnitrene intermediate. On heating $R_fSO_2NCl_2$ in the presence of alkene without zinc, 1:1 addition products were formed via a free-radical mechanism.

N,N-Dichlorotrifluoromethanesulfonylamide, $CF_3SO_2NCl_2$, was first prepared in 1974 [1]. Its chemistry, however, has received little study. Recently, Yagupol'skii *et al.* reported its reactions with diphenyl sulfide [2] and trifluoromethylphenyl sulfide [3].


Here, we report the preparation of some *N,N*-dichloroalkanesulfonylamides and their reactions with benzene, dimethyl sulfide, dimethyl sulfoxide and alkenes. The title compounds were conveniently prepared by the one-pot reaction of fluoroalkanesulfonylamides with KOH(aq.) and chlorine gas:



Compounds **2** were obtained as yellowish liquids. They are unstable, for example, after storage for 1 week in a flask at room temperature with nearly 50% of **2** decomposing to the corresponding fluoroalkanesulfonylamides (detected by ^{19}F NMR spectroscopy: the chemical shifts of $-CF_2S$ in compounds **1** and **2** were 40 ppm and 29 ppm, respectively; TFA external

standard; upfield positive). Heating **2** with benzene in the presence of zinc dust afforded 70% of $R_fSO_2NHC_6H_5$ (**3**). Similarly, treatment of **2** with dimethyl sulfide or dimethyl sulfoxide gave the sulfonium ylid $R_fSO_2N=SMe_2$ (**4**) and the sulfoxonium ylid $R_fSO_2N=S(O)Me_2$ (**5**), respectively. The *N*-fluoroalkanesulfonylaziridines **6** were obtained by refluxing **2** with styrene or 1,4-dimethyl-but-2-ene in CH_2Cl_2 . In these two reactions, $R_fSO_2NH_2$ was also formed in c. 15% yield.

These results appear to indicate that fluoroalkanesulfonylnitrene intermediates are involved in all the above questions [4, 5]. This is similar to the reactions of fluoroalkanesulfonylazides [6] (see Scheme 1).




[$R_1=R_2=R_3=H$, $R_4=C_6H_5$ (**6a**); $R_1=R_2=R_3=R_4=CH_3$ (**6b**)]

Reagents, conditions and yields: i, C_6H_6 , reflux, 8 h, 70%; ii, Me_2S , r.t., 8 h, 68%; iii, DMSO, 60 $^\circ\text{C}$, 8 h, 65%; iv, $PhCH=CH_2$, CH_2Cl_2 , r.t., 8 h, 62%, $Me_2C=CMe_2$, r.t., 8 h, 64%.

Scheme 1.


It was noteworthy that the reaction of **2** with styrene occurred rapidly without zinc powder, and gave 1:1 addition products:



[$R_1=H$; $R_2=C_6H_5$]

This reaction was demonstrated to be a free-radical reaction process since the radical intermediate $R_fSO_2NCICH_2CHPh$ could be captured by $Bu^{\bullet}NO$. The ESR spectrum showed triplet-doublet peaks ($a_N = 15.18$ G, $a_H = 3.04$ G, $g = 2.0052$).

Reduction of **8** by $NaHSO_3$, followed by elimination of HCl with alcoholic NaOH gave *N*-fluoroalkanesulfonylaziridine (**6a**) [7]:



*Author to whom correspondence should be addressed.

All the new compounds were fully characterized by their IR, ^1H NMR, ^{19}F NMR, MS spectra and by elemental analysis.

Further studies on the chemistry of **2** are in progress.

Experimental

General procedure for the preparation of *N,N*-dichloroperfluoroalkanesulfonylamides **2**

Perfluoroalkanesulfonylamides **1a** (6.3 g, 15 mmol), KOH (1.7 g, 30 mmol) and water (15 ml) were placed in a 50 ml three-necked flask and the mixture stirred at room temperature for 2 h. The flask was cooled down via an ice bath, then chlorine gas (4.1 g, 60 mmol) was introduced during which time the reaction temperature was kept at 0–5 °C. The reaction mixture was then stirred for another 2 h at 20 °C. A yellowish oil separated which was dried over Na_2SO_4 and distilled under vacuum to give **2a** (3.8 g, 52%), b.p. 42–44 °C/2 mmHg. IR (film) (ν_{max} , cm^{-1}): 1605 (w); 1420 (m); 1380 (s); 1320 (vs); 1292 (s); 1220–1120 (vs); 1092 (s); 990 (s); 713 (s); 600 (m); 530 (m). ^{19}F NMR (neat) (TFA external, upfield positive) δ : –11.8 (s, ICF_2); 5.20 (m, CF_2); 8.5 (m, CF_2O); 29.3 (s, CF_2S) ppm. MS m/z %: 492 (MH^+ , 0.62); 424 ($\text{M}^+ - \text{F} - \text{SO}$, 1.90); 364 ($\text{M}^+ - \text{I}$, 0.96); 343 ($\text{M}^+ - \text{SO} - \text{C}_2\text{F}_4$, 5.52); 329 ($\text{M}^+ - \text{SON} - \text{C}_2\text{F}_4$, 4.82); 227 (IC_2F_4^+ , 100); 205 (ISO_2N^+ , 15.7). Analysis: Calc. for $\text{C}_4\text{Cl}_2\text{F}_8\text{INO}_3\text{S}$: C, 9.76; N, 2.85; F, 30.89%. Found: C, 10.04; N, 2.99; F, 31.30%.

Typical reaction of **2** in the presence of zinc powder

A mixture of **2b** (2 g, 5 mmol), DMSO (10 ml) and zinc powder (1 g, 15 mmol) was stirred for 8 h at 60 °C. After filtration, the filtrate was fractionally distilled

under reduced pressure giving **5b** (1.3 g, 64%); b.p. 98–100 °C/2 mmHg. IR (film) (ν_{max} , cm^{-1}): 2971 (s); 2892 (s); 1580 (m); 1443 (m); 1380 (s); 1370 (s); 1277 (s); 1210–1100 (vs); 1005 (vs); 980 (s); 881 (m); 700 (m); 653 (s); 517 (s). ^1H NMR δ : 3.50 (s, 2CH_3) ppm. ^{19}F NMR δ : –1.0 (s, ClCF_2); 6.5 (m, OCF_2); 11.6 (m, CF_2O); 42.3 (s, CF_2S) ppm. MS m/z %: 408/410 (MH^+ , 0.75/0.37); 392/394 ($\text{M}^+ - \text{H} - \text{O}$, 25.96/16.09); 372 ($\text{M}^+ - \text{Cl}$, 2.64); 344/346 ($\text{M}^+ - \text{H} - \text{SO}_2$, 4.34/2.72); 332/334 ($\text{M}^+ - \text{H} - \text{SMe}_2 - \text{N}$, 20.22/8.40); 156 ($\text{M}^+ - \text{R}_f$, 11.09); 141 ($\text{M}^+ - \text{R}_f - \text{Me}$, 4.99); 140 ($\text{M}^+ - \text{R}_f - \text{O}$, 17.32); 135/137 (ClC_2F_4^+ , 15.39/7.45); 85/87 (ClCF_2^+ , 14.72/6.37); 80 (SOS^+ , 100); 78 (Me_2SO^+ , 5.74); 64 (SO_2^+ , 85.47); 61 (MeSN^+ , 12.90); 47 (MeS^+ , 5.14). Analysis: Calc. for $\text{C}_6\text{H}_6\text{F}_8\text{ClNO}_4\text{S}_2$: C, 17.67; H, 1.47; N, 3.44; F, 37.30%. Found: C, 18.03; H, 1.80; N, 3.13; F, 37.00%.

Acknowledgement

The authors wish to thank the Shanghai National Science and Technology Foundation for financial support.

References

- 1 V.P. Nazaretyan, O.A. Radchenko and L.M. Yagupol'skii, *Zh. Org. Khim.*, 10 (1974) 2460.
- 2 R.Yu. Gavrilova, N.V. Kondratenko, V.I. Popov and L.M. Yagupol'skii, *Zh. Org. Khim.*, 25 (1989) 1128.
- 3 N.V. Kondratenko, R.Yu. Gavrilova and L.M. Yagupol'skii, *Zh. Org. Khim.*, 24 (1988) 456.
- 4 R.A. Abramovitch, T.D. Bailey, T. Takaya and V. Uma, *J. Org. Chem.*, 39 (1974) 340.
- 5 D.S. Breslow and M.F. Sloan, *Tetrahedron Lett.*, (1968) 5349.
- 6 S.-z. Zhu, *Tetrahedron Lett.*, 33 (1992) 6503.
- 7 U.K. Nadir, R.L. Sharma and V.K. Koul, *J. Chem. Soc., Perkin Trans. I*, (1991) 2015.