

Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcar20>

SOME UNEXPECTED RESULTS IN DONDONI'S ONE-CARBON HOMOLOGATION PROCEDURE

José Marco-Contelles^a & Elsa de Opazo^a

^a Laboratorio de Radicales Libres (CSIC), C/Juan de la Cierva 3, Madrid, 28006, Spain

Published online: 16 Aug 2006.

To cite this article: José Marco-Contelles & Elsa de Opazo (2001) SOME UNEXPECTED RESULTS IN DONDONI'S ONE-CARBON HOMOLOGATION PROCEDURE, Journal of Carbohydrate Chemistry, 20:7-8, 637-646

To link to this article: <http://dx.doi.org/10.1081/CAR-100108278>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SOME UNEXPECTED RESULTS IN DONDONI'S ONE-CARBON HOMOLOGATION PROCEDURE¹

José Marco-Contelles* and Elsa de Opazo

Laboratorio de Radicales Libres (CSIC),
C/ Juan de la Cierva 3, 28006-Madrid, Spain

ABSTRACT

The formation of the unexpected secondary products (**3** and **9**) during Dondoni's one-carbon homologation of 6-deoxy-6-iodo-2,3:4,5-bis-*O*-isopropylidene-D-glucose (**1**) are described.


INTRODUCTION

Dondoni's one-carbon homologation procedure is one of the most successful protocols described to date for the homologation of aldehydes.² The mild and simple experimental conditions enhance the interest and synthetic potential of this efficient and reproducible process. In addition, the highly *anti*-stereoselective³ asymmetric induction applied to α -alkoxy aldehydes has been used in a number of synthetic approaches for the preparation of enantiomerically pure products.⁴ The mechanism of this useful transformation has been discussed by Dondoni and co-workers (Scheme 1).⁵

Very recently we wished to carry out a one-carbon homologation of 6-deoxy-6-iodo-2,3:4,5-bis-*O*-isopropylidene-D-glucose (**1**) and we chose Dondoni's method.² In this communication we describe in full the unexpected, secondary products obtained in a three-step protocol for the addition of 2-trimethylsilylthiazole (2-TST) to the aldehyde carbonyl of **1**.⁶


RESULTS AND DISCUSSION

Aldehyde **1** has been prepared in enantiomerically pure form and multigram quantities from commercially available D-glucose diethyl dithioacetal.⁷ The reac-




Scheme 1. Proposed mechanism for the 2-TST coupling with aldehydes (ref. 5).

tion of this compound with 2-TST under standard conditions³ afforded the expected 6-deoxy-6-iodo-2,3:4,5-bis-*O*-isopropylidene-1-(1',3'-thiazol-2'-yl)-D-glycero-D-gulose (**2**, 60%),⁷ and a secondary product (3a*R*,3b*S*,6a*R*,7a*R*)-tetrahydro-2,2,5,5-tetramethyl-6a*H*-cyclopenta[1,2-d:3,4-d']bis[1,3]dioxole-6a-carboxaldehyde (**3**) in 7% yield (Scheme 2). The structure of the latter product was firmly established after inspection of the analytical and spectroscopic data. In the ¹H NMR spectrum of **3** we observed a singlet aldehydic proton at 9.73 ppm (198.4 ppm for H—CO in the ¹³C NMR spectrum) and two significant protons at 2.50 ppm (dd, *J*_{5A,5B} = 15.2 Hz, *J*_{4,5bB} = 2.9 Hz, 1 H, H5B) and 2.17 ppm (dd, *J*_{5A,5B} = 15.2 Hz, *J*_{4,5A} = 6.2 Hz, 1 H, H5A) (C5: 39.9 ppm) forming the AB part of an ABX system with a proton at 4.91 ppm (td, *J*_{4,5B} = 2.9 Hz, *J*_{4,5A} = 6.2 Hz, 1 H, H4). In the IR spectrum of **3** no carbonyl band absorption was observed, the carbonyl function being in the hydrated form. The absolute configuration at C1 was easily assigned as *R* by NOE experiments. These data along with the elemental analysis and the MS spectral results clearly pointed out that product **3** contained a cy-



Scheme 2. Reaction of 2-TST with α-alkoxy aldehyde **1**.





Scheme 3. Possible mechanisms for the intramolecular alkylation of α -alkoxy aldehyde **1**.

cllopentane ring system resulting from the intramolecular α -alkylation of the α -alkoxy aldehyde **1**. Additional proof of structure was obtained upon reduction of compound **3** with sodium borohydride to give alcohol **4**, which was then acetylated to yield product **5** (Scheme 2).

Finally, we tested the direct alkylation of compound **1** with DBU (1.1 equiv) as base, under the same experimental conditions (Scheme 3). Very interestingly, a rapid reaction ensued (1 h) giving cleanly compound **3** in 75% isolated yield after flash chromatography.

It is noteworthy that compounds **3–5** are densely functionalized, chiral cyclopentane derivatives with some of the structural and functional moieties present in carbocyclic nucleosides⁸ and glycosidase inhibitors such as trehazolin aglycon (Figure 1).⁹

The present reactivity was completely unexpected and unprecedented. As it is well known, base-mediated alkylation of aldehydes is hampered by crotonization or polymerization due to aldol condensation-like competitive reactions. A series of aldehyde derivatives (imines, hydrazones, oximes or enamines) have been




Figure 1.




used for this purpose,¹⁰ whereas a literature search for α -alkylation of α -alkoxy aldehydes gave a limited number of examples.¹¹ In addition, a recent synthesis of chiral α -alkyl, α -trifluoromethyl α -alkoxy aldehydes used an indirect approach from ketones,^{12a} and an elegant synthesis of spirocyclopentenones from α -alkoxy aldehydes in furanose templates was only possible through enamine intermediates.^{12b} This is in sharp contrast with the α -alkoxy ketones which have proved to be excellent intermediates for aldol reactions *via* the enol borinates.¹³

In Scheme 3 we show two possible mechanisms to explain these results: (a) An intramolecular mechanism, where the presumed intermediate **A**, after proton abstraction gives the trimethylsilyl enol ether **B**, the real precursor for the intramolecular alkylation. Note that intermediate **A** can also yield the major product (**2**) in this reaction,¹⁴ and (b) Alternatively, the intermolecular 2-TST mediated abstraction of the α -proton in compound **1** could also afford the reactive intermediate **B** leading to compound **3**.

In any case β -elimination in intermediate **B** (Scheme 3) was not observed. This is really exceptional, considering the experimental conditions used (room temperature with 2-TST as the only reagent in the medium) compared with the necessary controlled conditions [low temperature reaction (-78°C), LDA as base, co-solvents such as HMPT or DMPU, and highly reactive alkylating agents] for the intermolecular alkylation of tartaric acid derivatives described by Seebach.¹⁵ In the present case, the stability of the enolate may also be due to the rigid acetonide skeleton which precludes the optimal geometrical arrangement for β -elimination to take place.¹⁵

In summary, we have reported the first example of an intramolecular alkylation of an ε -iodo α -alkoxy aldehyde, by using basic, extremely simple and mild reaction conditions. These results open the way for new exciting synthetic ventures directed to the important, synthetic intermolecular alkylation of chiral α -heteroatom *O* or *N*-substituted aldehydes, as a new strategy for the synthesis of enantiomerically pure molecules containing quaternary centers.¹⁶

Continuing with Dondoni's protocol for the synthesis of aldehyde **7** from intermediate **2** (Scheme 4), we first *O*-benzylated the free alcohol **2** to give compound **8**,⁷ and then submitted it to sequential reactions² with methyl iodide, sodium borohydride and mercuric chloride to give the desired product **7**, in 50% yield, and compound **9**, in 5% overall yield from product **8** (Scheme 4).




Scheme 4. Synthesis of compounds **7** and **9** from intermediate **8**.



The unexpected compound **9** was an extremely unstable product, and no reliable elemental analysis of the product could be obtained. However, the spectroscopic data strongly supported this structure. The IR spectrum of **9** contained a strong carbonyl ester band at 1760 cm^{-1} . In the ^1H NMR spectrum of compound **9** we could assign the same protons, with similar chemical shifts, as those bonded to carbons C3-C8, in compounds **7** or **8**: the iodide atom and the two isopropylidene groups were still present, but the thiazole nucleus was absent. The phenylmethyl protons were more shifted in compound **9** (ca. 0.6 ppm) than in the precursor **8**.⁷ These IR and ^1H NMR data were supported in the ^{13}C NMR spectrum plus the results from a DEPT experiment;⁹ a signal at 170.4 ppm was assigned to the ester carbonyl group, and the methylene carbon bearing the iodide appeared at 1.4 ppm. Finally, the mass spectrum of **9** showed an intense peak at m/z 461, suggesting that the structure of compound **9** corresponded to the benzylic ester of the oxidized form of aldehyde **1** (Scheme 2).

Looking very carefully at the three steps protocol for Dondoni's homologation of product **8**, we could see that this secondary product was already present in the first step of the protocol, the reaction with methyl iodide, and that it survived untouched the mild and quick reduction with sodium borohydride and the mercuric mediated deprotection of the masked aldehyde. To the best of our knowledge a similar product has not been documented in the reported Dondoni's systematic application of this method.² In fact, it is difficult to rationalize the formation of this molecule in the reaction with methyl iodide until formation of the quaternary salt!

In Scheme 5 we show a mechanism that we tentatively propose for the formation of ester **9**. Obviously, the equilibrium between **8** and **8''** is displaced to the



Scheme 5. Possible mechanism for the formation of compound **9** from intermediate **8**.



left, to the most stable aromatic “imine” **8**. However, we think that enamine **8** may react with small traces of water to give intermediate **A**, affording ester **9** after elimination of thiazolidine **B**.

In summary, in this report we have described the successful homologation of aldehyde **1** to aldehyde **7** using Dondoni’s method, showing also the formation of some secondary, unexpected products (**3** and **9**) in this protocol. These results enhance the generality of the method, but also point out the possible formation of secondary products when designing synthetic schemes using this methodology.

EXPERIMENTAL

General Methods. Reactions were monitored by TLC using precoated silica gel aluminium plates containing a fluorescent indicator (Merck, 5539). Detection was done by UV (254 nm) followed by charring with sulfuric-acetic acid spray, 1% aqueous potassium permanganate solution or 0.5% phosphomolybdic acid in 95% EtOH. Anhydrous Na₂SO₄ was used to dry organic solutions during work-ups and the removal of solvents was carried out under vacuum with a rotary evaporator. Flash column chromatography was performed using silica gel 60 (230–400 mesh, Merck) and hexane/ ethyl acetate mixtures as eluent unless otherwise stated. ¹H spectra were recorded with a Varian VXR-300(400)S spectrometer, using tetramethylsilane as internal standard and ¹³C NMR spectra were recorded with a Bruker WP-200-SY.

(3a*R*,3b*S*,6a*R*,7a*R*)-Tetrahydro-2,2,5,5-tetramethyl-6a*H*-cyclopenta[1,2-*d*:3,4-*d'*]bis[1,3]dioxole-6a-carboxaldehyde (3**).** *Method (A).* Compound **1** (197 mg, 0.53 mmol) dissolved in dry methylene chloride (1 mL) was treated with 2-TST (0.094 mL, 0.58 mmol, 1.1 equiv) at 0°C, under argon. The reaction mixture was warmed and stirred at room temperature for 18 h. The solvent was removed, and the residue was treated with Bu₄NF (0.53 mL, 1.1 M in THF) until desilylation was complete. The solvent was evaporated, the residue diluted with water and extracted with methylene chloride several times. The organic layer was dried, filtered and concentrated. Flash chromatography (hexane/ethyl acetate 10%) gave compounds **2**⁷ (141.4 mg, 60%) and **3** (9.3 mg, 7%). *Method (B).* Aldehyde **1** (104.4 mg, 0.28 mmol) was dissolved in dry CH₂Cl₂ (1 mL, 0.3 M), cooled at 0 °C and treated with DBU (0.05 mL, 0.31 mmol, 1.1 equiv). The mixture was stirred and warmed at rt for 1 h. The solvent was removed and the crude product was submitted to chromatography (hexane/ethyl acetate, 9/1) to give compound **3** (51 mg, 75%): oil; [α]_D²⁵ – 2 (c 0.7, CHCl₃); IR (film) ν 3470, 2986, 2942, 1744, 1644, 1376, 1254, 1166, 1060 cm^{–1}; ¹H NMR (300 MHz, CDCl₃) δ 9.73 (s, 1 H, CHO), 4.91 (td, $J_{4,5B}$ = $J_{4,3}$ = 6.2 Hz, $J_{4,5A}$ = 2.9 Hz, 1 H, H₄), 4.63 (d, $J_{3,4}$ = 6.2 Hz, 1 H, H₃), 4.61 (s, 1 H, H₂), 2.50 (dd, $J_{5A,5B}$ = 15.2 Hz, $J_{4,5A}$ = 2.9 Hz, 1 H, H_{5A}), 2.17 (dd, $J_{5A,5B}$ = 15.2 Hz, $J_{4,5B}$ = 6.2 Hz, 1 H, H_{5B}), 1.52, 1.46, 1.38, 1.31 [4 s, 12 H, 2 × OC(CH₃)₂O]; ¹³C NMR (75 MHz, CDCl₃) δ 198.4 (CHO), 113.5,



111.1 [$2 \times \text{OC}(\text{CH}_3)_2\text{O}$], 97.3 (C1), 86.6, 83.6 (C2, C3), 81.2 (C4), 39.9 (C5), 27.5, 26.7, 25.9, 24.7 [$2 \times \text{OC}(\text{CH}_3)_2\text{O}$]; MS (70 eV) m/z 243 ($\text{M}+1^+$, 19), 227 (M^+-15 , 64), 43 (100).

Anal. Calcd for $\text{C}_{12}\text{H}_{18}\text{O}_5$ (242.27): C, 52.94; H, 7.49. Found: C, 52.75; H, 7.38.

(3aR,3bS,6aR,7aR)-Tetrahydro-2,2,5,5-tetramethyl-6aH-cyclopenta[1,2-d:3,4-d']bis[1,3]dioxole-6a-methanol (4). To a solution of compound **3** (84.9 mg, 0.35 mmol) in methanol (2.1 mL, 0.17 M), cooled at 0°C , sodium borohydride (20 mg, 0.52 mmol, 1.5 equiv) was added, and the mixture was warmed at rt and stirred for 30 min. Acetone (0.5 mL) was added and the solvents removed by evaporation. The crude product was diluted with methylene chloride, washed with brine, dried and filtered. The solvent was evaporated and the residue was submitted to chromatography (hexane/ethyl acetate, 4/1) to give alcohol **4** (73 mg, 85%): oil; $[\alpha]_{\text{D}}^{25} - 13$ (c 0.47, CHCl_3); IR (film) ν 3468, 2986, 2944, 1380, 1210, 1168, 1058 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 4.81 (ddd, $J_{3,4} = 5.9$ Hz, $J_{4,5A} = 4.0$ Hz, $J_{4,5B} = 2.9$ Hz, 1 H, H4), 4.62 (d, $J_{3,4} = 5.9$ Hz, 1 H, H3), 4.40 (s, 1 H, H2), 3.78 (dd, $J_{1'A,1'B} = 12.1$ Hz, $J_{1'A,\text{OH}} = 4.9$ Hz, 1 H, H1'A), 3.69 (dd, $J_{1'A,1'B} = 12.1$ Hz, $J_{1'B,\text{OH}} = 8.2$ Hz, 1 H, H1'B), 2.18 (dd, $J_{1'B,\text{OH}} = 8.2$ Hz, $J_{1'A,\text{OH}} = 4.9$ Hz, 1 H, OH), 2.16 (d, $J_{5A,4} = 4.0$ Hz, 1 H, H5A), 2.10 (d, $J_{5B,4} = 2.9$ Hz, 1 H, H5B), 1.48, 1.46, 1.37, 1.29 [4 s, 12 H, $2 \times \text{OC}(\text{CH}_3)_2\text{O}$]; ^{13}C NMR (75 MHz, CDCl_3) δ 110.8, 109.8 [$2 \times \text{OC}(\text{CH}_3)_2\text{O}$], 92.1, (C1), 84.7 (C2), 82.3 (C3), 80.2 (C4), 65.0 (C1'), 41.2 (C5), 28.1, 26.4, 26.0, 23.6 [$2 \times \text{OC}(\text{CH}_3)_2\text{O}$]; MS (70 eV) m/z 229 ($\text{M}-15^+$, 100), 213 (4), 155 (10), 111 (56), 99 (17), 43 (74).

Anal. Calcd for $\text{C}_{12}\text{H}_{20}\text{O}_5$ (244.29): C, 59.00; H, 8.25. Found: C, 59.15; H, 8.36.

(3aR,3bS,6aR,7aR)-Tetrahydro-2,2,5,5-tetramethyl-6aH-cyclopenta[1,2-d:3,4-d']bis[1,3]dioxole-6a-methyl Acetate (5). Alcohol **4** (32 mg, 0.13 mmol) was treated with a mixture of acetic anhydride/pyridine (1 mL, 1/1) at rt overnight. The solvent was evaporated and the crude product was submitted to chromatography (hexane/ethyl acetate, 83/17) to give acetate **5** (35 mg, 93%): oil; $[\alpha]_{\text{D}}^{25} - 4$ (c 0.51, CHCl_3); IR (film) ν 2989, 2938, 1747, 1382, 1211, 1169, 1062 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 4.79 (dt, $J_{3,4} = J_{4,5B} = 5.6$ Hz, $J_{4,5A} = 1.5$ Hz, 1 H, H4), 4.59 (d, $J_{3,4} = 5.7$ Hz, 1 H, H3), 4.37/4.15 (2 d, $J = 12.1$ Hz, 2 H, 2 H1'), 4.32 (s, 1 H, H2), 2.17 (br d, $J_{5A,5B} = 15.0$ Hz, 1 H, H5A), 2.09 (s, 3 H, OCOCH_3), 2.07 (dd, $J_{5A,5B} = 15.0$ Hz, $J_{5B,4} = 5.6$ Hz, 1 H, H5B), 1.44 (s, 6 H, 1.37, 1.29 [3 s, 12 H, $2 \times \text{OC}(\text{CH}_3)_2\text{O}$]; ^{13}C NMR (75 MHz, CDCl_3) δ 170.1 (OCOCH_3), 111.2, 109.9 [$2 \times \text{OC}(\text{CH}_3)_2\text{O}$], 89.9, (C1), 85.0 (C2), 82.2 (C3), 80.2 (C4), 65.8 (C1'), 41.5 (C5), 27.9, 26.0 (2 C), 23.6 [$2 \times \text{OC}(\text{CH}_3)_2\text{O}$], 20.9 (OCOCH_3); MS (70 eV) m/z 286 (M, 1), 271 ($\text{M}-15^+$, 72), 228 (8), 171 (10), 153 (34), 111 (66), 81 (33), 59 (20), 43 (100).

Anal. Calcd for $\text{C}_{14}\text{H}_{22}\text{O}_6$ (286.32): C, 58.73; H, 7.74. Found: C, 58.65; H, 7.63.



2-*O*-Benzyl-7-deoxy-7-iodo-3,4:5,6-bis-*O*-isopropylidene-D-glycero-D-gulo-heptose (7) and Benzyl 6-deoxy-6-iodo-2,3:4,5-bis-*O*-isopropylidene-D-gluconoate (9). Product **8**⁷ (930 mg, 1.7 mmol) was dissolved in acetonitrile (10 mL) and the solution treated with methyl iodide (0.53 mL, 9.5 mmol, 15 equiv) at 80 °C (bath temperature) for 24 h in a sealed tube. The reaction was cooled, the solvent evaporated and the residue was dissolved in methanol (10 mL), cooled at 0 °C and treated with sodium borohydride (96.7 mg, 2.55 mmol, 1.5 equiv) under vigorous stirring for 30 min. Acetone was then added to destroy the excess of reagent, the solvent was evaporated in vacuo, and the crude product was dissolved in methylene chloride, washed with brine, dried, filtered, and the solvent was evaporated. The residue was dissolved in acetonitrile (1 mL) and to this solution, a solution of mercuric chloride (562 mg, 2.04 mmol, 1.2 equiv) in a mixture of acetonitrile/water (7.4 mL, 4/1) was slowly added. The mixture was stirred at rt for 15 min. The suspension was filtered over celite, which was washed with ethyl acetate. The organic layer was washed with brine, dried, filtered, concentrated and the residue was submitted to chromatography (hexane/ethyl acetate, 84/16) to give compounds **7** (420 mg, 50%) and **9** (30 mg, 5%).

7: oil; ¹H NMR (300 MHz, CDCl₃) δ 9.71 (d, *J*_{1,2} = 2.1 Hz, 1 H, H-1), 7.36/7.27 (m, 5 H, OCH₂C₆H₅), 4.76/4.60 (AB system, *J* = 11.7 Hz, 2 H, OCH₂C₆H₅), 4.54 (q, *J*_{5,6} = *J*_{6,7} = *J*_{6,7'} = 7.0 Hz, 1 H, H-6), 4.32 (dd, *J*_{3,4} = 7.9 Hz, *J*_{2,3} = 5.9 Hz, 1 H, H-3), 4.13 (q, *J*_{4,5} = 10.4 Hz, *J*_{3,4} = 7.9 Hz, 1 H, H-4), 4.12 (dd, *J*_{4,5} = 10.4 Hz, *J*_{5,6} = 7.0 Hz, 1 H, H-5), 3.88 (dd, *J*_{2,3} = 5.9 Hz, *J*_{1,2} = 2.1 Hz 1 H, H-2), 3.37–3.34 (m, 2 H, 2 H-7), 1.50, 1.43, 1.37, 1.35 [4 s, 2 × OC(CH₃)₂O].

Anal. Calcd for C₂₀H₂₇IO₆ (490.33): C, 48.99; H, 5.55. Found: C, 48.76; H, 5.67.

9: oil; [α]_D²⁵ – 21 (c 0.34, CHCl₃); IR (film) *ν* 2987, 2936, 1760, 1498, 1456, 1381, 1214, 1100, 1062 cm^{–1}; ¹H NMR (300 MHz, CDCl₃) δ 7.37–7.32 (m, 5 H, CO₂CH₂C₆H₅), 5.26/5.23 (2 d, *J* = 12.1 Hz, 2 H, CO₂CH₂C₆H₅), 4.57 (d, *J*_{2,3} = 8.1 Hz, 1 H, H2), 4.53 (q, *J*_{5,6A} = *J*_{5,6B} = *J*_{5B,4} = 7.0 Hz, 1 H, H5), 4.33 (dd, *J*_{3,4} = 1.5 Hz, *J*_{4,5} = 6.7 Hz, 1 H, H4), 4.25 (dd, *J*_{3,4} = 1.5 Hz, *J*_{3,2} = 8.0 Hz, 1 H, H3), 3.39 (dd, *J*_{6A,6B} = 10.1 Hz, *J*_{6A,5} = 6.7 Hz, 1 H, H6A), 3.33 (dd, *J*_{6A,6B} = 10.1 Hz, *J*_{6B,5} = 7.7 Hz, 1 H, H5B), 1.49, 1.48 1.42, 1.36 [3 s, 12 H, 2 × OC(CH₃)₂O]; ¹³C NMR (75 MHz, CDCl₃) δ 170.4 (CO₂CH₂C₆H₅), 135.3–128.5 (CO₂CH₂C₆H₅), 111.9, 109.6 [2 × OC(CH₃)₂O], 77.8, (C5), 76.8 (C3), 75.7 (C2), 75.2 (C4), 67.1 (CO₂CH₂C₆H₅), 26.8, 26.6, 25.9, 25.5 [2 × OC(CH₃)₂O], 1.4 (C6); MS (70 eV) *m/z* 461 (M-15⁺, 63), 283 (17), 235 (38), 183 (18), 91 (100).

HRMS: *m/z* Calcd for C₁₉H₂₅IO₆: 476.0695. Found: 476.0643.

ACKNOWLEDGMENTS

We thank Professor A. Dondoni for useful interchange of information and discussion. This was supported by COST Action No. D12 (European Union) and Mizutani Foundation for Glycoscience. E. de O. is a fellow of the Consejería de Educación (CAM).



REFERENCES

1. Dedicated to Prof. J. Thiem on the occasion of his 60th birthday.
2. Dondoni, A. The Thiazole Aldehyde Synthesis. *Synthesis* **1998**, 1681–1706.
3. Dondoni, A.; Merino, P. Diastereoselective Homologation of D-(*R*)-Glyceraldehyde Acetonide Using 2-(Trimethylsilyl)thiazole: 2-*O*-Benzyl-3,4-*O*-isopropylidene-D-erythrose. *Org. Synth.* **1995**, 72, 21–31.
4. Dondoni, A.; Fantin, G.; Fogagnolo, M.; Medici, A.; Pedrini, P. Iterative, Stereoselective Homologation of Chiral Polyalkoxyaldehydes Employing 2-(Trimethylsilyl)thiazole as a Formyl Anion Equivalent. The Thiazole Route to Higher Carbohydrates. *J. Org. Chem.* **1989**, 54, 693–702.
5. Dondoni, A.; Douglas, W.; Shinkai, I. Spirodioxolane Intermediates in the Reaction of 2-(Trimethylsilyl)thiazole with Aldehydes. Support for the 2-Ylide Mechanism. *J. Org. Chem.* **1993**, 58, 3196–3200.
6. For a preliminary communication, see: Marco-Contelles, J.; de Opazo, E. Intramolecular Alkylation of ϵ -Iodo, α -alkoxy Aldehydes. *Tetrahedron Lett.* **2000**, 41, 5507–5510.
7. Marco-Contelles, J.; de Opazo, E.; Arroyo, N. Synthesis of Higher-Carbon Sugars by Addition of Organometallic Reagents to Aldehydes or Lactols Derived from Carbohydrates. *Tetrahedron* **2001**, 57, 4729–4739.
8. Maycock, C.D.; Barros, M.T.; Godinho, L.C. An Application of Quinic Acid to the Synthesis of Cyclic Homochiral Molecules: A Common Route to Some Interesting Carbocyclic Nucleoside Precursors. *Tetrahedron Lett.* **1993**, 34, 7985–7988.
9. Kobayashi, Y.; Miyazaki, H.; Shiozaki, M. Synthesis of Trehazolin, Trehalamine, and the Aminocyclitol Moiety of Trehazolin: Determination of Absolute Configuration of Trehazolin. *J. Org. Chem.* **1994**, 59, 813–822.
10. Caine, D. Alkylation of Enols and Enolates. In *Comprehensive Organic Synthesis*; Trost, B.M., Fleming, I. Eds.; Pergamon Press: London, 1991; Vol. 3, 1–63.
11. (a) Hirota, K.; Monguchi, Y.; Sajiki, H.; Sako, M.; Kitade, Y. Novel Synthesis of Purine Acyclonucleosides Possessing a Chiral 9-Hydroxyalkyl Group by Sugar Modification of 9-D-Ribitylpurines. *J. Chem. Soc., Perkin Trans. 1* **1998**, 941–946; (b) Boehme, H.; Sutoyu, P.N. 3,5-Dideoxy-3,5-di-*C*-methyl-L-idose-trimethylenedithioacetal, ein Chiraler Baustein zur Macrolidantiika-Synthese. *Liebigs Ann. Chem.* **1982**, 1643–1655; (c) Redlich, H.; Neumann, H.-J. Zur Formylierung von Isochroman, Isothiochroman und deren Derivate. *Chem. Ber.* **1981**, 114, 2029–2050; (d) Schaffer, R. Branched-chain Sugars. II. A Diethylidene-octose. *J. Am. Chem. Soc.* **1959**, 81, 2838–2842; (e) Schaffer, R.; Isbell, H. Branched-chain Sugars. I. A 9-Aldo-4-*C*-formylnonose Derivative. *J. Am. Chem. Soc.* **1959**, 81, 2178–2183.
12. (a) Fernández, R.; Martín-Zamora, E.; Pareja, C.; Vázquez, J.; Díez, E.; Monge, A.; Lassaleta, J.M. Synthesis of Enantiopure α -Alkoxy- α -Trifluoromethyl Aldehydes and Carboxylic Acids from Trifluoromethyl Ketones. *Angew. Chem. Int. Ed. Engl.* **1998**, 37, 3428–3430; (b) Kuhn, C.; Le Gouadec, G.; Skaltsounis, A.L.; Florent, J.-C. Synthesis of Chiral 4-Alkyl 4-Hydroxy Cyclopentenones via Diastereoselective Tandem Aza-Cope/Mannich Cyclization Reaction from Aldehyde Sugar. *Tetrahedron Lett.* **1995**, 36, 3137–3140.
13. Paterson, I.; Tillyer, R.D. High π -Face Selectivity in *Anti* Aldol Reactions of *E*-Enol Borinates from Chiral Alkoxyethylketones: Stereocontrolled Synthesis of C₂₄-C₃₂ Polyol Subunit of Rapamycin. *J. Org. Chem.* **1993**, 58, 4182–4184.



14. Intermediates of type **A** (Scheme 3) have been proposed by Dondoni in the mechanism of the thiazole-carbonyl coupling (see reference 5).
15. (a) Seebach, D.; Sting, A.R.; Hoffmann, M. Self-Regeneration of Stereocenters (SRS)-Applications, Limitations, and Abandonment of a Synthetic Principle. *Angew. Chem. Int. Ed. Engl.* **1996**, *53*, 2708–2748; (b) Seebach, D.; Imwinkelried, R.I.; Weber, T. EPC Syntheses with C,C Bond Formation *via* Acetals and Enamines. In *Modern Synthetic Methods*, Scheffold, R. Ed.; Springer: New York, 1986, Vol. 4, p 125–259; (c) Naef, R.; Seebach, D. Direct Diastereoselective Alkylation of Tartaric Acid Through an Enolate. *Angew. Chem. Int. Ed. Engl.* **1981**, *20*, 1030–1031. See also: (d) Ladner, W. Stereoselektive Synthese von α -Alkyl- α,β -dihydroxy-carbonsäureester-Derivativen: Synthese der (+)-Viridiflorsäure. *Chem. Ber.* **1983**, *116*, 3413–3426; (e) Fräter, G.; Müller, U.; Günther, M. Synthesis of Enantiomerically Pure Enriched Atrolactic Acid and Other α -Hydroxy Acids. *Tetrahedron Lett.* **1981**, *22*, 4221–4224.
16. Martin, S.F. Methodology for the Construction of Quaternary Carbon Centers. *Tetrahedron* **1980**, *36*, 419–460.

Received February 5, 2001

Accepted July 19, 2001



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081CAR100108278>