Gray, M. W., and Lane, B. G. (1968), Biochemistry 7, 3441.

- Hall, R. H. (1971), The Modified Nucleosides in Nucleic Acids, New York, N. Y., Columbia University Press, p 109.
- Hudson, L., Gray, M., and Lane, B. G. (1965), *Biochemistry* 4, 2009.
- Hurst, R. O., and Butler, G. C. (1951), J. Biol. Chem. 193, 91.
- Ishikura, H., Yamada, Y., Murao, K., Saneyoshi, M., and Nishimura, S. (1969), *Biochem. Biophys. Res. Commun.* 37, 990.
- Kimura-Harada, F., Harada, F., and Nishimura, S. (1972a), FEBS (Fed. Eur. Biochem. Soc.) Lett. 21, 71.
- Kimura-Harada, F., von Minden, D. L., McCloskey, J. A., and Nishimura, S. (1972b), *Biochemistry* 11, 3910.
- Kirby, K. S. (1956), Biochem. J. 64, 405.
- Koerner, J. F., and Sinsheimer, R. L. (1957), J. Biol. Chem. 228, 1049.
- Lane, B. G. (1963), Biochim. Biophys. Acta 72, 110.
- Lane, B. G. (1965), Biochemistry 4, 212.
- Lane, B. G., Diemer, J., and Blashko, C. A. (1963), Can. J. Biochem. 41, 1927.

- McLennan, B. D., and Lane, B. G. (1968), Can. J. Biochem. 46, 93.
- Miller, J. P., and Schweizer, M. P. (1972), Anal. Biochem. 50, 327.
- Nishimura, S. (1972), Progr. Nucl. Acid Res. Mol. Biol. 12, 49.
- Powers, D. M., and Peterkofsky, A. (1972), J. Biol. Chem. 247, 6394.
- Ralph, R. K., and Bellamy, A. R. (1964), *Biochim. Biophys. Acta* 87, 9.
- Rushizky, G. W., and Sober, H. A. (1962), *Biochim. Biophys.* Acta 55, 217.
- Schweizer, M. P., Chheda, G. B., Baczynskyj, L., and Hall, R. H. (1969), *Biochemistry* 8, 3283.
- Schweizer, M. P., McGrath, K., and Baczynskyj, L. (1970), Biochem. Biophys. Res. Commun. 40, 1046.
- Singer, B., and Fraenkel-Conrat, H. (1963), Biochim. Biophys. Acta 72, 534.
- Singh, H., and Lane, B. G. (1964), Can. J. Biochem. 42, 1011.
- Takemura, S., Murakami, M., and Miyazaki, M. (1969), J. Biochem. (Tokyo) 65, 553.

3'-O-Aminoacyl-2'-deoxyadenosines and 2'-O-Aminoacyl-3'-deoxyadenosines Related to Charged Transfer Ribonucleic Acid Termini[†]

Morris J. Robins,* Roger A. Jones, and Malcolm MacCoss‡

ABSTRACT: Aminoacyl nucleosides derived from 2'-deoxyadenosine and 3'-deoxyadenosine have been isolated as pure solids and completely characterized for the first time. Reaction of 5'-O-(di-p-methoxytrityl)-2'-deoxyadenosine (1) with N-trityl blocked amino acid anhydrides (2a-c) (generated *in* situ from the corresponding N-tritylamino acid and dicyclohexylcarbodiimide) in the presence of 4-N,N-dimethylaminopyridine gave the 3'-O-(N-tritylaminoacyl)-5'-O-(di-p-methoxytrityl)-2'-deoxyadenosines (3a-c) in high yields. This coupling reaction was unsuccessful using pyridine. Analogous treatment of 5'-O-(mono-p-methoxytrityl)-3'-deoxyadenosine (6) gave the corresponding 2'-O-(N-tritylaminoacyl)-5'-O-(mono-p-methoxytrityl)-3'-deoxyadenosines (7a-c). The Lleucine (a), L-phenylalanine (b), and L-methionine (c) com-

Lt is well established that the aminoacyl nucleoside antibiotic puromycin inhibits protein biosynthesis by simulating a charged terminus of tRNA. Acting in this role it accepts the growing peptide chain from peptidyl-tRNA and the covalently linked peptidylpuromycin dissociates from the ribosomal complex (for example, see Nathans (1964), Traut and Monro pounds were prepared in each series. Complete deblocking was effected using formic acid-1-butanol-toluene (1:1:1) at room temperature. Under these conditions the 3'-O-(L-aminoacyl)-2'-deoxyadenosines (4a-c) and 2'-O-(L-aminoacyl)-3'-deoxyadenosines (8a-c) were obtained in high yield with no detectable hydrolysis of either the aminoacyl ester or glycosidic bonds.

N-Formylmethionyl and *N*-acetylphenylalanyl derivatives were prepared in each series by subsequent acylation of the free aminoacyl compounds with acetic formic anhydride and *p*-nitrophenyl acetate, respectively. Biochemical rationale for the use of these compounds in the study of protein biosynthesis and initiation processes and preliminary biochemical data are discussed.

(1964), and Coutsogeorgopoulos (1967); for a recent concise review, see Suhadolnik (1970)). Various 3'(2')-O-aminoacylnucleosides and nucleotide derivatives have been synthesized and evaluated as analogous peptide receptors in protein biosynthetic systems (see, for example, Rammler and Khorana, 1963; Waller *et al.*, 1966; Žemlička *et al.*, 1969; Rychlik *et al.*, 1969; Chládek *et al.*, 1970; Černá *et al.*, 1970a,b; Chládek, 1972, and other papers of the Czech group; Gottikh *et al.*, 1970; Tarusova *et al.*, 1971; Pozdnyakov *et al.* (1972), and other papers of the Russian group), and base as well as amino acid variation has been explored especially by the Czech group.

An example of such protein biosynthesis blockage by 2',3'bis(O-aminoacyl)adenosines (Černá *et al.*, 1970a) added

⁺ From the Department of Chemistry, The University of Alberta, Edmonton, Alberta, Canada T6G 2G2. *Received August 13, 1973*. This work was generously supported by Grant No. 9530-108 from the Defense Research Board of Canada, Grant No. A5890 from the National Research Council of Canada, and The University of Alberta. This paper is Nucleic Acid Related Compounds. 10. For the previous paper in this series, see Robins and Jones (1974).

[‡] University of Alberta Postdoctoral Fellow, 1972-present.

interest to a hypothesis of Neumann et al. (1968) that protein synthesis might occur by transfer of the growing peptide to the 2'(3')-hydroxyl of a 3'(2')-O-aminoacyl charged tRNA. This step would be followed by intramolecular peptidyl transfer from the 2'(3')-hydroxyl to the amino function of the 3'(2')-O-amino acid as the actual elongation step. The significant peptide acceptor activity of 2',3'-bis(O-aminoacyl)adenosines (Černá et al., 1970a), contrary to the conclusion of Pozdnyakov et al. (1972), does not preclude the Neumann et al. (1968) hypothesis since spontaneous intramolecular amino acid acylation of the adjacent amine function to give 3'(2')-O-dipeptidyladenosine has been observed (Neumann et al., 1968) and strong acceptor activity of such a dipeptidyl derivative demonstrated (Cerná et al., 1970a). The virtually complete lack of acceptor activity of 3'-O-(L-phenylalanyl)-2'-deoxyadenosine (Rychlik et al., 1969) was compatible with the Neumann et al. (1968) proposal, but such a negative result neither demands the presence of a 2'-OH function nor provides any definitive support. A recent communication (Pozdnyakov et al., 1972) reported the high puromycin-like activity of 3'-O-phenylalanyl-2'-O-methyladenosine. This interesting result would appear to obviate the necessity of a 2'hydroxyl function for proper binding in the acceptor vicinity (Coutsogeorgopoulos, 1967) and for accepting activity assuming that no demethylation occurs (for such a nullifying example, see Fox et al. (1971)).

Although the naturally occurring aminoacyl-tRNAs are thought to occur and function as 3' esters of the terminal adenosine (see Zachau and Feldmann (1965) and references therein), very facile and rapid 2'-O \rightleftharpoons 3'-O isomerization occurs in aqueous media (McLaughlin and Ingram, 1965) and, thus, 2'-O-aminoacyl involvement has not been precluded. Indeed, a recent communication (Sprinzl et al., 1973) has reported that a modified yeast tRNA^{Phe} terminating with 3'-deoxyadenosine (cordycepin) can be enzymatically aminoacylated whereas the corresponding 2'-deoxyadenosine terminated tRNA cannot. This suggests possible initial charging at the 2'-hydroxyl group of intact tRNA. The 2'-Naminoacyl isomer of puromycin has been reported to be inactive as an inhibitor of protein synthesis (Nathans' and Neidle, 1963). This result has been quoted (Zachau and Feldmann, 1965) as evidence for 3'-O-aminoacyl functionality in biological circumstances. Tarusova et al. (1971) have synthesized an ω -aminocaproic ester of adenosine 3',5'cyclic monophosphate but there appear to have been no specifically linked 2'-O-aminoacyl nucleosides available for evaluation previously.

Certain 5'-N-aminoacyl-5'-amino-5'deoxyadenosines have been shown to inhibit protein biosynthesis (Robins *et al.*, 1971). The aromatic amino acid phenylalanine was most effective in that system as has been previously observed in the puromycin reaction (Nathans and Neidle, 1963).

The initiation of protein synthesis in *Escherichia coli* (Clark and Marcker, 1966) and probably in prokaryotic systems in general (for a review, see Lucas-Lenard and Lipmann (1971)) is thought to proceed exclusively by a specific species of *N*-formylmethionyl-tRNA. Initiation by *N*-acetyl-phenylalanyl-tRNA in an *E. coli* system was demonstrated by Lucas-Lenard and Lipmann (1967). Eukaryotic systems, in contrast, employ other aminoacyl-tRNAs including a free methionine species. Thus, a biological species specificity at the initiation stage of protein biosynthesis apparently exists which could be of great potential significance in terms of selective inhibition.

In the present study, specific 3'-O-aminoacyl nucleosides

containing 2'-deoxyadenosine and 2'-O-aminoacyl nucleosides containing 3'-deoxyadenosine (cordycepin) have been prepared and fully characterized as analytically pure solids. The amino acids L-leucine, L-phenylalanine, and L-methionine were employed as examples of aliphatic, aromatic, and control methionine types, respectively. In addition, the *N*formylmethionyl and *N*-acetylphenylalanyl derivatives were synthesized in order to evaluate possible effects on the complex initiation process.

Experimental Section

Melting points were determined on Reichert microstage or Gallenkamp apparatus and are uncorrected. Ultraviolet (uv) spectra were determined on Cary 14 or 15 spectrophotometers. Nuclear magnetic resonance (nmr) spectra were recorded on a Varian HA-100 instrument. Optical rotations were measured using a Perkin-Elmer Model 141 polarimeter with a 10-cm 1-ml micro cell. Elemental analyses were obtained by the microanalytical laboratory of this department and Schwarzkopf Microanalytical Laboratory. Thin-layer chromatography (tic) was effected using Eastman Kodak sheets (silica gel No. 13181) in solvent systems indicated; developed chromatograms were detected by 2537-A light (chromophore), 20% perchloric acid spray (trityl, mono-, and di-*p*-methoxytrityl functions show yellow to red, respectively), and ninhydrin spray (slow development of yellow color upon warming, with free aminoacyl nucleosides). Column chromatography was performed on alumina (J. T. Baker No. 0537 aluminum oxide) or silica gel (Gebr. Herrmann, Kieselgel). Evaporations were effected using a Büchler rotating evaporator with a Dry Ice cooled dewar condensor under aspirator or mechanical oil pump vacuum at 30° or lower.

The preparations of 3'-deoxyadenosine (cordycepin) (Robins *et al.*, 1973), 5'-O-(di-*p*-methoxytrityl)-2'-deoxyadenosine (1) (Hogenkamp and Oikawa, 1964), and N-trityl-L-leucine, N-trityl-L-phenylalanine, and N-trityl-L-methionine (Zervas and Theodoropoulos, 1956; Stelakatos *et al.*, 1959) followed outlined procedures. Ethyl acetate was dried by distillation from "P₂O₆" and stored over Linde molecular sieves (4A, dried at 200°). Pyridine was dried by refluxing over and then distillation from CaH₂.

5'-O-(Mono-p-methoxytrityl)-3'-deoxyadenosine (6). To a solution of 2.5 g (0.01 mol) of 3'-deoxyadenosine in 75 ml of dry pyridine was added 4.6 g (0.015 mol) of mono-p-methoxytrityl chloride and the solution was stirred for 39 hr at 13°. MeOH (150 ml) was added and stirring was continued for 1.5 hr after which the solution was poured into 400 ml of ice-water. This mixture was extracted with CHCl3 and the combined extracts were evaporated to give 7.9 g of a brown gum. This material was dissolved in 10 ml of CHCl₃ and applied to a column (2×38 cm, 250 g, packed in CHCl₃) of alumina. The column was then washed with 2350 ml of CHCl₃ and the fast migrating materials (tlc, CHCl₃-MeOH 97:3) eluted were discarded. Solvent was changed to CHCl3-MeOH (95:5) and the first 275 ml of eluate containing primarily the same fast migrating (tlc) impurities was discarded. The following 725 ml of eluate was combined and evaporated. The residue was dissolved in a minimum volume of CHCl₃ and dripped slowly into pentane with vigorous stirring. The precipitate was filtered and air-dried to give 3.0 g (57%) of 6. A sample for analysis was recrystallized from MeOH to give 6: mp 110-114°; uv (MeOH) max 259 and 233 nm (e 16,700, 18,400), min 245 and 225 nm (e 13,700, 16,300); nmr (CDCl₃, Me₄Si internal) δ 2.20 (br m, 2, H_{3',2''}), 3.34 (m, 2, H_{5',5''}),

					E	ementa	ıl Anal.				
Wald				Calcd	(%)			Found	d (%)		
(%)	Mp ^b (°C)	Formula	C	Н	N	S	С	н	N	S	Uv Spectra nm $(\epsilon)^{c}$
62	119–120	$C_{56}H_{56}N_6O_6$	73.98	6.21	9.25		73.65	6.31	9.65		max 234 (32,600) sh 260 (17,300) min 225 (30, 300)
81	120–125	$C_{59}H_{54}N_6O_6$	75.14	5.77	8.91		74.90	5.65	8.79		max 233 (30, 300) sh 260 (16, 400) min 227 (29, 600)
90	120 softens 110-115	$C_{55}H_{54}N_6O_6S$	71.25	5.87	9 .07	3.46	71.16	5.79	9.01	3.45	max 233 (31,700) sh 260 (16,700) min 225 (30,000)
92	113–116	$C_{55}H_{54}N_6O_5$	75.14	6.19	9.56		74.68	6.36	9.78		max 230 (28,200) max 258 (17,000) min 226 (27,600) min 247 (15,700)
88	121–123	$C_{56}H_{52}N_6O_5$	76.29	5.74	9.21		76.59	5.89	9.15		$\begin{array}{c} \text{min} 247 (15,700) \\ \text{max} 228 (30,300) \\ \text{max} 259 (17,300) \\ \text{min} 247 (16,000) \end{array}$
84	110–114	$C_{54}H_{52}N_6O_5S$	72.30	5.84	9.37	3.58	72.38	6.22	9.10	3.61	max 231 (30,100) max 259 (17,900) min 226 (29,400) min 249 (16,500)
	Yield (%) 62 81 90 92 88 88 84	Yield (%) Mp ^b (°C) 62 119–120 81 120–125 90 120 softens 110–115 92 113–116 88 121–123 84 110–114	Yield ($\%$)Mp ^b (°C)Formula62119–120C_{56}H_{56}N_6O_681120–125C_{59}H_{54}N_6O_690120 softens 110–115C_{55}H_{54}N_6O_6S92113–116C_{55}H_{54}N_6O_588121–123C_{56}H_{62}N_6O_584110–114C_{54}H_{52}N_6O_5S	Yield ($?_{5}$) Mp ^b (°C) Formula C 62 119–120 C ₅₆ H ₅₆ N ₆ O ₆ 73.98 81 120–125 C ₅₉ H ₅₄ N ₆ O ₆ 75.14 90 120 C ₅₅ H ₅₄ N ₆ O ₆ S 71.25 softens 110–115 2 113–116 C ₅₅ H ₅₄ N ₆ O ₅ 88 121–123 C ₅₆ H ₅₂ N ₆ O ₅ 76.29 84 110–114 C ₅₄ H ₅₂ N ₆ O ₅ S 72.30	Yield Mp ^b (°C) Formula Calcd 62 119–120 C $_{56}H_{56}N_6O_6$ 73.98 6.21 81 120–125 C $_{59}H_{54}N_6O_6$ 75.14 5.77 90 120 C $_{55}H_{54}N_6O_6S$ 71.25 5.87 90 120 C $_{56}H_{54}N_6O_6S$ 71.25 5.87 92 113–116 C $_{56}H_{54}N_6O_5$ 75.14 6.19 88 121–123 C $_{56}H_{62}N_6O_5$ 76.29 5.74 84 110–114 C $_{54}H_{52}N_6O_5S$ 72.30 5.84	Yield (%)Mp b (°C)FormulaCHN62119–120C $_{56}H_{56}N_6O_6$ 73.986.219.2581120–125C $_{59}H_{54}N_6O_6$ 75.145.778.9190120 softens 110–115C $_{55}H_{54}N_6O_6$ 71.255.879.0792113–116C $_{55}H_{54}N_6O_5$ 75.146.199.5688121–123C $_{56}H_{52}N_6O_5$ 76.295.749.2184110–114C $_{54}H_{52}N_6O_5$ 72.305.849.37	Yield (%)Mp b (°C)FormulaCHNS62119–120C $_{56}H_{56}N_6O_6$ 73.986.219.2581120–125C $_{59}H_{54}N_6O_6$ 75.145.778.9190120 softens 110–115C $_{55}H_{54}N_6O_6S$ 71.255.879.073.4692113–116C $_{55}H_{54}N_6O_5$ 75.146.199.5688121–123C $_{56}H_{62}N_6O_5$ 76.295.749.2184110–114C $_{54}H_{52}N_6O_5S$ 72.305.849.373.58	Yield (%)Mp b (°C)FormulaCHNSC62119–120C $_{56}H_{56}N_6O_6$ 73.986.219.2573.6581120–125C $_{59}H_{54}N_6O_6$ 75.145.778.9174.9090120 softens 110–115C $_{55}H_{54}N_6O_6S$ 71.255.879.073.4671.1692113–116C $_{55}H_{54}N_6O_5$ 75.146.199.5674.6888121–123C $_{56}H_{52}N_6O_5$ 76.295.749.2176.5984110–114C $_{54}H_{52}N_6O_5S$ 72.305.849.373.5872.38	Yield (%)Mp ^b (°C)FormulaCHNSCH62119–120C $_{86}H_{86}N_6O_6$ 73.986.219.2573.656.3181120–125C $_{59}H_{54}N_6O_6$ 75.145.778.9174.905.6590120 softens 110–115C $_{55}H_{54}N_6O_6$ 71.255.879.073.4671.165.7992113–116C $_{55}H_{54}N_6O_5$ 75.146.199.5674.686.3688121–123C $_{56}H_{62}N_6O_5$ 76.295.749.2176.595.8984110–114C $_{64}H_{62}N_6O_5$ 72.305.849.373.5872.386.22	Yield (%) Mp ^b (°C) Formula C H N S C H N 62 119–120 $C_{56}H_{56}N_6O_6$ 73.98 6.21 9.25 73.65 6.31 9.65 81 120–125 $C_{56}H_{54}N_6O_6$ 75.14 5.77 8.91 74.90 5.65 8.79 90 120 $C_{35}H_{34}N_6O_6S$ 71.25 5.87 9.07 3.46 71.16 5.79 9.01 90 120 $C_{35}H_{34}N_6O_6S$ 75.14 6.19 9.56 74.68 6.36 9.78 92 113–116 $C_{36}H_{62}N_6O_5$ 76.29 5.74 9.21 76.59 5.89 9.15 88 121–123 $C_{56}H_{62}N_6O_5$ 72.30 5.84 9.37 3.58 72.38 6.22 9.10	Yield (%) Mp ^b (°C) Formula C H N S C H N S 62 119–120 $C_{56}H_{56}N_6O_6$ 73.98 6.21 9.25 73.65 6.31 9.65 81 120–125 $C_{50}H_{54}N_6O_6$ 75.14 5.77 8.91 74.90 5.65 8.79 90 120 $C_{56}H_{54}N_6O_6$ 71.25 5.87 9.07 3.46 71.16 5.79 9.01 3.45 92 113–116 $C_{56}H_{54}N_6O_5$ 75.14 6.19 9.56 74.68 6.36 9.78 88 121–123 $C_{56}H_{62}N_6O_5$ 76.29 5.74 9.21 76.59 5.89 9.15 84 110–114 $C_{54}H_{62}N_6O_5$ 72.30 5.84 9.37 3.58 72.38 6.22 9.10 3.61

TABLE I: Fully Blocked Aminoacyl Derivatives.^a

^{*a*} All analytical samples were recrystallized from methanol and dried for 15 hr at room temperature over phosphorus pentoxide and paraffin wax at 0.1 mm. ^{*b*} See footnote 3 in text. ^{*c*} Determined in methanol.

3.75 (s, 3, OCH₃), 4.74 (m, 2, H_{2'}, H_{4'}), 5.94 (d, $J_{1'-2'} = 3.0$ Hz, 1, H₁'), 6.21 (br, 2, 6-NH₂), 6.7–7.5 (m, 14, aromatic), 8.09 (s, 1, H₂), 8.30 (s, 1, H₈).

Anal. Calcd for $C_{30}H_{29}N_5O_4$: C, 68.52; H, 5.58; N, 13.38. Found: C, 68.62; H, 5.64; N, 13.45.

Preparation of 3'-O-(N-Tritylaminoacyl)-5'-O-(di-p-methoxytrityl)-2'-deoxyadenosines (3a-c) and 2'-O-(N-Tritylaminoacyl)-5'-O-(mono-p-methoxytrityl)-3'-deoxyadenosines (7a-c). These blocked derivatives were prepared from the appropriate N-tritylamino acid anhydride (2a-c) (generated *in situ* from the corresponding N-tritylamino acid by the general procedure of Rammler and Khorana (1963)) and 5'-O-(di-p-methoxytrityl)-2'-deoxyadenosine (1) (for 3a-c) or 5'-O-(mono-p-methoxytrityl)-3'-deoxyadenosine (6) (for 7a-c) (see Tables I and II for data) by the same general procedure as illustrated in the following.

3'-O-(N-Trityl-L-methionyl)-5'-O-(di-p-methoxytrityl)-2'-deoxyadenosine (3c). To a solution of 2.42 g (0.0062 mol) of N-trityl-L-methionine in 15.5 ml of dry EtOAc was added 1.28 g (0.0062 mol) of N,N'-dicyclohexylcarbodiimide (DCC). The precipitation of N,N'-dicyclohexylurea (DCU) began immediately and the mixture was stirred for 1 hr at room temperature. The mixture was filtered directly into a solution of 1.44 g (0.0026 mol) of dried 1 in 15.5 ml of dry EtOAc. The DCU filter cake (0.6 g) was washed with an additional 15 ml of dry EtOAc (filtrate collected into reaction solution). A 0.1-g (0.0008 mol) portion of 4-N,N-dimethylaminopyridine was added and the reaction was stirred for 21 hr at 15°.¹ Tlc (CHCl₃-Me₂CO, 80:20) showed essentially complete conversion of **1** to **3c**. An additional 0.29 g of DCU was removed by filtration and this filter cake was washed with 50 ml of EtOAc. The combined filtrate was washed with 2×100 ml of ice-cold saturated aqueous NaHCO₃ and 2×100 ml of H₂O and evaporated. The resulting solid foam was dissolved in 10 ml of CHCl₃ and precipitated into 350 ml of pentane with vigorous stirring. The white solid (3.19 g) was filtered, air-dried, dissolved in 12 ml of CHCl₃, and applied to a column (2.5 \times 33 cm, 154 g, packed in CHCl₃) of alumina. The column was developed with CHCl₃ and fractions were evaluated by tlc (CHCl₃-Me₂CO, 80:20). The first 250 ml of eluate contained fast migrating material and was discarded.

TABLE II: Nmr Data on Fully Blocked Aminoacyl Derivatives.^a

					÷	
Com-						Amino
pound	$6-NH_2$	$H_{1'}$	$H_{2'}$	$\mathbf{H}_{\mathbf{3'}}$	OCH_3	Acid
3a	6.05	6.21 ^b		4.82	3.76	0.91°
3b	5.87	5.88°		4.54	3.68	3.06^{d}
3c	5.94	6.20°		4.85	3.77	2.09^{e}
7a	5.89	5.63 ^f	5.08		3.77	0.92°
7b	5.95	5.47 ¹	4.87		3.72	3.12^{d}
7c	5.98	5.59 ^{,r}	5.08		3.74	2.08^{e}

^{*a*} Chemical shifts in δ from Me₄Si internal in CDCl₃. ^{*b*} ABX "quartet", $J_{1'-2',2''} \sim 6$ and 8.5 Hz. ^{*c*} Poorly resolved doublet, CH(CH₃)₂. ^{*d*} Poorly resolved doublet, CH₂Ph. ^{*e*} Singlet, SCH₃. ^{*f*} Doublet, $J_{1'-2'} \sim 1.5$ Hz.

¹ Several preliminary reactions were investigated using varying amounts of pyridine, without success. For example, this reaction in the presence of 2.5 ml of pyridine (but no 4-N,N-dimethylaminopyridine) failed to proceed observably (tlc) in 30 hr at room temperature.

TABLE III: Aminoacyl Deoxynucleosides.^a

					Elemental Anal.								
Com-	Yield				Calc	d (%)			Foun	d (%)			
pound	(%)	Mp [∂] (°C) Formula	С	Н	N	S	С	Н	N	S	Uv Spectra nm $(\epsilon)^{c}$	$[\alpha]_{\mathrm{D}}{}^{d}$
4a	90	130	$C_{16}H_{24}N_6O_4$	52.73	6.64	23.06		52.49	6.61	23.06		max 260 (14,600) min 227 (2600)	$-26.5^{\circ f}$ (c 0.97)
4b ^{<i>e</i>}	95	Slowly softens above 75	$\begin{array}{c} C_{19}H_{22}N_6O_4 \\ 0.5HCO_2H \end{array}$	55.57	5.50	19.94		55.27	5.53	19.92		max 260 (14,900) min 227 (2400)	$-7.58^{\circ f}$ (c 1.06)
4c	86	136-140	$C_{15}H_{22}N_6O_4S$	47.10	5.80	21.98	8.38	47.40	5.92	21.74	8.12	max 259 (13, 300) min 227 (2200)	$-32.8^{\circ f}$ (c 0.97)
8a	67	169–170	$C_{16}H_{24}N_6O_4$	52.73	6.64	23.06		52.83	6.79	22.97		max 258 (14,600) min 227 (2100)	$-58.2^{\circ 9}$ (c 0.73)
8b	88	110-113	$C_{19}H_{22}N_6O_4$	57.27	5.57	21.10		56.96	5.86	20.99		max 259 (14,400) min 227 (2900)	$-28.8^{\circ g}$ (c 1.03)
8c	86	132–137	$C_{15}H_{22}N_6O_4S$	47.10	5.80	21.98	8.38	47.01	5.82	21.70	8.55	max 259 (14,400) min 227 (2800)	$(c 1.01)^{\circ}$ (c 1.01)

^{*a*} All analytical samples were recrystallized from dry ethyl acetate and dried for 15 hr at 56° over potassium hydroxide pellets and paraffin wax at 0.1 mm. ^{*b*} See footnote 3 in text. ^{*c*} Determined in methanol. ^{*d*} Determined in dimethylformamide freshly distilled from phosphorus pentoxide at reduced pressure. ^{*e*} Dried at room temperature, 0.5 mole ratio of HCO₂H verified by nmr. ^{*f*} Determined at 20°. ^{*g*} Determined at 22°.

Com-					
pound	$6-\mathbf{NH}_2$	$H_{1'}$	$\mathbf{H}_{2'}$	$H_{3'}$	Amino Acid
4a	7.31	6.380		5.40	0.87 and 0.94°
4b	7.28	6.26 ^b		5.30	2.90 ^d
4c	7.30	6.38°		5.37	2.06 ^e
8a	7.25	6.08 ⁷	5.63		0.85 and 0.92°
8b	7.29	6.06 ^f	5.55		2.87 ^{<i>d</i>}
8c	7.27	6.09 ^f	5.61		2.01 ^e

^{*a*} Chemical shifts in δ from Me₄Si internal in Me₂SO-*d*₆. ^{*b*} ABX "quartet", $J_{1'-2',2''} \sim 6.3$ and 8.7 Hz. ^{*c*} Two doublets, $J \sim 3.0$ Hz, CH(CH₃)₂. ^{*d*} Doublet, $J \sim 7$ Hz,-CH₂Ph. ^{*e*} Singlet, SCH₃. ^{*f*} Doublet, $J_{1'-2'} \sim 2$ Hz.

The succeeding fractions containing only product were combined and evaporated. The resulting white solid foam was precipitated from 10 ml of CHCl₃ into 350 ml of pentane and the resulting white solid (**3c**, 2.17 g, 90%) was dried at room temperature over "P₂O₃"-paraffin wax. (See Tables I and II for data.)

Deblocking to Give the 3'-O-Aminoacyl-2'-deoxyadenosines (4a-c) and 2'-O-Aminoacyl-3'-deoxyadenosines (8a-c). These compounds were prepared from the blocked derivatives 3a-c and 7a-c, respectively, by the same general procedure as illustrated here for the following.

3'-O-(L-Methionyl)-2'-deoxyadenosine (4c). To 30 ml of a solution of 98% formic acid-1-butanol-toluene (1:1:1 by volume) was added 0.42 g (0.00045 mol) of 3c and the resulting orange-red solution was allowed to stand at room

temperature for 3 min.² An additional 100 ml of 1-butanol was added and the now colorless solution was evaporated to approximately 2 ml (vacuum pump, $<20^{\circ}$). Two successive coevaporations with 100-ml portions of 1-butanol-toluene (1:1) gave a gum which was stirred magnetically with four successive 50-ml portions of pentane. The supernatant solutions were decanted and the final portion was filtered to give 0.16 g (86%) of **4c** (as the hemiformate). A sample for analysis was recrystallized³ from dry EtOAc and dried at 56° (0.01 mm) over KOH pellets and paraffin wax. (See Tables III and IV for data.)

3'-O-(N-Acetyl-L-phenylalanyl)-2'-deoxyadenosine (5b). To a solution of 0.72 g (0.004 mol) of p-nitrophenyl acetate in 10 ml of absolute EtOH was added 0.21 g (0.0005 mol) of 4b hemiformate. The solution was stirred for 16 hr at room temperature at which time tlc (p-dioxane-acetonitrile, 9:1) indicated the reaction was complete and 25 ml of H₂O was added. This mixture was extracted with 4×25 ml of CHCl₃ and the combined organic phase was evaporated. The residue was dissolved in 4 ml of Me₂CO-CHCl₃ (20:80) and applied to a column (2.5 \times 23 cm, 35 g, packed in the same solvent mixture) of silica gel. Elution of fast migrating materials (tlc, pdioxane-acetonitrile, 9:1) was effected with 250 ml of Me₂CO-CHCl₃ (20:80) and the solvent was changed to absolute EtOH. The first 50 ml was uv transparent and the following 225 ml contained only 5b. These combined fractions were evaporated to give a white solid foam which was precipitated from acetone into pentane (or crystallized³ from Et₂O-CHCl₃) to give 0.17 g (76%) of **5b**. A sample for analysis was recrystallized³ from dry EtOAc-absolute EtOH-pentane and dried at 56° (0.01 mm) over KOH pellets and paraffin wax to

^a Reaction time was 20 min for the 5'-O-(mono-p-methoxytrityl) compounds 7a-c.

⁶ These products were subjected to the usual process of recrystallization (*i.e.*, dissolved by warming in solvent and found to separate from solution upon cooling). However, most of the samples did not exhibit diffracting properties under a polarizing microscope and are probably amorphous or microcrystalline solids.

give **5b**: mp 214–215°; $[\alpha]_D^{20} - 18.1^\circ$ (*c* 0.99, DMF); uv (MeOH) max 260 nm (ϵ 17,100) min 227 nm (ϵ 3200); nmr (Me₂SO-*d*₆, TMS internal) δ 1.85 (s, 3, COCH₃), 2.22–2.62 (m, 1, H_{2'}), 2.74–3.14 (m, 1, H_{2''}), 3.01 (d, *J* = 7.5 Hz, 2, CH₂Ph), 3.57 (m, 2, H_{5',5''}), 3.85 (m, 1, H_{4'}), 4.48 (d of t, *J* = 7.5 and 7.5 Hz, 1, -CH (NHAc)CH₂Ph), 5.24–5.54 (m, 2, 5'-OH, H_{3'}), 6.27 ("q," *J*_{1'-2',2''} = 8.4 and 5.6 Hz, 1, H_{1'}), 7.30 (m, 7, 6-NH₂, C₆H₃), 8.16 (s, 1, H₂), 8.34 (s, 1, H₈), 8.40 (d, *J* = 7.5 Hz, NHAc).

Anal. Calcd for $C_{21}H_{24}N_6O_5$: C, 57.26; H, 5.49; N, 19.08. Found: C, 57.16; H, 5.44; N, 18.97.

3'-O-(N-Formyl-L-methionyl)-2'-deoxyadenosine (5c). To a solution of 0.20 g (0.0005 mol) of 4c hemiformate in 20 ml of MeOH at 0° was added approximately 0.04 ml (0.0006 mol) of acetic formic anhydride (Krimen, 1970) and the solution was stirred for 10 min at 0° . Tlc (*p*-dioxane-acetonitrile, 9:1) indicated that reaction was complete. Addition of 30 ml of 1butanol-toluene (1:1) was followed by evaporation of the resulting solution (vacuum pump, $<20^{\circ}$). The residue was coevaporated with 15 ml of absolute EtOH and then 15 ml of CH₂Cl₂ to give a white powder. This material was recrystallized³ from CH₂Cl₂ (with a second crop obtained by adding pentane) to yield 0.15 g (73%) of 5c (dried at room temperature (0.01 mm) over KOH pellets and paraffin wax for 15 hr): mp 165–167°; $[\alpha]_D^{20}$ –43.8° (c 1.07, DMF); uv (MeOH) max 260 nm (e 17,100) min 227 nm (e 4500); nmr (Me₂SO-d₆, TMS internal) & 1.74-2.14 (m, 2, CH₂CH₂S), 2.05 (s, 3, SCH_3), 2.36–2.64 (m, 3, $H_{2'}$, CH_2S), 2.78–3.10 (m, 1, $H_{2''}$), 3.62 (m, 2, H_{5',5''}), 4.08 (m, 1, H_{4'}), 4.47 ("t," 1, CH(NH-COH)), 5.42 (m, 2, 5'-OH, $H_{3'}$), 6.36 ("q," $J_{1'-2',2''} = 8$ and 6 Hz, 1, H_{1'}), 7.31 (s, 2, 6-NH₂), 8.14 (s, 1, -NHCOH), 8.16 $(s, 1, H_2), 8.38 (s, 1, H_8), 8.62 (d, J = 8 Hz, 1, H_8)$ NHCOH).

Anal. Calcd for $C_{16}H_{22}N_6O_5S$: C, 46.82; H, 5.40; N, 20.48; S, 7.81. Found; C, 46.54; H, 5.49; N, 20.52; S, 7.53.

2'-O-(N-Acetyl-L-phenylalanyl)-3'-deoxyadenosine (9b). Acetylation of **8b** was effected by the procedure described above for the conversion of $4b \rightarrow 5b$. The product 9b was obtained in 80% yield and had mp 176–178°: $[\alpha]_D^{22} - 56.7^\circ$ (*c* 1.05, DMF); uv (MeOH) max 258 nm (ϵ 14,600) min 227 nm (ϵ 2500); nmr (Me₂SO-*d*₆, TMS internal) δ 1.80 (m, 1, H_{3'}), 1.82 (s, 3, COCH₃), 2.45 (m, 1, H_{3''}), 2.98 (d, J = 8 Hz, CH₂Ph), 3.47 and 3.56 (d and d, J = 3 and 3 Hz, 1 and 1, H_{5'} and H_{5''}, observed upon D₂O exchange), 4.11 (m, 1, H_{4'}) 4.50 (t, J = 8 Hz, 1, CHNHAc, observed upon D₂O exchange), 5.08 (m, 1, 5'-OH), 5.58 (m, 1, H_{2'}), 6.05 (d, J_{1'-2'} = 2.5 Hz, 1, H_{1'}), 7.25 (br s, 7, 6-NH_{2'} – C₆H₅), 8.17 (s, 1, H₂), 8.33 (s, 1, H₈), 8.41 (d, J = 8 Hz, 1, CHNHAc).

Anal. Calcd for $C_{21}H_{24}N_6O_5$: C, 57.26; H, 5.49; N, 19.08. Found: C, 57.12; H, 5.51; N, 19.11.

2'-O-(*N*-Formyl-L-methionyl)-3'-deoxyadenosine (9c). Formylation of 8c was effected with acetic formic anhydride in MeOH as described above for the conversion of $4c \rightarrow 5c$. The product 9c was obtained in 85% yield and had mp 189– 191°: $[\alpha]_{D}^{22} - 82.1^{\circ}$ (*c* 0.99, DMF); uv (MeOH) max 259 nm (ϵ 13,700) min 227 nm (ϵ 2400) nmr (Me₂SO-*d*₆, TMS internal) δ 1.84–2.24 (m, 3, CH₂CH₂S, H_{3'}), 2.05 (s, 3, SCH₃), 2.4–2.6 (m, 3, CH₂S, H_{3''}), 3.60 (m, 2, H_{5',5''}), 4.20–4.60 (m, 2, CHNHCOH, H_{4'}), 5.10 (t, *J* = 6 Hz, 1, 5'-OH), 5.67 (m, 1, H_{2'}), 6.09 (d, J_{1'-2'} = 2.0 Hz, 1, H_{1'}), 7.25 (s, 2, 6-NH₂), 8.09 (s, 1, NHCOH), 8.14 (s, 1, H₂), 8.32 (s, 1, H₈), 8.55 (d, *J* = 8 Hz, 1, CHNHCOH).

Anal. Calcd for $C_{16}H_{22}N_6O_5S$: C, 46.82; H, 5.40; N, 20.48; S, 7.81. Found: C, 47.02; H, 5.61; N, 20.48; S, 7.86.

Results

The fully protected aminoacyl 2'-deoxynucleosides 3a-c(see Reaction Scheme) were prepared by treatment of 5'-O-(di-*p*-methoxytrityl)-2'-deoxyadenosine (5'-O-di-*p*-anisylphenylmethyl-2'-deoxyadenosine) (1) (Hogenkamp and Oi-



kawa, 1964) with N-tritylamino acid anhydrides 2a-c in the presence of 4-N,N-dimethylaminopyridine. The anhydrides were generated in situ from the corresponding N-tritylamino acids (Zervas and Theodoropoulos, 1956; Stelakatos et al., 1959) and DCC by the general procedure of Rammler and Khorana (1963). Although the coupling reactions did not proceed under the normal conditions in pyridine solution (Rammler and Khorana, 1963), the use of 4-N,N-dimethylaminopyridine obviated this difficulty and 3'-O-(N-trityl-L-leucyl)-5'-O-(di-p-methoxytrityl)-2'-deoxyadenosine (3a), 3'-O-(N-trityl-L-phenylanyl)-5'-O-(di-p-methoxytrityl)-2'-deoxyadenosine (3b), and 3'-O-(N-trityl-L-methionyl)-5'-O-(di-p-methoxytrityl)-2'-deoxyadenosine (3c) were obtained in high yields (see Table I). Treatment of 3'-deoxyadenosine (cordycepin) (Robins et al., 1973) with monomethoxytrityl chloride in pyridine gave 5'-O-(mono-p-methoxytrityl)-3'deoxyadenosine (6) in 57% yield. Analogous condensation reactions of 6 with 2a-c gave high yields (see Table I) of 2'-O-(N-trityl-L-leucyl)-5'-O-(mono-p-methoxytrityl)-3'-deoxyadenosine (7a), 2'-O-(N-trityl-L-phenylalanyl)-5'-O-(mono*p*-methoxytrityl)-3'-deoxyadenosine (7b), and 2'-O-(N-1)trityl-L-methionyl)-5'-O-(mono-p-methoxytrityl)-3'-deoxyadenosine (7c). Deblocking of 3a-c and 7a-c proceeded smoothly in 98% formic acid-1-butanol-toluene (1:1:1) to give 3'-O-(and 2'-O-)L-leucyl-2'-(and 3'-)deoxyadenosines (4a and 8a), 3'-O-(and 2'-O-)L-phenylalanyl-2'-(and 3'-)deoxyadenosines (4b and 8b), and 3'-O-(and 2'-O-)L-methionyl-2'-(and 3'-)deoxyadenosines (4c and 8c), respectively, in good yields (see Table III).

Treatment of **4b** and **8b** with *p*-nitrophenyl acetate in alcohol effected N acetylation of the amino acid moiety to give 3'-O-(and 2'-O-)N-acetyl-L-phenylalanyl-2'-(and 3'-)-deoxyadenosines (**5b** and **9b**) in 76 and 80% yields, respectively. Acetic formic anhydride in methanol smoothly N formylated the amino acid portion of **4c** and **8c** to give 3'-O-(and 2'-O-)N-formyl-L-methionyl-2'-(and 3'-)deoxyadenosines (**5c** and **9c**) in 73 and 85% yields, respectively.

Discussion

The wide scope of interest in the processes of protein biosynthesis in general and the puromycin reaction in particular (for example, see recent reviews by Lucas-Lenard and Lipmann (1971), Suhadolnik (1970), and the entire *Cold Spring Harbor Symp. Quant. Biol.*, 34 (1969)) has stimulated the preparation and study of many aminoacyl derivatives of various nucleosides, nucleotides, and oligonucleotides as noted in the introduction. Most of these studies have used the general procedure of Rammler and Khorana (1963) to produce 3'(2') ester mixtures which were characterized by qualitative or semiquantitative spectral and chromatographic evaluation.

The ribonucleoside products are known to be quite unstable in aqueous solution with typical half-lives of 5–30 min at near neutrality and about 30° (Rammler and Khorana, 1963; Waller *et al.*, 1966; see discussion and references in Zachau and Feldmann (1965)). This instability has been attributed to hydrogen-bonding stabilization of the hydrolytic transition state by the neighboring (2' or 3') *cis*-hydroxyl group (Bruice and Fife, 1962; Zachau and Karau, 1960; Griffin and Reese, 1964). Removal of this vicinal hydroxyl group would be expected to confer added stability (Griffin and Reese, 1964) and make the resulting aminoacyl nucleosides more amenable to biochemical study in aqueous media.

Žemlička et al. (1969) reported the preparation of 3'-O-(L-phenylalanyl)-2'-deoxyadenosine (4b) using the symmetrical anhydride coupling with the carbobenzyloxy (Cbz) amino acid followed by catalytic hydrogenolysis for removal of the Cbz blocking group. Their product was characterized, however, only by qualitative chromatography and spectrophotometry and was biochemically evaluated as a "stock solution." Our initial studies employed Cbz amino acids. However, during catalytic hydrogenolysis of the Cbz group, the presence of significant quantities of free nucleoside plus amino acid were detected even before complete deblocking occurred as observed by others (Chládek et al., 1970). In this regard it is probably significant to note that a nucleoside/phenylalanine ratio of 1.3 was recorded for the previously reported preparation of 4b (Žemlička et al., 1969). Conditions reported for deblocking tert-butyloxycarbonyl (t-Boc) amino acids were expected to be too severe for the deoxynucleoside glycosidic bond and so N-tritylamino acids (Zervas and Theodoropoulos, 1956; Stelakatos et al., 1959) were chosen.

Attempted condensation of *N*-tritylamino acids with 5'-O-(di- and mono-*p*-methoxytrityl) deoxynucleosides **1** and **6** using the *in situ* generated symmetrical anhydride (Rammler and Khorana, 1963) with pyridine as catalyst failed completely presumably due to the steric effect of the trityl group. It had been reported (Steglich and Höfle, 1969) that 4-*N*,*N*dimethylaminopyridine markedly expedited the acylation of hindered alcohols. Indeed, addition of catalytic quantities of this compound to the above ethyl acetate reaction solution resulted in rapid and essentially quantitative conversion (tlc) of 1 or 6 to the desired *N*-tritylaminoacyl deoxynucleosides 3a-c or 7a-c, respectively (see Experimental Section).⁴

Although Chládek *et al.* (1970) fully characterized blocked derivatives of aminoacyl nucleosides and obtained certain deblocked products as solids, the latter were characterized only by chromatography and spectroscopy. Indeed, it appears that previous free aminoacyl nucleosides have been studied as "stock solutions" or freeze-dried powders of not rigorously characterized constitution. This arises primarily from the inherent instability of these compounds and sometimes unsatisfactory deblocking procedures. Since the present molecules were expected to have enhanced ester stability due to the absence of a vicinal hydroxyl (Zachau and Karau, 1960; Griffin and Reese, 1964), we sought a procedure which would allow high yield isolation and complete characterization of the free aminoacyl deoxynucleosides to make them available for carefully defined biochemical studies.

It was eventually found that complete deblocking of 3a-c and 7a-c was affected with no observable (tlc) cleavage of aminoacyl ester or glycosidic bonds using a solution of formic acid-1-butanol-toluene (1:1:1) at room temperature (see Experimental Section). The solution was further diluted with 1-butanol and toluene to prevent N formylation during evaporations. These solvents provide favorable azeotropic properties and *n*-butyl formate and the various *n*-butyl trityl ethers are highly lipophilic, enhancing separation from product nucleoside derivatives. The final chromatographically homogeneous products 4a-c and 8a-c were isolated in high yields and could be recrystallized.3 See Tables III and IV for yields, physical constants, and analytical data. It is interesting to note that oxidation problems were not encountered during the syntheses of the methionine compounds as previously observed by others (Chládek et al., 1970).

Since *E. coli* and other prokaryotic systems exclusively employ a specific tRNA (fMet-tRNA^{Met}_f) for protein biosynthesis initiation (Lucas-Lenard and Lipmann, 1971 and references therein) and *N*-acetylphenylalanyl-tRNA can simulate this initiation (Lucas-Lenard and Lipmann, 1967), the *N*-formylmethionyl derivatives **5c** and **9c** and *N*-acetylphenylalanyl derivatives **5b** and **9b** were prepared to evaluate biochemical consequences of these initiator models. The active ester approach of Bodánszky (1955) employing *p*nitrophenyl acetate in alcoholic medium gave smooth N acetylation of the phenylalanine amino group. Similarly, acetic formic anhydride in methanolic solution produced facile selective N formylation of the methionine amino moiety.

Compounds **4a-c**, **5b,c**, **8a-c**, and **9b,c** were evaluated in cell free protein biosynthesis systems and also against *E. coli* K-12 and *E. coli* B cell lines for antibacterial activity (M. J. Robins, C. Coutsogeorgopoulos, and A. Bloch, to be published). It is of considerable interest to note that the 3'-O-aminoacyl-2'-deoxy nucleosides **4a-c** and **5b,c** showed 50 % inhibition of the *E. coli* lines in the order of tenfold lower concentration than puromycin (*i.e.*, in the range of 10^{-6} M vs. 10^{-5} M for puromycin). The *N*-formylmethionine-containing **5c** and *N*-acetylphenylalanyl-containing **5b** showed comparable inhibitory activity to the free aminoacyl products **4c** and **4b**. However, the 2'-O-aminoacylcordycepin deriva-

⁴ The use of 4-N,N-dimethylaminopyridine dramatically facilitates other nucleoside acylations (to be published) and should be explored in peptide chemistry [*e.g.*, mixed anhydrides of bulky N-tritylamino acids failed to condense using triethylamine (Zervas and Theodoropoulos, 1956)].

tives **8a,b** had inhibitory levels comparable to puromycin whereas the methionine-containing **8c** was in the order of 100-fold less active and **9b,c** showed no inhibitory activity toward the two bacteria at levels tested. The recent finding by Sprinzl *et al.* (1973) that cordycepin-terminated tRNA^{Phe} can be charged whereas the 2'-deoxyadenosine-terminated analog cannot enhances the interest in the differential activity of the cordycepin derivatives **8a-c** and **9b,c**.

The aminoacyl products of this study do not appear to function in the "puromycin reaction," confirming the report of Rychlik *et al.* (1969) for 4b. Fox *et al.* (1966) have suggested an empirical proposal for the inhibition of protein biosynthesis by nucleoside antibiotics with two basic centers. However, it should be recognized that the present aminoacyl deoxynucleosides might function as antibacterial molecules by completely different mechanisms.

Details of biochemical studies on these products and the synthesis and evaluation of analogous 3'-O-aminoacyl-2'-O-methyladenosines and 2'-O-aminoacyl-3'-O-methyladenosines will be reported (M. J. Robins *et al.*, in preparation).

Acknowledgment

We thank Dr. Mason G. Stout for the preparation of compound $\mathbf{6}$ and Dr. A. Bloch for the rapid transmission of biochemical data.

References

- Bodánszky, M. (1955), Nature (London) 175, 685.
- Bruice, T. C., and Fife, T. H. (1962), J. Amer. Chem. Soc. 84, 1973.
- Černá, J., Chládek, S., Rychlik, I., and Žemlička, J. (1970a), Biochim. Biophys. Acta 199, 291.
- Černá, J., Rychlik, I., Žemlíčka, J., and Chládek, S. (1970b), Biochim. Biophys. Acta 204, 203.

Chládek, S. (1972), J. Org. Chem. 37, 2863.

- Chládek, S., Pulkrábek, P., Sonnenbichler, J., Žemlička, J., and Rychlik, I. (1970), *Collect. Czech. Chem. Commun.* 35, 2296.
- Clark, B. F. C., and Marcker, K. A. (1966), J. Mol. Biol. 17, 394.
- Coutsogeorgopoulos, C. (1967), Biochemistry 6, 1704.
- Fox, J. E., Sood, C. K., Buckwalter, B., and McChesney, J. D. (1971), *Plant Physiol.* 47, 275.
- Fox, J. J., Watanabe, K. A., and Bloch, A. (1966), Progr. Nucl. Acid Res. Mol. Biol. 5, 251.
- Gottikh, B. P., Krayevsky, A. A., Tarussova, N. B., Purygin, P. P., and Tsilevich, T. L. (1970), *Tetrahedron 26*, 4419.

- Griffin, B. E., and Reese, C. B. (1964), Proc. Nat. Acad. Sci. U. S. 51, 440.
- Hogenkamp, H. P. C., and Oikawa, T. G. (1964), J. Biol. Chem. 239, 1911.
- Krimen, L. I. (1970), Org. Syn. 50, 1.
- Lucas-Lenard, J., and Lipmann, F. (1967), Proc. Nat. Acad. Sci. U. S. 57, 1050.
- Lucas-Lenard, J., and Lipmann, F. (1971), Annu. Rev. Biochem. 40, 409.
- McLaughlin, C. S., and Ingram, V. M. (1965), *Biochemistry 4*, 1442, 1448.
- Nathans, D. (1964), Proc. Nat. Acad. Sci. U. S. 51, 585.
- Nathans, D., and Neidle, A. (1963), Nature (London) 197, 1076.
- Neumann, H., Shashoua, V. E., Sheehan, J. C., and Rich, A. (1968), Proc. Nat. Acad. Sci. U. S. 61, 1207.
- Pozdnyakov, V. A., Mitin, Yu. V., Kukhanova, M. K., Nikolaeva, L. V., Krayevsky, A. A., and Gottikh, B. P. (1972), FEBS (Fed. Eur. Biochem. Soc.) Lett. 24, 177.
- Rammler, D. H., and Khorana, H. G. (1963), J. Amer. Chem. Soc. 85, 1997.
- Robins, M. J., and Jones, R. A. (1974), J. Org. Chem. 39, 113.
- Robins, M. J., Mengel, R., and Jones, R. A. (1973), J. Amer. Chem. Soc. 95, 4074.
- Robins, M. J., Simon, L. N., Stout, M. G., Ivanovics, G. A., Schweizer, M. P., Rousseau, R. J., and Robins, R. K. (1971), J. Amer. Chem. Soc. 93, 1474.
- Rychlik, I., Černá, J., Chládek, S., Žemlička, J., and Haladová, Z. (1969), J. Mol. Biol. 43, 13.
- Sprinzl, M., Scheit, K.-H., Sternbach, H., von der Haar, F., and Cramer, F. (1973), *Biochem. Biophys. Res. Commun.* 51, 881.
- Steglich, W., and Höfle, G. (1969), Angew. Chem., Int. Ed. Engl. 8, 981.
- Stelakatos, G. C., Theodoropoulos, D. M., and Zervas, L. (1959), J. Amer. Chem. Soc. 81, 2884.
- Suhadolnik, R. J. (1970), Nucleoside Antibiotics, New York, N. Y., Wiley-Interscience, Chapter 1.1.
- Tarusova, N. B., Kuznetsova, L. I., Kraevskii, A. A., and Gottikh, B. P. (1971), *Izv. Akad. Nauk SSSR*, Ser. Khim., 1511.
- Traut, R. R., and Monro, R. E. (1964), J. Mol. Biol. 10, 63.
- Waller, J.-P., Erdös, T., Lemoine, F., Guttmann, S., and Sandrin, E. (1966), *Biochim. Biophys. Acta 119*, 566.
- Zachau, H. G., and Feldmann, H. (1965), Progr. Nucl. Acid Res. Mol. Biol. 4, 217.
- Zachau, H. G., and Karau, W. (1960), Chem. Ber. 93, 1830.
- Žemlička, J., Chládek, S., Haladová, Z., and Rychlik, I. (1969), Collect. Czech. Chem. Commun. 34, 3755.
- Zervas, L., and Theodoropoulos, D. M. (1956), J. Amer. Chem. Soc. 78, 1359.