


This article was downloaded by: [UQ Library]

On: 23 June 2015, At: 02:48

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcar20>

An Exceptionally Simple Chemical Synthesis of O-Glycosylated d-Glucosamine Derivatives by Heyns Rearrangement of the Corresponding O-Glycosyl Fructoses

Arnold E. Stütz^a, Gyula Dekany^a, Brigitte Eder^a, Carina Illaszewicz^a & Tanja M. Wrodnigg^a

^a Glycogroup, Institut für Organische Chemie, Technische Universität Graz, Stremayrgasse 16, A-8010, Graz, Austria

Published online: 16 Aug 2006.

To cite this article: Arnold E. Stütz, Gyula Dekany, Brigitte Eder, Carina Illaszewicz & Tanja M. Wrodnigg (2003) An Exceptionally Simple Chemical Synthesis of O-Glycosylated d-Glucosamine Derivatives by Heyns Rearrangement of the Corresponding O-Glycosyl Fructoses, Journal of Carbohydrate Chemistry, 22:5, 253-265, DOI: [10.1081/CAR-120023468](https://doi.org/10.1081/CAR-120023468)

To link to this article: <http://dx.doi.org/10.1081/CAR-120023468>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

An Exceptionally Simple Chemical Synthesis of *O*-Glycosylated D-Glucosamine Derivatives by Heyns Rearrangement of the Corresponding *O*-Glycosyl Fructoses[†]

Arnold E. Stütz, Gyula Dekany, Brigitte Eder, Carina Illaszewicz,
and Tanja M. Wrodnigg*

Glycogroup, Institut für Organische Chemie, Technische Universität Graz,
Graz, Austria

ABSTRACT

2-*N*-Acetyl-4-*O*-(β-D-galactopyranosyl)-D-glucosamine (*N*-acetyl-D-lactosamine), a very important building block of biologically relevant oligosaccharides such as sialyl Lewis^x, is easily accessible via the Heyns rearrangement of the corresponding *O*-glycosylated ketohexose, D-lactulose. This approach can also be extended to other glucosamine derivatives employing suitable *O*-glycosylated ketoses many of which are commercially available. For example, nigerosamine (3-*O*-α-D-glucopyranosyl-D-glucosamine) was prepared from turanose (3-*O*-α-D-glucopyranosyl-D-fructose). In combination with a recently introduced vinylogous amide type *N*-protecting group, [1,3-dimethyl-2, 4, 6 (1H, 3H, 5H)-trioxypyrimidine-5-ylidene] methyl (DTPM), this access is clearly superior to other routes and eminently suitable for scaling up.

[†]This paper is dedicated to Professor Dr. Kurt Heyns.

*Correspondence: Tanja M. Wrodnigg, Glycogroup, Institut für Organische Chemie, Technische Universität Graz, Stremayrgasse 16, A-8010 Graz, Austria; E-mail: wrodnigg@orgc.tu-graz.ac.at.




INTRODUCTION

The Heyns rearrangement, wherein ketoses react with suitable amines to form ketosylamines which subsequently isomerise to the corresponding 2-amino-2-deoxyaldoses, was first discovered by Fischer^[1,2] during his studies on the osazone formation of sugars and later further investigated by Heyns and Koch in the 1950s when they found that D-glucosamine **2** was formed in the reaction of D-fructose **1** with ammonia^[3] (Scheme 1).

Subsequently, this reaction was extended by Carson^[4–6] as well as the Heyns group^[7] to a wide range of different primary and secondary amines^[8] yielding the corresponding *N*-substituted glucosamine derivatives. In addition, amino acids^[9–12] and other ketoses^[13–16] have been employed in this reaction. Generally, despite many efforts,^[17] yields rarely exceeded 20% because this reaction suffers from a variety of problems such as competition between hydrolysis and rearrangement of the initial condensation product, epimer formation at position C-2 of the rearrangement product, separation problems, Amadori rearrangement as a side reaction, disubstitution when employing primary amines and chemical instability of some products.

Because of overestimation of these difficulties there has been practically no synthetic application of this rearrangement. In context with a demand for multigram quantities of *N*-acetyl-D-lactosamine **3**, which was first discovered by Freudenberg when he investigated blood substances,^[18,19] we became interested in this reaction. Compound **3** is a constituent of various glycoproteins and glycolipids and consequently plays a very important role in many biochemical processes.^[20,21] Amongst others, it is, for example, the key intermediate in most syntheses of Lewis^X^[22–25] **4** and sulfo-Le^x, which have been identified as ligands of selectins, carbohydrate recognising receptors on the surface of endothelial cells involved in cell adhesion phenomena^[26–29] (Figure 1).

Not surprisingly, synthetic approaches to **3** have been investigated for many years and it has remained an eminently interesting target for synthetic chemists. Consequently, many different approaches have been reported, either by multistep chemical syntheses,^[30–35] with the inherent need for protecting groups, or by enzymatic methods.^[36–38] In addition, solid-phase approaches were reported.^[23] The best chemical syntheses, concerning yields and scale-up potential starting from lactose and allowing access to **3** or derivatives thereof, require seven to nine steps with overall yields ranging from 10 to, at most, 30%.^[39–44] Enzymatic approaches have employed D-galactosidases from various sources^[36–38,45–51] or D-galactosyl transferases^[52–55] as components of multi-enzyme systems which require co-factor recycling. Thus far, they have been limited to relatively small scale, and require know-how which is not generally available in synthetically oriented laboratories.



Scheme 1. The Heyns rearrangement of D-Fructose.




Figure 1. NAHc-Lactosamine (3) and Lewis^X (4).

Following the Heyns rearrangement protocol, we have recently communicated a short and efficient synthesis of selected *N*-substituted D-lactosamine derivatives starting from D-lactulose.^[56] Due to limited space, this contribution could not address the general applicability to other suitable *O*-glycosylketoses, nor was it possible to include some experimental details.

Now we would like to exemplify the tremendous practicability of the Heyns rearrangement reporting two examples of wider interest, the syntheses of new *N*-protected derivatives of D-lactosamine such as **9** and D-nigerosamine **16** [3-*O*-(α -D-glucopyranosyl)-D-glucosamine], starting from the corresponding commercially available glycosylated ketoses, lactulose **5** and turanose (3-*O*- α -D-glucopyranosyl-D-fructose) **11**, respectively. The successful Heyns rearrangement of turanose is particularly interesting due to the fact that β -elimination of the substituent at C-3 is a feasible side-reaction potentially limiting the scope of this approach.

In the case of *N*-protected lactosamine, the Heyns rearrangement in combination with the *N*-protecting group [1,3-dimethyl-2, 4, 6 (1H, 3H, 5H)-trioxypyrimidine-5-ylidene]methyl (DTPM), a vinylogous amide type (23), was found to be eminently intriguing, allowing access to the product in a straightforward procedure without chromatography in an overall yield of 65 % starting from lactulose. This approach is, because of the simple preparative operations and easy handling, clearly suitable for the preparation of large amounts at low costs and in a short time.


RESULTS AND DISCUSSION

Lactulose **5** (4-*O*- β -D-galactopyranosyl-D-fructose), by reaction with benzylamine following the Heyns rearrangement pathway initially gives *N*-benzylactulosylamine **6**, which is converted under acidic conditions to the rearrangement product *N*-benzylactosamine **7**. Hydrogenolysis of **7** leads to lactosamine hydrochloride **8** which chemoselectively undergoes *N*-acetylation to yield *N*-acetylactosamine **3** (38 – 45%). This reaction sequence can be performed in a one-pot procedure. Nevertheless, a final chromatographic purification step is necessary.^[56]

We have now optimized the purification of the intermediates and, by employing a suitable *N*-protecting group, the corresponding lactosamine derivative precipitates from the reaction mixture in the final step of the synthesis. Following this route, lactulose **5**




reacted with commercial grade benzylamine to give the corresponding ketosyl amine **6**, which could be obtained together with unreacted lactulose and side products by precipitation from diethyl ether. The rearrangement was conducted in methanol in the presence of glacial acetic acid (10:1) at room temperature within two hours. The desired *N*-benzylactosamine **7** as well as unreacted lactulose **5** and side products were obtained as a crude solid by treatment of the reaction mixture with diethyl ether. Hydrogenolysis of the *N*-benzyl group was performed in water at pH 1–2 employing Pearlman's catalyst (20 %) to yield a mixture of lactosamine hydrochloride **8** and unreacted **5**. Fortunately, side products from previous steps do not survive these reaction conditions, which allows for an "indirect" partial purification of the mixture. Finally, *N*-protection takes place in methanol employing triethylamine and 1,3-dimethyl-5-[(dimethylamino)methylene] 2,4,6 (1H, 3H, 5H)-trioxypyrimidine (DTPM-reagent).^[57] This protecting group is stable during most reaction conditions commonly used in carbohydrate chemistry such as acetylation, Zemplén conditions, alkylation, hydrogenolysis, acetal formation, silylation as well as glycosylation and can be easily removed with ammonia, hydrazine or primary amines at room temperature in a few minutes. Employing this protecting group, the desired lactosamine derivative **9** precipitates from the reaction mixture, whereas unreacted **5** remains in solution. Following this method, **9** could be obtained as a white powder in an overall yield of 65% from lactulose **5**. Per-*O*-acetylation in pyridine with acetic anhydride allowed,



Scheme 2. Synthesis of lactosamine via Heyns rearrangement.

after recrystallisation from diisopropyl ether, access to the fully protected derivative **10** in almost quantitative yield, ready for further transformations. Interestingly, formation of the epimer at C-2, the *D-manno* configured 2-amino-2-deoxysugar, was not observed under the particular conditions employed (Scheme 2).

Likewise, turanose **11** (3-*O*- α -D-glucopyranosyl-D-fructose), an easily available and cheap constituent of melezitose, a highly crystalline non-reducing trisaccharide found in various types of honey,^[58,59] could also be shown to serve as a substrate for the Heyns rearrangement, despite the fact that the glucosyl residue at position *O*-3 can give rise to a highly undesired β -elimination during the reaction. Nevertheless, it was possible to obtain the Heyns rearrangement product, nigerosamine **14**. Yields are somewhat lower than in the case of the lactulose-to-lactosamine conversion. In this synthesis, turanose **11** was stirred with benzylamine for 3 days until TLC showed the corresponding ketosylamine **12** as the main product. The reaction mixture was diluted with methanol and stirred into ether to give a precipitate containing a mixture of **12**, unreacted starting material as well as side products. For the rearrangement reaction, this mixture was



Scheme 3. Synthesis of nigerosamine via Heyns rearrangement.



dissolved in methanol/glacial acetic acid, stirred at room temperature for one to two hours and added to an excess of diethyl ether, to give a precipitate of *N*-benzyl-nigerosamine **12**, turanose and side products. Hydrogenolysis afforded the free amine **14** as the hydrochloride together with the starting material. Its treatment with the DTPM-reagent gave the *N*-protected nigerosamine **15**. Unfortunately, due to the presence of pyranoid and furanoid tautomers at the reducing end as well as side products, this product failed to precipitate. Consequently, per-*O*-acetylation followed by conventional column chromatography was necessary to yield pure nigerosamine derivative **16** (Scheme 3).

CONCLUSION

In summary, starting from commercially available lactulose it was possible to synthesise the DTPM-protected lactosamine derivative **9** via the Heyns rearrangement in four steps without chromatography and in overall yields of 65%. The vinylogous amide type protecting group employed allows for easy purification by crystallization, leading to free sugar **9** which is ready for further transformations. Following this approach, the corresponding nigerosamine **16** was prepared from turanose. The entire sequence is simple and relies on cheap and commercially available reagents only. Other glycosylated ketoses reacted accordingly,^a but the observed equilibrium of furanoid and pyranoid conformations as well as side product formation demanded purification by column chromatography.

EXPERIMENTAL

¹H NMR spectra were recorded on a Varian INOVA 500 operating at 499.925 MHz. ¹³C NMR spectra were recorded at 75.47 or 50.29 MHz on a BRUKER MSL 300 or on a Varian Gemini 200. Residual non-deuterated solvent was used as internal standard. Signals of protecting groups were found in the expected regions and are not listed explicitly. Melting points were measured on a Büchi 530 apparatus and are uncorrected. Mass spectra were recorded on an HP 1100 series MSD, Hewlett Packard. Samples were dissolved in acetonitrile or acetonitrile/water mixtures. The scan mode for positive ions (mass range 100–1000 D) was employed varying the fragmentation voltage from 50 to 250 V with best molecular peaks observed at 150 V. Analytical

^aMaltulose and palatinose could be converted accordingly. In case of maltulose, only the monohydrate was commercially available. The presence of water led to lower yields, shifting the equilibrium of the condensation reaction, the first step of the Heyns rearrangement. This was reached at around 60% conversion of maltulose into the corresponding ketosylamine. Palatinose is an excellent substrate for the Heyns reaction, which is driven by the ring expansion from the furanoid fructose to the pyranoid glucosamine derivative. Both glycosylated glucosamine derivatives were found to be mixtures of pyranoid and furanoid tautomers creating the need for column chromatography of the *N*-protected derivatives which did not precipitate well under the conditions employed.



TLC was performed on precoated aluminum plates silica gel 60 F254 (Merck 5554), detected with UV light (254nm), as well as staining with 5% vanillin/sulfuric acid or ceric ammonium molybdate (100g ammonium molybdate/4g cerium sulfate in 1L 10% H₂SO₄) and heating on a hotplate. For column chromatography, silica gel 60 (230–400 mesh, Merck 9385) was employed.

GENERAL METHODS

Heyns rearrangement of glycosylated ketoses with benzylamine: Benzylamine (7.8 equiv) was added at 0°C to the respective ketose. The reaction mixture was allowed to reach room temperature and was subsequently stirred at 40°C until TLC (MeOH/CHCl₃/NH₄OH 2:1:1) showed the ketosylamine as the main product. Methanol (same volume as benzylamine) was added and this mixture was stirred into diethyl ether and kept at 0°C for 3 h. The resulting precipitate consisting of the product and unreacted starting material as well as side products was collected by filtration and dried under reduced pressure.

The crude product was dissolved in methanol/glacial acetic acid (8:1), and stirred at room temperature for 1–2 hours, slowly added to excess ether, and kept at 0°C for 5 h. The precipitate thus obtained containing the glycosylated *N*-benzylglucosamine, unreacted starting material and side products was collected by filtration and dried under reduced pressure.

Hydrogenolysis: The above crude material was dissolved in deionized water, the pH value was brought to 1 by dropwise addition of concd HCL, Pearlman's catalyst [Pd(OH₂)/C, 20%, (5 % by weight)] was added and this reaction mixture was kept on a Parr apparatus at 3 bar until TLC (MeOH/CHCl₃/NH₄OH 2:1:1) showed completed removal of the *N*-benzyl group. The reaction could be performed under ambient pressure but requiring reaction times of up to 5 days.

The catalyst was filtered off, the solution was concentrated under reduced pressure and the residue was treated with benzene/ethanol (1:1) followed by removal of the solvent under reduced pressure. The so obtained yellowish residue contained the glycosylated glucosamine hydrochloride and unreacted starting material.

DTPM protection: The hydrochloride of the free aminosugar was dissolved in a small amount of methanol, triethylamine (2 equiv) and DTPM reagent (1.1 equiv in methanol) were added and the reaction mixture was kept at room temperature for 2 h. The solid product formed was collected by filtration and washed with methanol twice.

***N*-Benzylactosamine (7).** Following the general procedures for the Heyns rearrangement, in a typical experiment lactulose **5** (50 g, 146.1 mmol) was reacted with benzylamine (125mL, 1144.3 mmol) and stirred at 40°C for 3 days, until TLC showed the condensation product as the main component in the mixture. Methanol (125 mL) was added and this solution was slowly stirred into ether (5000 mL). The resulting precipitate was collected by filtration and dried under reduced pressure. A crude product (71.1 g) containing *N*-benzylactulosylamine (**6**) and side products were obtained.



This mixture was dissolved in methanol (250 mL) containing glacial acetic acid (30 mL), kept at room temperature for two h and this solution was stirred into ether (4500 mL). A precipitate containing *N*-benzylactosamine **7**, lactulose **5** and side products (77.9 g in total) was obtained.

Lactosamine hydrochloride (8). The crude material (30.6 g) from the rearrangement reaction was dissolved in deionized water (150 mL), acidified dropwise with concd HCl (7.6 mL) to pH 1, Pd(OH)₂/C (1.5 g) was added and following the general procedures, a mixture of **8** with lactulose **5** could be obtained (26.5 g).

***N*-[1,3-Dimethyl-2,4,6 (1H, 3H, 5H)-trioxopyrimidine-5-ylidene]methyl lactosamine (9).** The mixture obtained after hydrogenolysis (6.8 g) was dissolved in methanol (70 mL), Et₃N (3.7 mL, 26.7 mmol) and DTPM reagent (4.5 g, 21.3 mmol) in methanol (20 mL) were added and the reaction mixture was stirred at ambient temperature for 2 h. The white, amorphous precipitate formed was collected by filtration and washed with methanol twice to give pure **9** (4.6 g, 65% from **5**). Recrystallization from CH₃CN/H₂O gave an analytical sample, mp 265–267°C; [α]_D²⁰ +67.7 (*c* 0.68, DMSO); ¹³C NMR (DMSO-*d*₆) δ 164.64, 162.56, 160.09, 152.01 (DTPM), 104.35 (C-1'), 90.41, 90.29 (C-1, DTPM), 81.03 (C-4'), 76.08 (C-5'), 73.61 (C-3'), 71.08, 70.67, 70.14 (3 C, C-2', C-3, C-5), 68.66 (C-4), 63.88, 60.99, 60.72 (3 C, C-2, C-6, C-6'), 27.83, 27.21 (DTPM). MS: (150V): *m/z* : 508.48 [M⁺-H].

Anal. Calcd for C₁₉H₂₉O₁₃N₃ · ½ H₂O (516.61): C, 44.17; H, 5.86. Found: C, 43.91; H, 5.75.

1,3,6,2',3',4',6'-Hepta-*O*-acetyl-*N*-[1,3-dimethyl-2,4,6 (1H, 3H, 5H)-trioxopyrimidine-5-ylidene]methyl lactosamine (10). To a 10% solution of **9** (10.5 g, 20.7 mmol) in pyridine, acetic anhydride (50 mL, 528.9 mmol) was added dropwise at 0°C, a catalytic amount of dimethylaminopyridine was added and the reaction kept at room temperature for 16 h. The solvent was removed under reduced pressure, the residue dissolved in CH₂Cl₂ and the solution was consecutively washed with 6% HCl and sat aqueous NaHCO₃, then dried over MgSO₄. The crude product was precipitated from diisopropyl ether, collected by filtration and dried over P₂O₅ to give 15.6g (94 %) of **10**. Recrystallization from CHCl₃/cyclohexane gave an analytical sample, mp 145–147°C; [α]_D²⁰ +64.8 (*c* 0.66, CHCl₃); ¹H NMR (CDCl₃) δ 10.16 (dd, 1H, NH DTPM), 8.06 (d, 1H, HC=CDTPM), 6.18 (d, 1 H, J_{1,2} 3.3 Hz, H-1), 5.37 (t, 1 , J_{2,3} = 3,4 9.8 Hz, H-3), 5.33 (d, 1H, H-4'), 5.08 (dd, 1H, H-2'), 4.93 (m, 1 H, J_{2',3'} 10.3 Hz, J_{3',4'} 3.0 Hz, H-3'), 4.46 (d, 1 H, J_{1',2'} 7.9 Hz, H-1'), 4.39 (m, 1 H, J_{6a,6b} 12.1 Hz, H-6a), 4.14 – 4.04 (m, 3H, H-6b, H-6'a, H-6'b), 3.97 (m, 1H, H-5), 3.87 – 3.79 (m, 2H, H-2, H-4), 3.66 – 3.61 (ddd, 1 H, J_{5',6'a} = 4',5' 9.7 Hz, J_{5',6'b} 3.5 Hz, H-5'), 3.2 (2s, 6H, 2 NCH₃); ¹³C NMR (CDCl₃) δ 170.09, 169.98, 169.89, 169.84, 169.38, 168.86, 168.53 (acetyl), 164.45, 162.41, 158.41, 151.68 (DTPM), 100.86 (C-1'), 92.41 (DTPM), 89.82 (C-1), 74.86 (C-4), 70.68, 70.60, 70.52 (3 C, C-2, C-5, C-3'), 70.09 (C-3), 68.89 (C-2'), 66.40 (C-4'), 61.36, 61.26, 60.63 (3 C, C-5', C-6, C-6'), 27.67, 27.05 (DTPM), 20.63, 20.60, 20.51, 20.46, 20.29 (acetyl). MS (150V): *m/z* : 802.7 [M⁺-H];

Anal. Calcd for C₃₃H₄₃O₂₀N₃ (801.8): C, 49.44; H, 5.41. Found: C, 49.27; H, 5.48.

***N*-Benzylnigerosamine (13).** Following the general procedure for the Heyns rearrangement in a typical experiment, turanose **11** (3 g, 8.8 mmol) was reacted with



benzylamine (7.5 mL, 68.7 mmol) and the mixture stirred at 40°C for 4 days, when TLC showed the condensation product as the main component in the mixture. Methanol (7.5 mL) was added and this solution then stirred into ether (400 mL). The precipitate was collected by filtration and dried. A slightly yellow material (3.0 g) containing *N*-benzylturanosyl amine **12** and side products was obtained. This crude material (2.9 g) was dissolved in MeOH/HOAc (14/2 mL), kept at room temperature for 90 min and this solution was stirred into 900 mL of ether. A solid (2.7 g) containing *N*-benzyl-nigerosamine **13**, turanose **11** and side products was collected.

Nigerosamine hydrochloride (14). The rearrangement mixture (4.14 g) were dissolved in 50 mL deionised water, acidified dropwise with concd HCl (0.5 mL), 10% by weight of Pd(OH)₂/C (20 wt.%) was added and, following the general procedures, 3.84 g of a mixture of **14** and turanose **11** were obtained.

1,4,6,2',3',4',6'-Hepta-*O*-acetyl-*N*-(1,3-dimethyl-2,4,6 (1H, 3H, 5H)-trioxopyrimidine-5-ylidene)methyl nigerosamine (16). The above mixture (3.27 g) was dissolved in methanol (30 mL). Et₃N (1.8 mL, 13 mmol) and DTPM reagent (2.7 g, 12.8 mmol) in methanol (5 mL) were added and the reaction mixture was stirred at room temperature for 2 h. For purification and identification, the compounds were per-*O*-acetylated. The residue (7.81 g) was dissolved in pyridine (30 mL, 371 mmol), acetic anhydride (15 mL, 158.5 mmol), and a catalytic amount of DMDP was added, and the reaction was stirred at room temperature for 3 h. Methanol was added, the volume was reduced to 50% under reduced pressure and the resulting solution was diluted with CH₂Cl₂, washed with 6% aqueous HCl, satd NaHCO₃, dried over Na₂SO₄ and concentrated under reduced pressure. Column chromatography employing ethyl acetate/cyclohexane 3:1 gave a mixture of furanoid and pyranoid tautomers of per-*O*-Ac-NHDTM nigerosamine (**16**) in an overall yield of 15% from turanose. Recrystallization from CHCl₃/cyclohexane gave an analytical sample of **16** (pyranose form), $[\alpha]_D^{20} +126.9$ (c 0.63, CHCl₃); ¹H NMR (CDCl₃) δ 10.18 (m, 1H, NHDTM), 8.27 (d, 1 H, J 13.68 Hz, HC=CDTPM), 6.25 (d, 1 H, J_{1,2} 3.4 Hz, H-1), 5.36 (d, 1 H, J_{1',2'} 3.9 Hz, H-1'), 5.26 (m, 1H, H-3'), 5.20 (m, 1H, H-3), 4.98 (m, 1 H, J_{3',4'} 9.8 Hz, H-4'), 4.80 (dd, 1 H, H-2'), 4.26 (m, 1 H, J_{3,4} 10.3 Hz, J_{4,5} 9.3 Hz, H-4), 4.19 (dd, 1 H, J_{5,6a} 4 Hz, J_{6a,6b} 12.2 Hz, H-6a), 4.06 (dd, 1 H, J_{5',6'a} 2.4 Hz, J_{6'a,6'b} 12.7 Hz, H-6'a), 4.01 – 3.29 (m, 3H, H-6b, H-5, H-6'b), 3.83 (ddd, 1 H, J_{2,3} 10.3 Hz, H-2), 3.61 (ddd, 1H, H-5'); ¹³C NMR (CDCl₃) δ 170.90, 170.78, 170.69, 169.75, 169.53, 169.40, 168.66 (acetyl), 165.37, 162.11, 159.74, 152.04 (DTPM), 95.73 (C-1'), 93.01 (DTPM), 90.67 (C-1), 72.94 (C-4), 70.41 (C-5), 70.27, 70.17 (2 C, C-2', C-3'), 69.22 (C-3), 68.21 (C-5'), 67.81 (C-4'), 62.57 (C-2), 61.40 (C-6), 61.15 (C-6'), 28.18, 27.47 (DTPM), 20.99, 20.92, 20.80, 20.76, 20.45 (acetyl). MS (150V): *m/z* : 802.7 [M⁺-H].

Anal. Calcd for C₃₃H₄₃O₂₀N₃ (801.8): C, 49.44; H, 5.41. Found: C, 49.30; H, 5.47.

ACKNOWLEDGMENTS

Glycogroup appreciate financial support by the Austrian *Fonds zur Förderung der Wissenschaftlichen Forschung* (FWF), Vienna (Projects P-13593 CHE and 15726-N03) as well as Hertha-Firnberg-Fellowship T-18 CHE and Erwin Schrödinger-Stipend J2092 to T. M. W. Dr. Günther Gradnig (Fresenius Austria, Linz) is acknowledged for



a generous gift of lactulose. G. D. thanks TU Graz for a guest professorship (March–July 2001).

REFERENCES

1. Fischer, E. Verbindungen des Phenylhydrazins mit den Zuckerarten. *Ber.* **1884**, *17*, 579–584.
2. Fischer, E. Über Isoglucosamin. *Ber.* **1886**, *19*, 1920–1924.
3. Heyns, K.; Koch, W. Formation of an amino sugar from D-fructose and ammonia. *Z. Naturforsch.* **1952**, *7b*, 486–488.
4. Carson, J.F. The reaction of fructose with isopropylamine and cyclohexylamine. *J. Am. Chem. Soc.* **1955**, *77*, 1881–1884.
5. Carson, J.F. The reaction of fructose with aliphatic amines. *J. Am. Chem. Soc.* **1955**, *77*, 5957–5960.
6. Carson, J.F. Reaction of fructose with benzylamine. *J. Am. Chem. Soc.* **1956**, *78*, 3728–3731.
7. Heyns, K.; Meinecke, K.-H. The formation and preparation of D-glucosamine from fructose and ammonia. *Chem. Ber.* **1953**, *86*, 1453–1462.
8. Heyns, K.; Pflughaupt, K.-W.; Müller, D. Ketosylamine rearrangement in the reaction of D-fructose with pyrrolidine. *Chem. Ber.* **1968**, *101*, 2807–2814.
9. Heyns, K.; Paulsen, H.; Breuer, H. Umsetzung von Fructose mit Aminosäuren zu Glucosaminosäuren. *Angew. Chem.* **1956**, *68*, 334–335.
10. Heyns, K.; Breuer, H.; Paulsen, H. Preparation and Behaviour of 2-(N-amino acid substituted) 2-Deoxyglucose (“Glucose Amino Acid”) from glycine, alanine, leucine, and fructose. *Chem. Ber.* **1957**, *90*, 1374–1386.
11. Heyns, K.; Paulsen, H.; Breuer, H. Darstellung und Verhalten weiterer N-substituierter 2-Amino-2-desoxy-aldosen aus D-Fructose und Aminosäuren. *Chem. Ber.* **1958**, *91*, 2750–2762.
12. Heyns, K.; Noack, H. Die Umsetzung von L-Tryptophan und L-Histidin mit Hexosen. *Chem. Ber.* **1964**, *97*, 415–418.
13. Heyns, K.; Eichstedt, R.; Meinecke, K.H. Reaction of fructose and sorbose with ammonia and amines. *Chem. Ber.* **1955**, *88*, 1551–1555.
14. Heyns, K.; Paulsen, H.; Eichstedt, R.; Rolle, M. The formation of 2-aminoaldoses by rearrangement of ketosylamines. *Chem. Ber.* **1957**, *90*, 2039–2049.
15. Heyns, K.; Pflughaupt, K.-W.; Paulsen, H. Ketosylamine rearrangement in the reaction of D-threo-pentulose (D-xylulose) with amines and amino acids to give 2-alkylamino-2-deoxypentoses. *Chem. Ber.* **1968**, *101*, 2800–2806.
16. Heyns, K.; Beilfuß, W. Ketosylamine rearrangement of D-threo-pentulose (“D-Xylulose”) with α -amino acids. *Chem. Ber.* **1970**, *103*, 2873–2876.
17. Wrodnigg, T.M.; Eder, B. The amadori and heyns rearrangements: landmarks in the history of carbohydrate chemistry or unrecognized synthetic opportunities? In *Glycoscience – epimerization, isomerization, rearrangement reactions of carbohydrates*; Stütz, A.E., Ed.; Springer: Heidelberg, 2000; 115–152.
18. Freudenberg, K.; Eichel, H. Über spezifische Kohlenhydrate der Blutgruppen. *Liebigs Ann. Chem.* **1934**, *510*, 240–248.



19. Freudenberg, K.; Eichel, H. Über spezifische Kohlenhydrate der Blutgruppen II. Liebigs Ann. Chem. **1935**, *518*, 97–102.
20. Fukuda, M. Carbohydrate-dependent cell adhesion. Bioorg. Med. Chem. **1995**, *3*, 207–215.
21. Feizi, T.; Childs, R.A. Carbohydrates as antigenic determinants of glycoproteins. Biochem. J. **1987**, *245*, 1–11.
22. Ernst, B.; Dragic, Z.; Marti, S.; Muller, C.; Wagner, B.; Jahnke, W.; Magnani, J.L.; Norman, K.E.; Oehrlein, R.; Peters, T.; Kolb, H.C. Design and synthesis of E-selectin antagonists. Switz. Chimia **2001**, *4*, 268–274.
23. Zhu, T.; Boons, G.-J. A novel and efficient synthesis of a dimeric le^x oligosaccharide on polymeric support. J. Am. Chem. Soc. **2000**, *41*, 10222–10223.
24. Sallas, F.; Nishimura, S.-I. Chemo-enzymatic synthesis of glycopolymers and sequential glycopeptides bearing lactosamine and sialyl Lewis x unit pendant chains. J. Chem. Soc., Perkin Trans. **2000**, *1* (13), 2091–2103.
25. Defrees, S.A.; Gaeta, F.C.A.; Gaudino, J.J.; Zheng, Z.; Hayashi, M. Preparation of Sialyl Lewis X Analogs as Inhibitors of Cellular Adhesion. PCT Int. Appl. WO 9426760, 1994.
26. Phillips, M.L.; Nudelman, E.; Gaeta, F.C.A.; Perez, M.; Singhal, A.K.; Hakomori, S.; Paulson, J.C. ELAM-1 mediates cell adhesion by recognition of a carbohydrate ligand, sialyl- le^x . Science **1990**, *250*, 1130–1132.
27. Brown, R.R.; Nguyen, T.; Lasky, L.A. Characterisation of a human homologue of the murine peripheral lymph node homing receptor. J. Cell. Biol. **1989**, *109* (1), 421–427.
28. Kansas, G.S. Selectins and their ligands: current concepts and controversies. Blood **1996**, *88*, 3259–3287.
29. Varki, A. Biological roles of oligosaccharides: all theories are correct. Glycobiology **1993**, *3* (2), 97–130.
30. Paulsen, H. Synthesis, conformation, and x-ray analysis of saccharide chains of glycoprotein core regions. Angew. Chem. Int. Ed. Engl. **1990**, *29*, 823–938.
31. Toshima, K.; Tatsuta, K. Recent progress in *O*-glycosylation methods and its application to natural product synthesis. Chem. Rev. Ed. Engl. **1993**, *93* (4), 1503–1531.
32. Schmidt, R.R.; Kinzy, W. Anomeric-oxygen activation for glycoside synthesis: the trichloroacetimidate method. Adv. Carbohydr. Chem. Biochem. **1994**, *50*, 21–123.
33. Garegg, P.J. Enzymes in organic synthesis: application to the problems of carbohydrate recognition. Part 2. Adv. Carbohydr. Chem. Biochem. **1997**, *52*, 179–266.
34. Deshpande, P.P.; Kim, H.M.; Zatorski, A.; Park, T.K.; Ragupathi, G.; Livingston, P.O.; Live, D.; Danishefsky, S.J. Strategy in oligosaccharide synthesis: an application to a concise total synthesis of the KH-1 (adenocarcinoma) antigen. J. Am. Chem. Soc. **1998**, *120*, 1600–1614.
35. Sakar, A.K.; Brown, J.R.; Esko, J.D. Synthesis and glycan priming activity of acetylated disaccharides. Carbohydr. Res. **2000**, *329*, 287–300.
36. Wong, C.-H.; Halcomb, R.L.; Ichikawa, Y.; Tetsuya, K. Enzymes in organic synthesis: application to the problem of carbohydrate recognition (part 2). Angew. Chem. **1995**, *107*, 569–593.



37. Wong, C.-H.; Halcomb, R.L.; Ichikawa, Y.; Tetsuya, K. Enzymes in organic synthesis: application to the problem of carbohydrate recognition (part 2). *Angew. Chem. Int. Ed. Engl.* **1995**, *34* (5), 521–546, and references therein.
38. Wong, C.-H. Hydrolysis and formation of glycosidic bonds. In *Enzyme Catalysis in Organic Synthesis*, 2nd Ed.; Drauz, K., Waldmann, H., Eds.; VCH-Wiley: Weinheim, 2002; Vol II, 609–653, chapt 11.3.
39. Kuhn, R.; Kirschenlohr, W. Amino sugar synthesis. IV. The preparation of *N*-acetylactosamine (4- β -Galactopyranosyl-2-deoxy-2-acetamidoglucopyranose) from lactose. *Liebigs Ann. Chem.* **1956**, *600*, 135–143.
40. Lee, R.T.; Lee, Y.C. A Simple preparation of 2-Acetamido-2-deoxy-4-*O*- β -D-galactopyranosyl-D-glucose and D-mannose. *Carbohydr. Res.* **1979**, *77*, 270–274.
41. Alais, J.; Veyrières, A. A convenient synthesis of *N*-acetylactosamine. *Carbohydr. Res.* **1981**, *93* (1), 164–165.
42. Lattová, E.; Petrus, L. Synthesis of *N*-acetylactosamine via ozonolysis of a nitro derivative. *Carbohydr. Res.* **1992**, *235*, 289–293.
43. Kaji, E.; Lichtenthaler, F.W. Expedient conversion of lactose into versatile derivatives of lactosamine and β -D-galactosyl-(1 \rightarrow 4)-D-mannosamine. *J. Carbohydr. Chem.* **1995**, *14* (6), 791–803.
44. Kretzschmar, G.; Stahl, W. Large scale synthesis of linker-modified sialyl Lewis^x, Lewis^x and *N*-acetylactosamine. *Tetrahedron* **1998**, *54*, 6341–6358.
45. Fang, J.; Xie, W.; Li, J.; Wang, P.G. Chemical and enzymatic synthesis of glycoconjugates: synthesis of lactosamine by thermophilic galactosidase catalyzed galactosylation on a multigram scale. *Tetrahedron Lett.* **1998**, *39*, 919–922.
46. Sakai, K.; Katsumi, R.; Ohi, H.; Usui, T.; Ishido, Y. Enzymatic syntheses of *N*-acetylactosamine and *N*-acetylallolactosamine by the use of β -D-galactosidases. *J. Carbohydr. Chem.* **1992**, *11* (5), 553–565.
47. Usui, T.; Kubota, S.; Ohi, H. A convenient synthesis of β -*O*-galactosyl disaccharide derivatives using the β -D-galactosidase from *Bacillus Circulans*. *Carbohydr. Res.* **1993**, *244* (2), 315–323.
48. Hermann, G.F.; Ichikawa, Y.; Wandrey, C.; Gaeta, F.C.A.; Paulson, J.C.; Wong, C.-H. A new multi-enzyme system for a one-pot synthesis of sialyl oligosaccharides: combined use of β -galactosidase and a (2,6)-sialyltransferase coupled with regeneration in situ of CMI-sialic acid. *Tetrahedron Lett.* **1993**, *34* (19), 3091–3094.
49. Kimura, T.; Takayama, S.; Huang, H.; Wong, C.-H. A practical method for the synthesis of *N*-Acetyl-D-lactosamine derivatives by the tandem use of galactose oxidase and β -galactosidase. *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 2348–2350.
50. Hermann, G.F.; Kragl, U.; Wandrey, C. Continuous catalytic synthesis of *N*-acetylactosamine. *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 1342–1343.
51. Yoon, J.H.; Rhee, J.S. The efficient enzymatic synthesis of *N*-acetylactosamine in an organic co-solvent. *Carbohydr. Res.* **2000**, *327* (4), 377–383.
52. Wong, C.-H.; Haynie, S.L.; Whitesides, G.M. Enzymatic-catalyzed synthesis of *N*-acetylactosamine with in situ regeneration of uridine 5'-diphosphate glucose and 5'-diphosphate galactose. *J. Org. Chem.* **1982**, *47*, 5416–5418.
53. Thiem, J.; Wiemann, T. Synthesis of galactose-terminated oligosaccharides by use of galactosyltransferase. *Synthesis* **1992**, 141–145.
54. Ichikawa, Y.; Lin, Y.-C.; Dumas, D.P.; Shen, G.-J.; Garcia-Junceda, E.; Williams,



- M.A.; Bayer, R.; Ketcham, C.; Walker, L.E.; Paulson, J.C.; Wong, C.-H. Chemical-enzymatic synthesis and conformational analysis of sialyl Lewis X and derivatives. *J. Am. Chem. Soc.* **1992**, *114*, 9283–9298.
55. Zervosen, A.; Elling, L. A Novel three-enzyme cycle for the synthesis of *N*-acetyllactosamine with in situ regeneration of uridine 5'-diphosphate glucose and uridine 5'-diphosphate galactose. *J. Am. Chem. Soc.* **1996**, *118*, 1836–1840.
56. Wrodnigg, T.M.; Stütz, A.E. The heyns rearrangement revisited: an exceptionally simple two-step chemical synthesis of D-lactosamine from lactulose. *Angew. Chem. Int. Ed. Engl.* **1999**, *38* (6), 827–828.
57. Dekany, G.; Bornaghi, L.; Papageorgiou, J.; Taylor, S. A Novel amino protecting group: DTPM. *Tetrahedron Lett.* **2001**, *42* (7), 3129–3132.
58. Siddiqui, I.R. The sugars of honey. *Adv. Carbohydr. Chem. Biochem.* **1970**, *25*, 285–309.
59. Hudson, C.S. Melizitose and turanose. *Adv. Carbohydr. Chem.* **1946**, *2*, 1–36.

Received October 11, 2002

Accepted March 21, 2003